## **Original Article**



# Human Papillomavirus Type 16 Based L1, L2, E6, and E7 Peptide Microspheres Induce Encapsulated Peptide Mixture Specific Cytotoxic T Lymphocytes and Tumor Regression in a Murine Model of Cervical Cancer

Md. Asad Khan<sup>1\*</sup>, Kashif Ali<sup>2</sup> and M. Moshahid A. Rizvi<sup>3</sup>

<sup>1</sup>Department of Biochemistry, Faculty of Dentistry, Jamia Millia Islamia, New Delhi, India; <sup>2</sup>Department of Physiology, Faculty of Dentistry, Jamia Millia Islamia, New Delhi, India; <sup>3</sup>Department of Biosciences, Jamia Millia Islamia, New Delhi, India

Received: February 01, 2024 | Revised: June 12, 2024 | Accepted: June 18, 2024 | Published online: June 25, 2024

## Abstract

**Background and objectives:** Infection with HPV16, a high-risk human papillomavirus (HPV), can cause cervical cancer in humans. These infections carry a high risk of morbidity and mortality globally in females. This study aimed to conduct an *in vivo* comparison of Poly (D,L-lactic-co-glycolide) (PLGA)-encapsulated peptide mixture nanoparticles and PLGA microspheres as delivery systems for vaccines.

**Methods:** PLGA polymers were used to form microspheres for a therapeutic vaccine against cervical cancer. The target antigens were the L1 and L2 capsid proteins and the E6 and E7 oncoproteins from HPV16. These antigens were selected based on their immunogenicity, allergenicity, and toxicity. We predicted epitopes for cytotoxic T lymphocytes (CTLs) and helper T lymphocytes. In our investigation of CTL epitopes, we employed synthetic chimeric PLGA microsphere peptides, consisting of multiple H-2Db-restricted HPV16 peptides, coupled with other immune-potentiating adjuvants as predicted by our work.

**Results:** H-2Db-restricted HPV16 peptides, when administered subcutaneously, enabled CTLs to eliminate *in vitro* TC-1 tumor cells expressing E6 and E7 of HPV16. Additionally, *TC-1* cells protected C57BL/6 mice against *in vivo* challenges. To address this problem, peptide-based vaccines, which are among the most effective vaccine systems, have been extensively studied. Combining peptide-based vaccinations with microsphere peptide mixture particles and delivery technologies enhances their efficacy in stimulating cellular immune responses and eliminating tumor cells.

**Conclusions:** This approach may provide a potential therapeutic candidate vaccine based on microsphere-encapsulated peptides for the prevention of cervical cancer caused by HPV.

## Introduction

One type of malignant tumor connected to the cervix is cervical cancer.<sup>1</sup> The human papillomavirus (HPV) is a virus associated

with different types of malignancies and causes deadly cancer. It has been observed in approximately 99.7% of cases of cervical cancer worldwide.<sup>2</sup> Approximately, 91% of deaths from HPV-associated cancer are attributable to cervical cancer. It is the second most common disease among women aged 15 to 44 and the fourth most common malignancy overall.<sup>3,4</sup> Additionally, almost 70% of cervical malignancies are caused by HPV genotypes 16 and 18.<sup>5</sup> E6 and E7 are important viral proteins that play a key role in eliciting oncogenesis in infected cells, leading to uncontrolled cell division and unrestrained telomerase activity, which result in the progression of cervical cancer.<sup>6</sup> In addition to the cervix, HPV can also lead to cancer in the penis, vagina, oropharynx, and vulva.<sup>1,7</sup> Squamous cells typically make up the outer layer of the cervix, while columnar gland cells make

© 2024 The Author(s). This article has been published under the terms of Creative Commons Attribution-Noncommercial 4.0 International License (CC BY-NC 4.0), which permits noncommercial unrestricted use, distribution, and reproduction in any medium, provided that the following statement is provided. "This article has been published in *Cancer Screening and Prevention* at https://doi.org/10.14218/CSP.2024.00003 and can also be viewed on the Journal's website at https://www.xiahepublishing.com/journal/csp".

Keywords: Cervical cancer; Microsphere peptide mixture; Human papillomavirus (HPV); Therapeutic vaccine; Cytotoxic T lymphocyte response; CTL.

<sup>\*</sup>Correspondence to: Md. Asad Khan, Department of Biochemistry, Faculty of Dentistry, Jamia Millia Islamia, New Delhi 110025, India. ORCID: https://orcid. org/0000-0001-6299-3616. Tel: +91-7053342467, E-mail: asad1amu@gmail.com How to cite this article: Khan MA, Ali K, Rizvi MMA. Human Papillomavirus Type 16 Based L1, L2, E6, and E7 Peptide Microspheres Induce Encapsulated Peptide Mixture Specific Cytotoxic T Lymphocytes and Tumor Regression in a Murine Model of Cervical Cancer. *Cancer Screen Prev* 2024;3(2):85–90. doi: 10.14218/CSP. 2024.00003.

up the interior layer. In particular, the first infection may result in dysplasia, called in situ adenocarcinoma. Various methods such as surgery, radiation therapy, and/or chemotherapy have been used for the treatment of cervical cancer.8 Without the risk of serious side effects, vaccination against HPV infection is a promising new alternative therapeutic.<sup>9</sup> Vaccinations use live-attenuated pathogenic antigens generated from pathogenic bacteria or viruses to activate the immune response and induce neutralizing antibodies.<sup>10</sup> The targeted cellular immune responses are generated by novel vaccinations for the treatment of cancer.<sup>11</sup> By employing sequences of immunogenic antigens against HPV-associated proteins, peptidebased vaccines are an excellent option for immunization, as they can effectively eradicate HPV infections and elicit cellular immune responses. Peptide-based vaccines are advantageous due to their ability to elicit cellular immune responses and eradicate HPV infections, making them an ideal option for effective vaccination. Peptide-based vaccines can be quickly obtained through straight forward methods and generally exhibit enhanced stability during storage and transportation, as well as bioavailability.12 The stimulatory effect on the immune system is mediated by microspheres due to their surface-anchored targeting moieties, hydrophobicity, size, and charge.<sup>13</sup> Improved nanoparticles can more successfully transfer associated antigens to antigen-presenting cells (APCs) initiating a robust and effective immune response. In addition, certain cytokines (interleukin (IL)-4, IL-10, tumor necrosis factor (TNF)-B and interferon (IFN)- $\gamma$ ) are secreted during the adaptive immune response. These cytokines activate cellular signaling pathways that regulate immune responses against specific antigens by binding to particular receptors on the surface of immune cells.<sup>14</sup> When phagocytized or endocytosed cells are displayed on APCs, they break the incoming pathogens (or antigens) into tiny peptides. Major Histocompatibility Complex (MHC-I or MHC-II, allowing CD8 or CD4 cells to detect them, subsequently stimulating humoral and/or cellular defense against invasive infections.<sup>11</sup> Targeting intracellular antigens, cancer immune therapies, and vaccinations aim to elicit cellular immune responses. In particular, vaccination candidates must be delivered as endogenous antigens that trigger a cell-mediated immune response when they are displayed by MHC I to CD8+ T-lymphocyte cells after being broken down in the host cytoplasm by the proteasome. HPV-infected cells are cleared by cell-mediated immunity.<sup>15</sup> Adjuvants such as PLGA microspheres were utilized to test the possibility of eliciting immunological responses against HPV16-associated malignancies. The introduction of a mixture of cytotoxic T lymphocyte (CTL) peptides in combination with microsphere adjuvants led to immunogenicity against TC-1 cellinduced tumors in vivo. Generally, APCs phagocytose and transport the peptide combination (L1, L2, E6, and E7) into draining lymph nodes. The usage of adjuvant enables the peptide microspheres to be released continuously over several days or even weeks. We have demonstrated that HPV16 chimeric CTL peptide immunization with an H-2Db-restricted antigen, in conjunction with adjuvant microspheres, generated an immune response specific to peptides and demonstrated in vivo anticancer efficacy against TC-1 cells that expressed HPV16 E6 and E7 in C57BL/6 mice. Before being applied in a clinical setting, these techniques must still be optimized for efficacy, safety, and specificity. We believe that a targeted gene therapy approach will benefit cervical cancer patients as long as the programmable structure and functions are further understood.

#### **Materials and methods**

We designed in silico, chimeric peptides L1165-173 (9 mer), L2108-120

(13 mer), E6<sup>48–57</sup> (10 mer), and E7<sup>48–57</sup> (10 mer) of HPV16, using the algorithmic prediction software CTLpred (available at: http:// www.imtech.res.in/raghava), each containing a CTL epitope. Target epitopes were predicted using a neural network along with the physicochemical properties of the antigen.<sup>16</sup> The small peptides (9–13 mer) were formed into a single large chimeric peptide associated with a linker of two glycines to form {E6<sup>48–57</sup>}GG{E7<sup>48–57</sup>}GG{L1<sup>165–173</sup>}GG{L2<sup>108–120</sup>} (48 mer), E6–E7–L1–L2 containing multiple CTL epitopes. The chimeric peptide swere synthesized commercially by Genscript Corporation, USA (purity 95% by high performance liquid chromatography, dissolved in phosphate-buffered saline (PBS), and stored at –70°C until use.

#### Adjuvants

PLGA was used for microsphere formation and obtained from Boehringer Ingelheim (Ingelheim, Germany).<sup>17</sup> The solvent double evaporation method (water/oil/water) was used to encapsulate the synthetic peptides of HPV16 L1, L2, E6, and E7 in the PLGA. The percentage entrapment of each peptide in the PLGA microsphere was determined by double solvent extraction followed by a BCA protein estimation assay (Pierce).

#### Mice and cell lines

C57BL/6 (H-2b) female mice (5–6 weeks old) were subcutaneously (s.c.) immunized with the encapsulated PLGA peptide microsphere mixture (200 µg/100 µL PBS per mouse). Control mice received the same volumes of PBS & PLGA microsphere. Each group of mice received two booster shots at 10-day intervals. Within 7–10 days after the last booster shot, the mice were euthanized, and their spleens were removed for *in vitro* work. The American Type Culture Collection (USA) provided the tumor cell line *TC-1*, created by retrovirally transducing lung fibroblasts of C57BL/6 origin with the HPV16 E6/E7 virus and the c-Hras *oncogene*. These cells were cultivated in RPMI-1640 medium with 10% (v/v) fetal calf serum, supplemented with nonessential amino acids (2 mM), and G418 (0.4 mg mL<sup>-1</sup>) at 37°C with 5% CO<sub>2</sub>.

#### In vivo tumor regression assay

To achieve *in vivo* tumor regression, viable *TC-1 cells*  $(1 \times 10^5 \text{ cells in } 100 \ \mu\text{L}$  PBS per mouse) were injected s.c. into the left flank of C57BL/6 mice on the first day of the experiment. On day 8, each peptide microsphere was injected s.c. with 200  $\mu$ g of the PLGA microsphere mixture, followed by two booster shots on days 18 and 28. Tumor size was measured every 3–5 days for 35 days using calipers once palpable tumors (7–10 days post-TC-1 inoculation) formed.<sup>18</sup> Tumor volumes greater than 2,000 mm<sup>3</sup> resulted in the death of the mice.

#### Enzyme linked immunosorbant assay (ELISA)

As previously mentioned, ELISA was performed on polystyrene modules with a flat bottom.<sup>19</sup> To ascertain the degree of peptide-specific antibodies in vaccinated mice, 100  $\mu$ L of a peptide microsphere cocktail (containing L1, L2, E6, and E7) was coated with 0.5  $\mu$ g of pure encapsulated PLGA antigens for 16 h at 4°C. The antigens were diluted in carbonate-bicarbonate buffer (0.05 M, pH 9.6). After washing the plates with Tris buffer saline-trixon (TBS-T) (20 mM Tris, 150 mM NaCl, pH 7.4 containing 0.05% Tween-20) and blocking the unoccupied sites with 150  $\mu$ L of 5% fat-free milk in TBS (10 mM Tris, 150 mM NaCl, pH 7.4) for 4–6 h, serum samples (diluted 1:100) were added in triplicate and incu-



**Fig. 1. Tumor challenge and** *in vivo* **tumor growth with TC-1 tumor cells.** Mean tumor volume ± SD (mm<sup>3</sup>) from mice (n = 8) immunized with chimeric peptide microspheres was compared to the respective control mice (injected with PBS along with microspheres) on the indicated days following the tumor challenge. \**P* < 0.05. PBS, phosphate-buffered saline; PLGA, Poly (D,L-lactic-co-glycolide); SD, standard deviation.

bated at  $37^{\circ}$ C for 1 h.<sup>20</sup> The plates were washed four times using TBS-T. To assess the bound antibodies, goat anti-mouse immunoglobulin G (IgG) horseradish peroxidase-conjugated antibody (diluted 1:2,000; Invitrogen) was added and incubated at  $37^{\circ}$ C for 1 h. Following a final wash, the enzyme-substrate TMB (Sigma) was applied, and immunoreactivity was measured using an ELISA plate reader (Biotek) to detect absorbance at 450 nm.

## Lymphocyte proliferation assay

Following *in vitro* stimulation with chimeric encapsulated PLGA peptide microspheres (10 µg per well), lymphocyte proliferation was measured as previously reported.<sup>20</sup> Splenocytes ( $5 \times 10^5$  cells per well) were aliquoted at 100 µL in 96-well plates with RPMI-1640 supplemented with 10% ( $\nu/\nu$ ) Foetal Calf Serum, IL-2 (10 Uml<sup>-1</sup> recombinant-mouse), and 2-mercaptoethanol (0.05 µM). Promega Corp. USA's Cell Titer was utilized to gauge lymphocyte proliferation from various cohorts by (3-(4, 5-dimethylthiazolyl-2)-2, 5-diphenyltetrazolium bromide) (MTT) assay. Cytotoxicity percentage was computed as follows:

#### % Cytotoxicity =

## Cytolytic activity against TC-1 tumor cells

As previously reported,<sup>15</sup> effector-mediated cytolysis of *TC-1 cells* was analyzed using the CytoTox 96 Non-Radioactive test kit (Promega Corp. USA) and a lactate dehydrogenase assay.<sup>21</sup>

## Estimation of cytokines

To conduct a lymphocyte proliferation experiment, 96-well plates containing 10  $\mu$ g of chimeric encapsulated PLGA peptide microspheres were used to cultivate splenocytes from both the control and immunized groups. After 48 h, supernatants were collected for the cytokine test. The amounts of cytokines were measured using Endogen mouse *IFN-y*, *IL-2*, *IL-10*, and *IL-4* ELISA kits (Pierce Biotechnology, Inc. USA).



Fig. 2. ELISA for serum IgG levels of microspheres and peptide mixture encapsulated microsphere wells were coated with the respective antigens. Data are reported as mean  $\pm$  SEM; \**P* < 0.05. ELISA, enzyme linked immunosorbant assay; IgG, immunoglobulin G; PM, PLGA Microsphere; PmM, PLGA encapsulated peptide mixture.

#### Statistical analysis

The Student's *t-test*, Mann-Whitney U test, and Wilcoxon signedrank test were used for statistical analysis. The population-averaged model of the generalized estimating equation was used to examine the tumor regression percentage. *P-values* of  $\leq 0.05$  were considered significant.

## Results

## Regression of tumors against post-TC-1 by Chimeric encapsulated PLGA peptide microsphere

The encapsulated PLGA chimeric peptide microsphere was able to regress E7-related tumors after *s.c.* injection of the *TC-1 cell line* on day 1, for the tumor regression test. Tumor development was observed from the 7th day onward. The encapsulated PLGA chimeric peptide was injected post-tumor assay to assess tumor regression. The *TC-1 cell line* showed tumor progression to 1,895.64 mm<sup>3</sup> but the encapsulated PLGA peptide mixture showed tumor regression to 858.34 mm<sup>3</sup>. The data showed a significant increase in tumor regression compared to controls, being 54.72% for the encapsulated PLGA peptide mixture (Fig. 1).

## Immunoglobulin G responses by subcutaneous immunization of HPV16 encapsulated PLGA peptides (L1, L2, E6 & E7) cocktail

Subcutaneous immunization of the PLGA microsphere peptide cocktail was found to induce antibodies in sera capable of binding to the HPV16 PLGA peptide mixture (L1, L2, E6, and E7) as shown in Figure 2. Serum was collected on day 27 after the first immunization. Encapsulated PLGA peptide-specific IgG antibodies in serum were higher compared to the control PLGA microsphere in mice. The serum IgG level of the PLGA peptide cocktail was significantly higher compared to the IgG level of the control PLGA microsphere.

## Chimeric encapsulated PLGA peptide mixture induced lymphocyte proliferation

Harvesting of splenocytes for *in vitro* stimulation with chimeric peptides of PLGA showed a statistically significant increased proliferation ( $P \le 0.05$ ; Mann–Whitney U test) compared to control microspheres. In response, the lymphocyte proliferation, measured



Fig. 3. *In vitro* lymphocyte proliferative response and antigen-induced cytotoxic T lymphocyte generation in C57BL/6 mice immunized with chimeric peptide. Proliferation of lymphocytes was determined by MTT assay. Data are reported as mean ± SEM; \**P* < 0.05. MTT, 3-(4, 5-dimethylthiazolyl-2)-2, 5-diphenyltetrazolium bromide); PHA, phytohaemagglutinin; PM, PLGA microsphere; PmM, PLGA encapsulated peptide mixture.

at OD<sub>490</sub> by MTT assay, yielded mean values (stimulation indices  $\pm$  SEM) of 0.856  $\pm$  0.104, 0.331  $\pm$  0.082, and 0.819  $\pm$  0.01, respectively, with PLGA peptide mixture and control PLGA microsphere (Fig. 3). When paired analysis (i.e., unstimulated vs. stimulated lymphocytes from the immunized group) was performed, the proliferation level was shown to be considerably ( $P \leq 0.05$ ; Mann–Whitney U test or Wilcoxon signed-rank test).

## Cytotoxicity for peptide mixture against TC-1 cells

After *in vitro* stimulation, viable effector cells from the vaccinated and control groups with the PLGA peptide combination were evaluated for their cytotoxic efficacy against *TC-1 cells*. Figure 4 illustrates the percentage difference in cytotoxicity resulting from effector cell-mediated lysis of *TC-1 cells*. Using the PLGA microsphere peptide mixture, the percentage cytotoxicity was 70.50%, which was significantly greater than the target: effector ratio of unstimulated effector cells from control groups at 1:40.

## Th-1 type response against chimeric peptide by cytokines estimation

In comparison to the control group, stimulated splenocytes exhibited reduced levels of *IL-10* and moderate amounts of *IL-2* and *IL-4*. Additionally, they displayed greater levels of *IFN-y*. Immunized groups had noticeably increased levels of *IFN-y* secretion. A notably elevated release of *IL-2* was seen in the microsphere. PLGA microspheres were used as an adjuvant, and among the Th-2 subtype, significant levels of *IL-10* and *IL-4* were found in the culture supernatant in the PLGA peptide combination (Table 1).

#### Discussion

Peptide antigens can be more specifically and effectively deliv-



Fig. 4. In vitro CTL cytotoxicity of TC-1 tumor cells: CTLs were measured by LDH estimation at 7 days after the last booster. Pooled effector splenocytes of mice immunized with encapsulated peptide mixture and the control group (n = 5 in each group) were co-cultured with target TC-1 cells, and cytolysis of the latter was determined. Data are expressed as mean ± SE of three separate experiments; \*P < 0.05, Wilcoxon rank sum test. CTL, cytotoxic T lymphocyte; LDH, Lactate Dehydrogenase; PM, PLGA Microsphere; PmM, PLGA encapsulated peptide mixture.

ered via microspheres, which also enhance immunogenicity stability and durability. Controlling the biophysical features of microspheres such as size, shape, and surface qualities is relatively easy and can influence the immune responses against the corresponding antigen.<sup>22</sup> One important factor influencing antigen retention and the immunogenicity of peptide-based vaccines is the nanostructure manufacturing method.<sup>23</sup> Generally, microspheres possess immune-stimulating qualities that facilitate antigen delivery and enhance absorption and recognition by antigen-presenting cells.<sup>17</sup> Moreover, antigens prepared or encapsulated in microspheres are resistant to enzymatic breakdown. Characteristics such as longer half-lives and increased permeability through barriers, such as mucosal tissues, are also observed in microspheres within the body.<sup>24</sup> An appealing method to activate the immune response against high-risk HPV varieties associated with infections is therapeutic vaccination.<sup>25</sup> Targeting antigens against HPV that are constitutively expressed in HPV-associated cancers is crucial for therapeutic vaccination,<sup>26</sup> particularly for HPV16 L1, L2, E6, and E7, which play roles in tumor etiology, cellular transformation, and virus replication.<sup>27</sup> The production of peptide vaccines is simple, safe, and restricted to a particular major histocompatibility complex. Initially, we identified unique immunogenic epitopes of HPV16 antigens using software, which overlapped with immunodominant epitopes for L1<sup>165-173</sup>, 28 L2<sup>108-120</sup>, E6<sup>48-57</sup>, 29 and E7<sup>49–57</sup> in murine (H-2Db) CTLs.<sup>30</sup> Combining these epitopes into a single large chimeric peptide enhances antigen absorption and synchronizes the activation of antigen-presenting cells. We also

Table 1. Cytokine estimation of immunized culture supernatant splenocyte and control group

S.No	Groups	Cytokines (pg/mL)			
		IFN-γ	IL-2	IL-4	IL-10
1	Control PM	46.71 ± 0.73	45.31 ± 0.93	72.31 ± 0.002	2.59 ± 0.63
2	PmM	1,234.46 ± 125.30	74.07 ± 53.06	187.34 ± 105.77	12.86 ± 3.13

For n = 6 mice, the data are presented as mean ± SE in pg/mL. *P*-values were determined using the Mann-Whitney U test to compare cytokine levels between the immunized and control groups. *P* < 0.05 was considered significant. IFN, interferon; IL, interleukin; PM, PLGA Microsphere; PmM, PLGA encapsulated peptide mixture.

Khan M.A. et al: HPV based PLGA encapsulated peptide vaccine against cervical cancer

investigated how adjuvants and delivery methods affect these peptide antigens. The E6 and E7 proteins, epitopes of cytotoxic T lymphocytes found in capsid proteins L1 and L2, are known to elicit weakened immune responses in cells. Strong models have been developed to select HPV16 E6 and E7 oncoproteins for creating peptide-based anticancer vaccines.<sup>31</sup> By using cutting-edge adjuvants and delivery methods, the production of CTLs and antibodies induced by peptide immunogens can be significantly improved. Mice were immunized using Freund's complete adjuvant, demonstrating immunogenic potential for peptides Q19D and Q15L from HPV16 E6 and E7 oncoproteins (E643-57 and E744-62, respectively).<sup>32</sup> Subcutaneous vaccination with an H-2Db-restricted HPV16 chimeric PLGA-encapsulated peptide produced peptide-specific CTL-mediated cytolysis for TC-1 tumor cells in vitro. Additionally, C57BL/6 mice were protected against an in vivo challenge by TC-1 cells. With the tested microsphere adjuvant, this peptide effectively induced CTLs that showed good effectiveness. Furthermore, these adjuvants activated splenocytes, triggering peptide-specific lymphocyte proliferation to varying degrees, while showing lower levels of IL-10 and moderate levels of IL-2, IL-4, and IFN-y in the culture supernatant. In vivo testing revealed that both the PLGA microsphere adjuvant and chimerically encapsulated peptide had similar anticancer effectiveness, likely due to the ongoing immune system-tumor cell conflict during the malignant transformation process, where tumors develop defense mechanisms to evade immunosurveillance, particularly with persistence.33 Therefore, lengthy overlapping peptides may be used in future peptide microsphere vaccines to increase the variety of antigenic determinants and decline the barrier for restriction of major histocompatibility complexes. Successful T-cell immune responses have been demonstrated with overlapping peptides in both preclinical animal models and human clinical trials.<sup>34,35</sup>

#### Conclusion

Cervical cancer, caused by HPV, affects millions worldwide, yet there is currently no effective therapeutic vaccine. In this study, we attempted to develop a multi-epitope cervical cancer vaccination. We have developed a peptide-based microsphere candidate vaccination for HPV16-related infections and tumors, showing significant antigen-specific therapeutic effects in a mouse model of cervical cancer. Thus, our preclinical findings provide a solid foundation for future clinical research.

## Acknowledgments

This study was supported by the Department of Biochemistry, Faculty of Dentistry, Jamia Millia Islamia, New Delhi.

## Funding

None.

#### **Conflict of interest**

None.

#### **Author contributions**

Study concept and design (MAK, MAR); acquisition of data (MAK); assay performance and data analysis (MAK, KA, MAR);

drafting of the manuscript (MAK, KA); critical revision of the manuscript (MAR); supervision (MAK). All authors have made a significant contribution to this study and have approved the final manuscript.

## **Ethical statement**

This study was carried out in accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the Indian Council of Medical Research (ICMR) New Delhi, India. The protocol was approved by the Committee on the Ethics of Animal Experiments of the Jawaharlal Nehru Medical College and Hospital, Aligarh Muslim University, Uttar Pradesh, India (AECN: 234/18/JN/AMU). All surgery was performed under sodium pentobarbital anesthesia, and all efforts were made to minimize suffering.

#### **Data sharing statement**

Not available additional data.

#### References

- Crosbie EJ, Einstein MH, Franceschi S, Kitchener HC. Human papillomavirus and cervical cancer. Lancet 2013;382(9895):889–899. doi:10.1016/S0140-6736(13)60022-7, PMID:23618600.
- [2] Walboomers JM, Jacobs MV, Manos MM, Bosch FX, Kummer JA, Shah KV, et al. Human papillomavirus is a necessary cause of invasive cervical cancer worldwide. J Pathol 1999;189(1):12–19. doi:10.1002/ (SICI)1096-9896(199909)189:1<12::AID-PATH431>3.0.CO;2-F, PMID:10451482.
- [3] World Health Organization. WHO guideline for screening and treatment of cervical pre-cancer lesions for cervical cancer prevention: web annex A: syntheses of Mevidence. Geneva: World Health Organization 2021;.
- [4] Khairkhah N, Bolhassani A, Najafipour R. Current and future direction in treatment of HPV-related cervical disease. J Mol Med (Berl) 2022;100(6):829–845. doi:10.1007/s00109-022-02199-y, PMID:354 78255.
- [5] Bruni L, Albero G, Serrano B, Mena M, Collado JJ, Gómez D, et al. ICO/IARC Information Centre on HPV and Cancer (HPV Information Centre). Human Papillomavirus and Related Diseases in the World. Summary Report 10 March 2023.
- [6] Okunade KS. Human papillomavirus and cervical cancer. J Obstet Gynaecol 2020;40(5):602–608. doi:10.1080/01443615.2020.1713592, PMID:31500479.
- [7] Bolhassani A, Mohit E, Rafati S. Different spectra of therapeutic vaccine development against HPV infections. Hum Vaccin 2009;5(10):671–689. doi:10.4161/hv.5.10.9370, PMID:19684468.
- [8] Roden R, Wu TC. Preventative and therapeutic vaccines for cervical cancer. Expert Rev Vaccines 2003;2(4):495–516. doi:10.1586/14760584.2.4.495, PMID:14711335.
- Zepp F. Principles of vaccine design-Lessons from nature. Vaccine 2010;28(Suppl 3):C14–C24. doi:10.1016/j.vaccine.2010.07.020, PMID:20713252.
- [10] Kelly HG, Kent SJ, Wheatley AK. Immunological basis for enhanced immunity of nanoparticle vaccines. Expert Rev Vaccines 2019;18(3):269– 280. doi:10.1080/14760584.2019.1578216, PMID:30707635.
- [11] Zhang J, Fan J, Skwarczynski M, Stephenson RJ, Toth I, Hussein WM. Peptide-Based Nanovaccines in the Treatment of Cervical Cancer: A Review of Recent Advances. Int J Nanomedicine 2022;17:869–900. doi:10.2147/IJN.S269986, PMID:35241913.
- [12] Skwarczynski M, Toth I. Peptide-based synthetic vaccines. Chem Sci 2016;7(2):842–854. doi:10.1039/c5sc03892h, PMID:28791117.
- [13] Al-Halifa S, Gauthier L, Arpin D, Bourgault S, Archambault D. Nanoparticle-Based Vaccines Against Respiratory Viruses. Front Immunol 2019;10:22. doi:10.3389/fimmu.2019.00022, PMID:30733717.

Cancer Screen Prev

- [14] Netea MG, Schlitzer A, Placek K, Joosten LAB, Schultze JL. Innate and Adaptive Immune Memory: an Evolutionary Continuum in the Host's Response to Pathogens. Cell Host Microbe 2019;25(1):13–26. doi:10.1016/j.chom.2018.12.006, PMID:30629914.
- [15] McComb S, Thiriot A, Akache B, Krishnan L, Stark F. Introduction to the Immune System. Methods Mol Biol 2019;2024:1–24. doi:10.1007/978-1-4939-9597-4\_1, PMID:31364040.
- [16] Gomez-Gutierrez JG, Elpek KG, Montes de Oca-Luna R, Shirwan H, Sam Zhou H, McMasters KM. Vaccination with an adenoviral vector expressing calreticulin-human papillomavirus 16 E7 fusion protein eradicates E7 expressing established tumors in mice. Cancer Immunol Immunother 2007;56(7):997–1007. doi:10.1007/s00262-006-0247-2, PMID:17146630.
- [17] Sharma C, Khan MA, Mohan T, Shrinet J, Latha N, Singh N. A synthetic chimeric peptide harboring human papillomavirus 16 cytotoxic T lymphocyte epitopes shows therapeutic potential in a murine model of cervical cancer. Immunol Res 2014;58(1):132–138. doi:10.1007/ s12026-013-8447-2, PMID:24174302.
- [18] Tang A, Dadaglio G, Oberkampf M, Di Carlo S, Peduto L, Laubreton D, Desrues B, Sun CM, Montagutelli X, Leclerc C. B cells promote tumor progression in a mouse model of HPV-mediated cervical cancer. Int J Cancer 2016;139(6):1358–1371. doi:10.1002/ijc.30169, PMID:27130719.
- [19] Khan MA, Alam K, Mehdi SH, Rizvi MMA. Genotoxic effect and antigen binding characteristics of SLE auto-antibodies to peroxynitrite-modified human DNA. Arch Biochem Biophys 2017;635:8–16. doi:10.1016/j.abb.2017.10.008, PMID:29051068.
- [20] Khan MA, Dixit K, Jabeen S, Moinuddin, Alam K. Impact of peroxynitrite modification on structure and immunogenicity of H2A histone. Scand J Immunol 2009;69(2):99–109. doi:10.1111/j.1365-3083.2008.02200.x, PMID:19144075.
- [21] Chai W, Wang X, Wang W, Wang H, Mou W, Gui J. Decreased glycolysis induced dysfunction of NK cells in Henoch-Schonlein purpura patients. BMC Immunol 2020;21(1):53. doi:10.1186/s12865-020-00382-9, PMID:33036556.
- [22] Marasini N, Giddam AK, Khalil ZG, Hussein WM, Capon RJ, Batzloff MR, et al. Double adjuvanting strategy for peptide-based vaccines: trimethyl chitosan nanoparticles for lipopeptide delivery. Nanomedicine (Lond) 2016;11(24):3223–3235. doi:10.2217/nnm-2016-0291, PMID:27830630.
- [23] Demento SL, Siefert AL, Bandyopadhyay A, Sharp FA, Fahmy TM. Pathogen-associated molecular patterns on biomaterials: a paradigm for engineering new vaccines. Trends Biotechnol 2011;29(6):294– 306. doi:10.1016/j.tibtech.2011.02.004, PMID:21459467.
- [24] Zhao G, Chandrudu S, Skwarczynski M, Toth I. The application of self-assembled nanostructures in peptide-based subunit vaccine development. Eur Polym J 2017;93:670–681. doi:10.1016/j.eurpolymj.2017.02.014, PMID:32226094.

- [25] Rai M, Santos C, editors. Nanotechnology Applied To Pharmaceutical Technology. 1st ed. Cham: Springer; 2017.
- [26] Schiller JT, Castellsagué X, Villa LL, Hildesheim A. An update of prophylactic human papillomavirus L1 virus-like particle vaccine clinical trial results. Vaccine 2008;26(Suppl 10):K53–K61. doi:10.1016/j.vaccine.2008.06.002, PMID:18847557.
- [27] Chabeda A, Yanez RJR, Lamprecht R, Meyers AE, Rybicki EP, Hitzeroth II. Therapeutic vaccines for high-risk HPV-associated diseases. Papillomavirus Res 2018;5:46–58. doi:10.1016/j.pvr.2017.12.006, PMID:29277575.
- [28] Ohlschläger P, Osen W, Dell K, Faath S, Garcea RL, Jochmus I, et al. Human papillomavirus type 16 L1 capsomeres induce L1-specific cytotoxic T lymphocytes and tumor regression in C57BL/6 mice. J Virol 2003;77(8):4635–4645. doi:10.1128/jvi.77.8.4635-4645.2003, PMID:12663770.
- [29] Peng S, Ji H, Trimble C, He L, Tsai YC, Yeatermeyer J, et al. Development of a DNA vaccine targeting human papillomavirus type 16 oncoprotein E6. J Virol 2004;78(16):8468–8476. doi:10.1128/ JVI.78.16.8468-8476.2004, PMID:15280455.
- [30] Feltkamp MC, Smits HL, Vierboom MP, Minnaar RP, de Jongh BM, Drijfhout JW, et al. Vaccination with cytotoxic T lymphocyte epitopecontaining peptide protects against a tumor induced by human papillomavirus type 16-transformed cells. Eur J Immunol 1993;23(9):2242– 2249. doi:10.1002/eji.1830230929, PMID:7690326.
- [31] Ressing ME, Sette A, Brandt RM, Ruppert J, Wentworth PA, Hartman M, et al. Human CTL epitopes encoded by human papillomavirus type 16 E6 and E7 identified through in vivo and in vitro immunogenicity studies of HLA-A\*0201-binding peptides. J Immunol 1995;154(11):5934–5943. PMID:7538538.
- [32] Sarkar AK, Tortolero-Luna G, Nehete PN, Arlinghaus RB, Mitchell MF, Sastry KJ. Studies on in vivo induction of cytotoxic T lymphocyte responses by synthetic peptides from E6 and E7 oncoproteins of human papillomavirus type 16. Viral Immunol 1995;8(3):165–174. doi:10.1089/vim.1995.8.165, PMID:8833270.
- [33] Piersma SJ. Immunosuppressive tumor microenvironment in cervical cancer patients. Cancer Microenviron 2011;4(3):361–375. doi:10.1007/s12307-011-0066-7, PMID:21626415.
- [34] Vambutas A, DeVoti J, Nouri M, Drijfhout JW, Lipford GB, Bonagura VR, et al. Therapeutic vaccination with papillomavirus E6 and E7 long peptides results in the control of both established virus-induced lesions and latently infected sites in a pre-clinical cottontail rabbit papillomavirus model. Vaccine 2005;23(45):5271–5280. doi:10.1016/j. vaccine.2005.04.049, PMID:16054734.
- [35] Welters MJ, Kenter GG, Piersma SJ, Vloon AP, Löwik MJ, Berends-van der Meer DM, et al. Induction of tumor-specific CD4+ and CD8+ T-cell immunity in cervical cancer patients by a human papillomavirus type 16 E6 and E7 long peptides vaccine. Clin Cancer Res 2008;14(1):178– 187. doi:10.1158/1078-0432.CCR-07-1880, PMID:18172269.