Opinion



A Data-driven, Dynamic and Flexible Approach to Safely Lifting Mask Mandate: A Proposal



Cynthia Chen and Shiqian Shen^{*}

Department of Anesthesia, Critical Care, and Pain Medicine, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

Received: February 15, 2022 | Revised: February 20, 2022 | Accepted: February 21, 2022 | Published: February 25, 2022

Abstract

The governors of New Jersey, New York, California, Connecticut, Delaware and Oregon announced early in the week of February 7 that selecting mask mandates in their states would end in two to six weeks. These states together account for 77.9 million Americans, or ~23.5% of the U.S. population, and therefore, these changes in policy could have a significant impact on the U.S. economy, as well as education and healthcare systems in each state. As counts of COVID-19 cases, hospitalizations and deaths decrease, mask mandates should be reassessed. We propose that a data-driven, dynamic and flexible approach may help lift mask mandates safely and facilitate a smooth transition to post-pandemic normalcy.

The governors of New Jersey, New York, California, Connecticut, Delaware and Oregon announced early in the week of February 7 that selecting mask mandates in their states would end in 2-6 weeks, with the extent of policy changes varying by state.¹ According to 2020 census data, these six states together account for 77.9 million Americans or ~23.5% of the U.S. population, such that these policies could have a significant impact on the U.S. economy, as well as education and healthcare systems in these states.² Public health officials, however, stated that it was too early to remove mask mandates.3 Strikingly, scientists had also cautioned not to lift mask mandates too early or quickly last May, when the U.S. and other countries began lifting mask mandates. Then, due to the omicron variant, came the largest surge of COV-ID-19 cases, despite increasing numbers of people being vaccinated.^{5,6} In this context, we recommend here a data-driven, dynamic and flexible approach to safely lifting mask mandates (Fig. 1). Briefly, mask mandate policies should be based on assessments of the COVID-19 burden, including COVID-19 metrics such as transmission rate, the impact of COVID-19 on the healthcare system and society, and risks of a resurgence.

According to the Centers for Disease Control and Prevention, there were 108,159 new cases of COVID-19 reported on Feb 18,

Abbreviations: COVID-19, coronavirus disease 2019.

2022, with a seven-day death-rate of 4.1 per 100,000. Currently, nearly all public health policies are based on data, but the challenges lie in assessing the quality of that data and deciding how best to use it. COVID-19 data have the following characteristics that policy makers must be aware of. First, recent case counts of COVID-19 may be underestimated. Indeed, the need for more rigorous collection of COVID-19 case counts by the U.S. government has previously been highlighted.7 COVID-19 case counts were unreliable early in the pandemic. The Biden administration started to distribute free at-home test kits to Americans. Even though this policy has benefits, self-diagnosis at home is likely to make case counts less reliable.8 Although counts of hospitalizations and deaths are more reliable than case counts, many new cases diagnosed or misdiagnosed at home will be unlikely to be reported or collected by public health agencies. Thus, the case count is lower than the true count and should be interpreted with great caution. Second, in order for trends in COVID-19 case counts and reproduction number to be assessed robustly, other monitoring tools must be considered. For example, studies have shown that internet search interests and Farr's law for COVID-19 cases hold predictive value, and these could be considered as additional methods for determining case counts.9,10 Third, high-quality data and expertise in economic and health sciences are required for a sound analysis of the risks and benefits of mask mandates. The effectiveness of mask mandates for preventing and controlling the spread of COVID-19 has been well demonstrated.¹¹⁻¹³ However, there are also well-documented risks and potential harms associated with long-term mask mandates, such as depression, anxiety, and mental health problems.14,15 Other perceived harms, such as physiologic decompensation, seem questionable.¹⁶ Finally, we need more and better data to increase rates of vaccination. Notably, while vacci-

© 2022 The Author(s). This article has been published under the terms of Creative Commons Attribution-Noncommercial 4.0 International License (CC BY-NC 4.0), which permits noncommercial unrestricted use, distribution, and reproduction in any medium, provided that the following statement is provided. "This article has been published in *Exploratory Research and Hypothesis in Medicine* at https://doi.org/10.14218/ERHM.2022.00025 and can also be viewed on the Journal's website at https://www.xiahepublishing.com/journal/erhm".

Keywords: Masking mandates; COVID-19; Policy.

^{*}Correspondence to: Shiqian Shen, Department of Anesthesia, Critical Care, and Pain Medicine, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA. ORCID: https://orcid.org/0000-0003-0062-6831. Tel: +1-617-726-4177, Fax: +1-617-726-2000, E-mail: sshen2@mgh.harvard.edu

How to cite this article: Chen C, Shen S. A Data-driven, Dynamic and Flexible Approach to Safely Lifting Mask Mandate: A Proposal. *Explor Res Hypothesis Med* 2022;7(2):131–133. doi: 10.14218/ERHM.2022.00025.




Fig. 1. A data-driven, dynamic and flexible approach to lifting mask mandates. The mask mandate policy in a given community is based on burden assessment and supported by the majority of stakeholders. Burden assessment consists of three main categories: 1) COVID metrics, including transmission rate, hospitalization rate, intensive care unit admission rate, COVID-related death rate, and vaccination rate; 2) impact of COVID, including workforce shortage, school opening, *etc.*; and 3) risk of COVID resurgence based on transmission rates of new variants and the effectiveness of vaccination. When COVID-related burdens are considered high, a universal masking mandate is appropriate. When COVID-related burdens are considered moderate, masking is required for at-risk populations, including people who are immunocompromised, unvaccinated or have an unknown vaccination status. When COVID-related burdens are considered low, masking is optional.

nation does not completely prevent COVID-19, it can reduce rates of COVID-19-related hospitalization, death, and to a lesser extent, incidence.^{17,18} Research suggests that mask mandates for health-care workers increases vaccination rates and decreases hospital visits for influenza, and it is possible that such mandates may have similar effects for COVID-19.¹⁹ Since vaccination mandates have been the subject of litigation, a reasonable alternative could be to dynamically and gradually lift mask mandates while reinforcing vaccination recommendations.

Cultural preferences, vaccination rates and COVID-19 case counts vary greatly by state and at the local community level in the U.S. Therefore, it will be practical and effective to adopt a dynamic approach to gradually lifting mask mandates. Recently, Rowland et al. proposed a burden-metric based dynamic approach to mask mandates that holistically assesses the impact of masking policies on transmission within schools, student absenteeism and staff capacity.²⁰ Their proposed policy includes masking requirements spanning four levels, namely, making masking universal, making it optional when more than 80% of the eligible population is vaccinated, having masking required for people who are not vaccinated and leaving masking as an optional individual decision.²⁰ Such an approach is practical and metric-based and will be widely acceptable to the public. It is noteworthy that having masking as optional, the lowest level of masking requirement, could greatly reduce discrimination against, and the psychological burden of people who are willing to wear a mask. An optional mask policy may also encourage some people to wear a mask. A similar approach can be considered by policy makers at the state, county and municipal levels. Notably, state and county officials may consider providing municipal officials with greater flexibility to determine local masking policies or mask mandates. Presumably, a masking policy primarily made by local officials should better satisfy the local community's needs, be more likely to be supported by its constituents, and be effectively adopted.

Since COVID-19 cases, hospitalizations and deaths have decreased significantly after the U.S. winter holiday surge, mask mandates should be reassessed.^{21,22} Social distancing policies could also be reassessed if long-term data support safely lifting mask mandates in a given community. Our proposed data-driven, dynamic and flexible approach may help lift mask mandates safely and facilitate a smooth transition to post-pandemic normalcy.

Acknowledgments

The authors acknowledge the department of anesthesia, critical care and pain medicine at Massachusetts General Hospital for support.

Funding

Shiqian Shen received support from the National Institutes of Health (GM128692, NS116423, AG067947, AG070141, AG065606).

Conflict of interest

Dr. Shiqian Shen has been an associate editor of *Exploratory Research and Hypothesis* since March 2021. The authors have no other conflicts of interest to note.

Author contributions

Proposal concept and design (CC, SS), manuscript preparation and revision (CC, SS). All authors have made a significant contribution to this study and have approved the final manuscript.

References

[1] Ferré-Sadurní L, Azi Paybarah A. Covid Updates: New York Is Lifting

Explor Res Hypothesis Med

Mask Mandate, Too. New York Times. 2022. Available from: https:// www.nytimes.com/live/2022/02/08/world/covid-test-vaccine-cases. Accessed February 22, 2022.

- [2] State Population Totals and Components of Change: 2020-2021. 2022 vol: U.S. Census Bureau; 2022. Available from: https://www.census. gov/data/tables/time-series/demo/popest/2020s-state-total.html. Accessed February 22, 2022.
- [3] McCarthy L. While several states lifted mask mandates, public health officials said not so fast: The week in Covid news. New York Times. 2022. Avialable from: https://www.nytimes.com/live/2022/02/12/ world/covid-19-tests-cases-vaccine/while-several-states-lifted-maskmandates-public-health-officials-said-not-so-fast-the-week-in-covidnews. Accessed February 22, 2022.
- Peeples L. What the science says about lifting mask mandates. Nature 2021;593(7860):495–498. doi:10.1038/d41586-021-01394-0, PMID: 34035531.
- Wadman M. New Omicron begins to take over, despite late start. Science 2022;375(6580):480–481. doi:10.1126/science.ada0852, PMID: 35113702.
- [6] Kang YJ. COVID-19 in South Korea: Proper Timing for Easing Mask Mandates After COVID-19 Vaccination. Disaster Med Public Health Prep 2021:1–3. doi:10.1017/dmp.2021.258, PMID:34346306.
- [7] Topol E. It's not too late. Science 2022;375(6578):245. doi:10.1126/science.abo1074, PMID:35040668.
- [8] The White House. Fact Sheet: The Biden Administration to Begin Distributing At-Home, Rapid COVID-19 Tests to Americans for Free. Available from: https://www.whitehouse.gov/briefing-room/statementsreleases/2022/01/14/fact-sheet-the-biden-administration-to-begindistributing-at-home-rapid-covid-19-tests-to-americans-for-free/. Accessed February 22, 2022.
- [9] Yuan X, Xu J, Hussain S, Wang H, Gao N, Zhang L. Trends and Prediction in Daily New Cases and Deaths of COVID-19 in the United States: An Internet Search-Interest Based Model. Explor Res Hypothesis Med 2020;5(2):1–6. doi:10.14218/ERHM.2020.00023, PMID:32348380.
- [10] Xu J, Cheng Y, Yuan X, Li WV, Zhang L. Trends and prediction in daily incidence of novel coronavirus infection in China, Hubei Province and Wuhan City: an application of Farr's law. Am J Transl Res 2020;12(4):1355– 1361. PMID:32355547.
- [11] Xu J, Hussain S, Lu G, Zheng K, Wei S, Bao W, et al. Associations of Stay-at-Home Order and Face-Masking Recommendation with Trends in Daily New Cases and Deaths of Laboratory-Confirmed COVID-19 in the United States. Explor Res Hypothesis Med 2020;5(3):77–86. doi:10.14218/ERHM.2020.00045, PMID:32734759.

- [12] Wong AK, Balzer LB. State-Level Masking Mandates and COVID-19 Outcomes in the United States: A Demonstration of the Causal Roadmap. Epidemiology 2022;33(2):228–236. doi:10.1097/EDE.00000000 00001453, PMID:34907975.
- [13] Tam KM, Walker N, Moreno J. Influence of state reopening policies in COVID-19 mortality. Sci Rep 2022;12(1):1677. doi:10.1038/s41598-022-05286-9, PMID:35102196.
- [14] Daly M, Robinson E. Depression and anxiety during COVID-19. Lancet 2022;399(10324):518. doi:10.1016/S0140-6736(22)00187-8.
- [15] Hafstad GS, Sætren SS, Wentzel-Larsen T, Augusti EM. Adolescents' symptoms of anxiety and depression before and during the Covid-19 outbreak - A prospective population-based study of teenagers in Norway. Lancet Reg Health Eur 2021;5:100093. doi:10.1016/j.lanepe.2021.100093, PMID:34557820.
- [16] Czypionka T, Greenhalgh T, Bassler D, Bryant MB. Masks and Face Coverings for the Lay Public : A Narrative Update. Ann Intern Med 2021;174(4):511–520. doi:10.7326/M20-6625, PMID:33370173.
- [17] Borchering RK, Viboud C, Howerton E, Smith CP, Truelove S, Runge MC, et al. Modeling of Future COVID-19 Cases, Hospitalizations, and Deaths, by Vaccination Rates and Nonpharmaceutical Intervention Scenarios - United States, April-September 2021. MMWR Morb Mortal Wkly Rep 2021;70(19):719–724. doi:10.15585/mmwr.mm7019e3, PMID:33988185.
- [18] Chao CW, Chia-Rong Hsieh V, Tan CY, Yuan MH. Universal masking to prevent SARS-CoV-2 transmission from Taiwan's practices. Glob Health Promot 2021:17579759211033831. doi:10.1177/17579759211033831, PMID:34553622.
- [19] Batabyal RA, Zhou JJ, Howell JD, Alba L, Lee HH, Furuya EY, et al. Impact of New York State Influenza Mandate on Influenza-Like Illness, Acute Respiratory Illness, and Confirmed Influenza in Healthcare Personnel. Infect Control Hosp Epidemiol 2017;38(11):1361–1363. doi:10.1017/ ice.2017.182, PMID:28826427.
- [20] Rowland LC, Klinkhammer MD, Ramirez DWE. Dynamic Masking: A Proposal of Burden-Based Metrics for Masking in K-12 Schools During the COVID-19 Pandemic. J Sch Health 2022;92(1):11–19. doi:10.1111/ josh.13099, PMID:34750833.
- [21] Worldometer. United States's COVID-19 data. Available from: https:// www.worldometers.info/coronavirus/country/us/. Accessed February 22, 2022.
- [22] Mary K. Doctors warn ending school mask mandates will lead to rise in COVID cases as several states lift requirements. Available from: https://abcnews.go.com/Health/doctors-warn-ending-school-maskmandates-lead-rise/story?id=82722570. Accessed February 22, 2022.