



Review Article

The Association between Consumption of Bitter-taste Vegetables in Asian Culture and Metabolic Syndrome Risk Factors in Children: A Narrative Review



Wai Yew Yang^{1*} , Kah Yen Lim¹, Pei Ling Yen¹, Shu Hwa Ong², Nenad Naumovski^{3,4} and Rati Jani⁵

¹Division of Nutrition and Dietetics, School of Health Sciences, Faculty of Medicine and Health, International Medical University, Kuala Lumpur, Malaysia; ²Institute of Research, Development and Innovation, International Medical University, Kuala Lumpur, Malaysia; ³Discipline of Nutrition and Dietetics, School of Rehabilitation and Exercise Sciences, Faculty of Health, University of Canberra, Canberra, ACT, Australia; ⁴Functional Foods and Nutrition Research (FFNR) Laboratory, University of Canberra, Bruce, Ngunnawal Country, ACT, Australia; ⁵School of Health Sciences and Social Work, Griffith University, Gold Coast 4215, Australia

Received: December 22, 2022 | Revised: February 16, 2023 | Accepted: May 06, 2023 | Published online: June 09, 2023

Abstract

Childhood obesity has been escalating in Asian countries in recent decades resulting in the younger age groups being diagnosed with metabolic syndrome (MetS). *Brassicaceae* vegetables that contain high bioactive compounds with anti-inflammatory and anti-oxidative properties might be beneficial in preventing MetS. This narrative review presents (a) the consumption of vegetables in the world population and the availability of bitter-taste vegetables in Asian culture; (b) the interaction between food preference and childhood obesity and (c) potential associations between the consumption of bitter-taste vegetables in Asian culture and clinical outcomes. A number of online searches were conducted for publications in the English language from the year 1990 until October 2022 with a two-step search strategy adopted: initial searches were conducted in four electronic databases (MEDLINE, CINAHL, EMBASE, and Cochrane Library), and a second search using all identified keywords and indexes by including two additional electronic databases (ProQuest and Scopus). The keywords included “bitter”; “vegetables”; “weight status”; “metabolic profile”, “Asia”, “culture”, and “children”. *Brassica* vegetables in Asian countries are abundantly available and commonly consumed, yet the overall vegetable intake in children was inadequate or below the recommended daily intake. Childhood obesity can be influenced by their preference for and consumption of bitter-taste vegetables, and excessive body weight is associated with the risk of developing MetS. It remains inconclusive whether *brassicaceae* vegetables play a dominant role in the group. Future longitudinal studies to investigate the taste sensitivity, vegetable acceptance, and effect of *brassicaceae* vegetables on the risk of MetS in Asian children are warranted.

Introduction

The worldwide rise in the prevalence of childhood obesity over the last decade continues to be a serious health concern, impacting

approximately 340 million children and adolescents between 5 and 9 years old.¹ This was observed in both genders and with similar percentages of prevalence (18% in girls and 19% in boys).² For children below 5 years of age, overweight (weight-for-height >2 standard deviation (SD)) and obesity (weight-for-height >3 SD) cases were reported at 39 million in 2020.¹ In 2016, a global estimate of more than 41 million overweight children below 5 years of age was reported, with the Asian sub-population contributing up to 50%.² These findings were similar in the adolescent group, which recorded the highest obesity rates across developing countries in Asia. Furthermore, there were large variations within the different countries of the Asian continent with the lowest found in rural Bangladesh (3.5%), followed by China (12.5%), Iran, Saudi Arabia (30% respectively), and the Maldives (65%).³

Metabolic syndrome (MetS) is a cluster of cardiovascular risk factors that include abdominal obesity (WC: Waist Circumference

Keywords: Bitter; Vegetables; Weight status; Metabolic profile; Asia; Children.

Abbreviations: AVI, alanine-valine-isoleucine; GSL, glucosinolates; HDL-C, high-density lipoprotein-cholesterol; LDL-C, low-density lipoprotein-cholesterol; MetS, metabolic syndrome; PAV, proline-alanine-valine; PROP, 6-n-propylthiouracil; PTC, phenylthiocarbamide; TG, triglyceride.

***Correspondence to:** Wai Yew Yang, Division of Nutrition and Dietetics, School of Health Sciences, Faculty of Medicine and Health, International Medical University, 57000 Kuala Lumpur, Malaysia. ORCID: <https://orcid.org/0000-0001-6829-8526>. Tel: +60-327317579, Fax: +60-386567229, E-mail: waiyew_yang@imu.edu.my

How to cite this article: Yang WY, Lim KY, Yen PL, Ong SH, Naumovski N, Jani R. The Association between Consumption of Bitter-taste Vegetables in Asian Culture and Metabolic Syndrome Risk Factors in Children: A Narrative Review. *Explor Res Hypothesis Med* 2024;9(1):47–54. doi: 10.14218/ERHM.2022.00129.




Fig. 1. Aetiology of childhood obesity and metabolic syndrome. BP, Blood Pressure; HDL, High-Density Lipoprotein; TG, Triglycerides.

≥90th percentile), elevated blood pressure (SBP: Systolic Blood Pressure ≥130 mmHg or DBP: Diastolic Blood Pressure ≥85 mmHg), impaired fasting blood glucose (FBG ≥ 100 mg/dL), high triglycerides (TG ≥ 150 mg/dL) and low high-density lipoprotein-cholesterol (HDL-C < 40 mg/dL).⁴ Although there are some variations in the diagnostic criteria for MetS in children and adolescents in different countries, there is a strong association between childhood obesity and MetS where there is 11.9% and 29.2% of MetS prevalence among overweight and obese children, respectively (Figure 1).⁵⁻²⁵ These numbers were also higher in Hispanics than their Caucasians and African American equivalents.^{19,26} Almost 90% of obese children and adolescents develop at least one MetS characteristic.²⁷

In South Asia, nearly one-third of urban-dwelling children and adolescents present MetS characteristics,²⁸ with 30% of Asian Indians presenting with insulin resistance.²⁹ Furthermore, a review by Misra *et al.* indicated that MetS was more prevalent among Asian Indian adolescents when fasting hyperinsulinemia was accounted for in its defining criteria, from 0.8% to 4.2% (modified NCEP, ATP III definition).³⁰ An overall MetS prevalence of 2.3% was also recorded in China, which increased with age and peaked at 17 years of age (3.9%).³¹ Dyslipidaemia defined by total cholesterol ≥170–200 mg/dL, low-density lipoprotein cholesterol (LDL-C) ≥110–130 mg/dL, high-density lipoprotein cholesterol (HDL-C) <35–45 mg/dL, TG ≥ 75–100 mg/dL for 0–9 years and ≥90–130 mg/dL for 10–19 years is one of the many components of MetS and had the greatest prevalence among Chinese school-aged children (21%), followed by hypertension (14%),³² and obesity (6.7%).³³ In Taiwan, the prevalence of MetS was higher in boys than girls (5.56% vs 6.39%) in a group of 6 to 12-year-olds.³⁴ In contrast, MetS among Thai adolescents had a prevalence of 4.27%, with the prevalence among females being greater than that of their male counterparts (5.22% vs 3.36%).³⁵

Taste sensitivity differs individually based on age, sex, ethnicity, body weight status, taste bud development, taste concentration,

and saliva composition.³⁶⁻⁴⁰ It is proposed that obese subjects must consume more to compensate for their impaired sensitivity and gain the same stimulation of taste and oral somatosensory system. This lacking sensitivity is hypothesized to have relationships with food intake and body weight variation with implications on long-term health outcomes. However, data concerning correlations between taste sensitivity and obesity are inconsistent⁴¹ and centered mainly on bitter-taste responsiveness, whereas little is currently known about other taste qualities, especially in children.

Sensitivity to bitter taste has broad implications for taste perception, food preferences, and dietary behavior, with potential impacts on nutritional status and health outcomes.⁴² Based on 6-*n*-propylthiouracil (PROP) or phenylthiocarbamide (PTC) sensitivity, individuals can be classified into three PROP taster categories: non-tasters (not at all or taste PROP at a high concentration of 0.32 mmol/l), medium tasters, and PROP super-tasters (perceive extreme bitterness when tasting PROP).⁴³ It has been proposed that those individuals identified as super-tasters are more responsive to other taste qualities including fats⁴⁴ and that PROP tasting is associated with variations in food acceptability, selection of vegetables and fruits, and increased health risk parameters for overweight and obesity.⁴⁵⁻⁴⁷ A recent study amongst 156 Caucasians and 67 Asians aged 18–65 years in the UK reported a higher proportion of Asians were super-tasters as compared to Caucasians (55% vs. 24%, $p < 0.01$), however, evidence is scarce amongst children from Asia.⁴⁸ Our previous pilot study among Malay children aged 7 to 12 years in Kuala Lumpur found no difference between normal and obese children in taste sensitivity and preferences (All p 's > 0.05).⁴⁹ Comparatively, there are more varieties of bitter-taste vegetables used in Asian than in European cuisine, however, their impact on children's weight status and metabolic profile is still relatively limited.^{50,51}

While genetics is an important determinant, it is beyond the scope of this study. Given the lack of existing reviews specifically focusing on bitter-taste vegetables available in Asian culture and

weight status and metabolic profile in children, the aim of the present narrative review was to describe the availability of bitter-taste vegetables in Asian culture and consumption of vegetables in the world population, discuss the potential interaction between food preference on childhood obesity and metabolic syndrome, and establish the potential associations between consumption of bitter-taste vegetables in Asian culture and clinical outcomes.

An online search was conducted for publications in English from 1990 until October 2022 with no restriction on the study design but only included studies conducted in humans. The keywords included “bitter”; “vegetables”; “weight status”; “metabolic profile”; “Asia”; “culture”, and “children”. A two-step search strategy was adopted with initial searches performed in four electronic databases (MEDLINE, CINAHL, EMBASE, and Cochrane Library) followed by an analysis of the text words contained in the title and abstract. A second search using all identified keywords and index terms was then undertaken in two additional electronic databases (ProQuest and Scopus). The systematic search strategy conducted in this review is shown in [Supplementary Table 1](#).

Consumption of vegetables in the world population

Data from the Food and Agriculture Organization of the United Nations regarding global vegetable consumption in 2013 indicated the highest yearly consumption rate per capita in Asia (176.83 kg), followed by Europe (115.10 kg), North America (113.42 kg), Oceania (101.47), Africa (67.57 kg), and the lowest being South America (52.6 kg). Among different vegetable types, the intake of the *Brassica* family seems to mirror that of the global trend with Europe (85 million tonnes) and America (67 million tonnes) following behind Asia’s consumption at 70% or 540 million tonnes. With 355 million tonnes, China was ranked first among the top 10 Brassica-consuming nations in the world, followed by India, the United States, Turkey, the Russian Federation, Japan, Egypt, the Islamic Republic of Iran, Italy, and the Republic of Korea. The population of Greece, however, had the highest per capita consumption (275 kg/person/year), with China coming in a close second (270 kg/person/year).⁵²

Among children aged 6 to 23 months, the UNICEF Infant and Young Child Feeding (IYCF) Global Database showed a 47% and 27% global consumption of vitamin A-rich and other fruits and vegetables, respectively.⁵³ The intake of vitamin A-rich fruits and vegetables, when observed from the Demographic and Health Surveys data, was also 5.7 times more apparent in children from wealthier families as compared to their poorer counterparts.⁵⁴

Despite inadequate fruit and vegetable consumption being common across all geographical regions, it was most apparent amongst South Asians, where 97% and 90% of girls consume fruits and vegetables below the recommended daily servings, respectively.⁵⁵ These numbers vary between countries, ranging from 60% in China,⁵⁶ 85% in India, 75% in Indonesia, 83% in Myanmar, 77% in Sri Lanka, 67% in Thailand, to over 95% in Nepal.^{57,58} Similarly, the Malaysian Adolescent Nutrition Survey 2017 mirrored these findings where a majority of adolescents (10–17 years) did not achieve sufficient vegetable intake (92%).⁵⁹ However, compared to children from Asia, a lower number of European adolescent children from 12 to 17 years old evidenced a daily fruit and vegetable intake lower than the recommended amount at 35% (boys) and 21% (girls).^{60,61}

Availability of bitter-taste vegetables in Asian culture and their nutritional composition

The *Brassicaceae*, often referred to as the *Cruciferae*, is one of

the most commercially significant plant families globally with a diverse worldwide distribution of 372 genera and 4,060 species.⁶² Around the world, various *Brassicaceae* vegetable species are cultivated, which include *Brassica oleracea* (cabbage, broccoli, and cauliflower), *Brassica rapa* (Chinese cabbage, pak choy, choy-sum, and turnips), *Brassica juncea* (mustard greens), and *Raphanus sativus* (daikon radish).⁶²

Among the Asian varieties, *B. oleracea* and *B. rapa* are the most widely grown and consumed. In different countries and regions, the production and consumption of different *Brassicaceae* vegetable types are attributed to social and economic factors. In China, *R. sativus* is the main *Raphanus* species consumed, which mirrors the consumption in Japan, in addition to *B. oleracea*, *B. rapa*, and *B. juncea*.⁶² Within Southeast Asia, the Chinese leafy vegetable kailan and choy-sum are the primary crops grown in varying climates, particularly in Thailand.⁶³ In 2017, the Thailand market supply identified the *Brassicaceae* varieties (*B. juncea*, *B. oleracea*, and *B. rapa*) to possess the greatest diversity in terms of species and cultivars.⁶⁴ Cruciferous vegetables of the *B. oleracea* species *i.e.*, cabbage, broccoli, and cauliflower were among the most common vegetables consumed in the Malaysian diet.^{65,66} These vegetable crop types were also similarly yielded in central Taiwan, together with kailan, Chinese cabbage, and radish.⁶⁷

The consumption of *Brassica* vegetables has been associated with anticarcinogenic, antioxidant, and anti-inflammatory properties and they serve as a source of vitamins, minerals, and several phytochemicals.⁶⁸ These vegetables, particularly broccoli, contain high quantities of carotenoid, tocopherol, vitamin C, and folic acid all of which have been associated with a reduction in the risk of the development of chronic diseases.⁶⁹ In a study by Wills and Ranga,⁷⁰ the analysis of seven leafy Chinese vegetables identified sixteen carotenoids in Chinese cabbage, pak choy, and choy-sum with lutein (20–36%) and β -carotene (16–21%) being the most abundant. Among the vegetable folate sources, broccoli was reported to have the highest levels (110–135 $\mu\text{g}/100\text{g}$).⁷¹ Folate is also present in raw cauliflower (696 \pm 111 $\mu\text{g}/100\text{g}$).⁷²

In addition to being a rich source of vitamins, *Brassica* vegetables are also rich in dietary minerals.⁶⁹ One of the highest mineral sources includes kale with its high contents of potassium (24.2 to 40.8 g/kg), phosphorus (3.2 to 6.4 g/kg), calcium (2.8 g/kg), selenium (1.0 to 1.29 $\mu\text{g}/\text{kg}$), iron (23.5 to 45.7 mg/kg), and zinc (35.8 mg/kg).^{73,74} Several other *Brassicaceae*, such as Chinese cabbage, white cabbage, Brussels sprouts, broccoli, and cauliflower also contain substantial amounts of essential minerals.⁷³ Besides being highly bioavailable in calcium, this plant family possesses high selenium levels, especially when cultivated and grown on selenium-rich soils. In vegetables such as broccoli, selenium is stored as selenocysteine⁶⁹ before being relatively quickly absorbed into the systemic circulation. As a prominent antioxidant, selenium has been associated with a reduction in obesity and the subsequent development of MetS through its influence on adipocyte physiology.⁶

The high antioxidant capacity of *Brassicaceae* is attributed to their high phenolic contents.⁷ Flavonoids, being one of the most common phenolic compounds in these vegetables, play a vital role in maintaining good health via a reduction in the development of reactive oxygen species. Furthermore, the high polyphenolic content has also been associated with the reduction in oxidative stress-related diseases such as various types of cancer, obesity, and cardiovascular disease.⁸ A study by Mian and Mohamed reported that, out of 62 vegetables analyzed, Chinese cabbage, green-white cabbage, kailan, broccoli, and cauliflower are high in flavonoids, ranging from 148–219 mg/kg.⁹

Brassicaceae vegetables have been identified as one of the richest

sources of glucosinolates (GSL) in the human diet. These compounds are secondary plant metabolites responsible for the bitterness and pungency of several different plants such as kailan, kale, cabbage, turnip, broccoli, and cauliflower.⁶² GSL-based research is predominately orientated towards their anticarcinogenic properties. However, relatively recent emerging evidence on the antioxidative impact of GSL-derived isothiocyanates in decreasing the risk of cardiometabolic disorders when incorporating cruciferous vegetables into the diet has garnered special attention.^{12,75} Several potential mechanisms of action have been associated with these compounds: in particular, changes in insulin sensitivity and glycaemic response management, lower blood pressure, and improved endothelial function as well as reduced atherosclerotic plaque development and its progression.¹²

Bitter-taste-related genes and their association with food preference

Differences in taste perception and sensitivity may be explained by genetic variations,^{36–38} as polymorphisms of the genes coding for taste are closely linked to the inter-individual differences.⁴⁰ In humans, bitter taste perception is controlled by the TAS2R family of genes, which has 25 functional genes. Of all these genes, TAS2R38 has been proven to be the PTC/PROP taste receptor accounting for a quarter of the total phenotypic variance in bitter-taste. Three functional single nucleotide polymorphisms in gene nucleotide positions 145 (rs713598 C/G), 785 (rs126866 C/T) and 886 (rs10246939 G/A) encode amino acids at position 49 (alanine/proline, A49P), 262 (valine/alanine, V262A), and 296 (isoleucine/valine, I296V) explaining the differences between the non-taster allele of PTC genes from the taster allele.⁷⁶ The two predominant haplotypes globally are the major taster form [known as Proline-Alanine-Valine (PAV)], and the major non-taster form [Alanine-Valine-Isoleucine (AVI)]. The worldwide variations in non-tasters have been observed in Africa (3%), China (between 6% and 23%), North America (about 30%), and India (as high as 40%).⁷⁶

A systematic review exploring the genetic background of taste perception and preferences and their nutritional implications suggests a significant association exists between TAS2R38 variants (rs713598, rs1726866, rs10246939) and bitter taste preference while fat taste responsiveness is related to rs176667 (CD36).⁷⁷ As the AVI/AVI homozygotes possess lesser bitter sensitivity than heterozygous or homozygous PAV carriers, it has been hypothesized that individuals with increased bitter-taste sensitivity might avoid a wide range of bitter compounds such as coffee, brassica vegetables, and spinach.¹⁹ In the adult studies, higher vegetable consumption was reported amongst the AVI/AVI diplotype compared to those from the PAV/AVI and PAV/PAV diplotype,⁷⁸ especially regarding the intake of brassica vegetables.⁷⁹ However, amongst a sample of preschoolers, PROP taster children demonstrated a greater preference for sweets than non-taster children,⁸⁰ highlighting differences between children and adults in their taste sensitivity driven by both genetic and developmental factors. The interaction between bitterness sensitivity and actual food consumption that can predispose to obesity and other health risks in children has yet to be fully understood and calls for more exploration.

Effect of food preference on childhood obesity and metabolic syndrome

At the nexus between hereditary and environmental components, children's food preference is considered an integral factor in the

development of childhood obesity. Their preference for discretionary foods (sweet-tasting over bitter/sour-natured fruits and vegetables) may be due to the underlying genetic-environmental determinants such as an inherent evolutionary preference for sweet foods, hereditary genotype markers, and bitter-taste endophenotype vs parental feeding practices, food availability, accessibility, and exposure.^{10,11,13,18} In some children, an endophenotype sensitivity towards bitter-taste can become a probable cause of their food rejections, including bitter cruciferous vegetables, with a simultaneous preference for sweeter food choices.^{10,81}

Children and adolescents require adequate energy intake for optimal growth and development, however, excessive energy intake combined with a decrease in activity can lead to the development of obesity.¹⁴ Moreover, the consumption of ultra-processed and refined diets and sugar-sweetened beverages has been purported to be the contributor to rising obesity prevalence. A recent systematic review of the worldwide consumption of ultra-processed foods concluded high variability in the intake with young people, men, and overweight/obese generally having higher levels of consumption.¹⁶ In turn, environmental changes leading to higher caloric consumption have been accompanied by factors predisposing the young to decreased energy expenditure such as reduced physical activity levels and increased time spent in sedentary activities.¹⁷ Moreover, it has been demonstrated that children's eating habits and the risk of childhood obesity were associated with parental feeding practices.^{82–84}

A short-term shift in food type consumption can eventually lead to a chronic increased energy intake, resulting in a subsequent body weight increment over time. Thus, increasing vegetable consumption as a replacement for that of energy-dense foods should be the primary strategy for family food practices and behavioral weight management programs in targeting the obesity epidemic amongst children and adolescents.^{85,86} The fundamentals of this weight control concept stem from reduced caloric consumption through the lower energy density and satiety-enhancing properties of water and fiber within the vegetables.^{87,88}

Potential associations between consumption of bitter-taste vegetables in Asian culture and clinical outcomes

Increased adiposity is closely associated with increased biomarkers of oxidative stress and inflammation.⁸⁹ The phytochemicals found in fruits and vegetables have been shown to have anti-obesity properties due to their possible role in suppressing adiposity-associated metabolic biomarkers.^{90,91} To date, the richest sources of vegetables for potential anti-obesity phytochemicals include the red varieties of onion (*Allium cepa*), lettuce (*Lactuca sativa*), capsicum (*Capsicum annuum*), curly kale (*Brassica oleracea var sabellica*) and orange-fleshed type of sweet potato (*Ipomoea batatas*).²⁰ In contrast to the modern diet, which is strong in wheat, processed meat, and fast food, the traditional Chinese diet, which includes a high intake of rice, vegetables, poultry, pork, and fish, was found to be inversely associated with later obesity.²¹ This study was conducted in China over 5 years from 2006 to 2011 and followed 489 participants aged 6–14 years.

In a recent narrative review of more than 60 prospective cohorts, most included studies found an inverse or no association between intake of various vegetable groups (green leafy vegetables, cruciferous vegetables, allium vegetables, yellow-orange-red vegetables, and legumes) and risk of developing atherosclerotic vascular diseases, heart disease, and stroke.²² However, in the meta-analysis of 23 studies by Gan *et al.*, when compared with the

lowest consumption levels of total fruits and vegetables, the relative risk of coronary heart disease was 0.84 (95% CI: 0.79–0.90), 0.86 (95% CI: 0.82–0.91), 0.87 (95% CI: 0.81–0.93), respectively. Specifically, the inverse associations between fruit and/or vegetable consumption and risk of coronary heart disease were not observed in Asian populations, contrary to Western populations,²³ despite Food and Agriculture Organization of the United Nations reporting the highest consumption of vegetables in Asia compared to other parts of the world. This valuable finding could be partially explained by the cooking methods and overall intake of fruit and vegetables in Asia. Asian cooking primarily involves the boiling and steaming of vegetables, which may cause the loss of water-soluble, heat-sensitive, and oxygen-labile nutrients while higher salt usage during home cooking may also reduce the benefits of vegetables.⁸⁶ More studies are warranted to further investigate the cultural differences in fruit and vegetable intake, and health risks.

The DONALD cohort study compared data on flavonoid intake from vegetables and fruits during adolescence (females: 9–15 years; males: 10–16 years) with fasting blood samples provided in adulthood (18–39 years).²⁴ Results revealed that a higher flavonoid intake from the consumption of fruits and vegetables was related to higher homeostasis model assessment insulin sensitivity (HOMA-2%S) among females ($p = 0.03$) but not males ($p = 0.05$). The HOMA-2%S is a method used to quantify cellular sensitivity to insulin. The authors concluded that flavonoid intake during adolescence is associated with a favorable risk profile for T2DM in early adulthood,²⁴ yet data on the relationship between vegetable consumption as an individual dietary component and MetS remain inconsistent. This discrepancy in results may be caused by variations in the amounts and distinct vegetable subgroups employed in different studies. A cohort study that followed 424 Iranian children and adolescents over 3.6 years reported a negative correlation between overall vegetable consumption and the risk of MetS. Among vegetable subgroups and participants with 1 component of MetS, the consumption of green leafy and allium vegetables was negatively related to the risk of MetS, OR = 0.23, 95% CI [0.07–0.71]; OR = 0.29, 95% CI [0.07–0.71], respectively. The associations were observed after accounting for the major potential confounders such as demographic characteristics (age, gender, physical activity, family history of diabetes), total energy and cholesterol intake, and BMI at baseline.²⁵

Future directions

Vegetable intake, particularly if it is high in bioactive components yet bitter in taste, might be beneficial for preventing MetS, however, these findings can only be attributed to the associations rather than the cause and effect. Thus, future longitudinal studies investigating taste sensitivity, vegetable acceptance, and the effect of brassicas vegetables on the metabolic syndrome risk in Asian children are warranted. In the ethnically diverse population of children in Asia, in-depth research exploring genetics and taste sensitivity is fundamentally important. The body of evidence can be strengthened by investigating children with differing weight status as weight is a complex phenotype and strong determinant predisposing obese children to chronic diseases including MetS.

Conclusion

Evidently, MetS can be prevented through a healthy lifestyle including active living and a balanced diet. There is supportive evidence showing that an overall increased vegetable intake was

negatively associated with the development of MetS in children and adolescents, but the intakes could be influenced by bitter-taste vegetable preference and consumption.

Acknowledgments

None to declare.

Funding

None to declare.

Conflict of interest

One of the co-authors, Nenad Naumovski, is the Associate Editor of *Exploratory Research and Hypothesis in Medicine*. Other authors declare they have no conflict of interests related to this publication.

Author contributions

WYY conceptualized the manuscript, contributed to the methodology, supervision, project administration, and investigation, conducted the formal analysis and data curation, and prepared the original draft of the manuscript. SHO, KYL, and PLY contributed to the investigation, contributed to the methodology, conducted the formal analysis and data curation, and reviewed and edited the manuscript. NN and RJ: conceptualized, reviewed, and edited the manuscript. All authors read and approved the final version of the manuscript.

Data sharing statement

The data used to support the findings of this study are included in the article.

References

- [1] World Health Organization. Obesity and overweight. Available from: <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>. Accessed October 15, 2022.
- [2] NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults. *Lancet* 2017;390(10113):2627–2642. doi:10.1016/S0140-6736(17)32129-3, PMID:29029897.
- [3] Mazidi M, Banach M, Kengne AP, Lipid and Blood Pressure Meta-analysis Collaboration Group. Prevalence of childhood and adolescent overweight and obesity in Asian countries: a systematic review and meta-analysis. *Arch Med Sci* 2018;14(6):1185–1203. doi:10.5114/aoms.2018.79001, PMID:30393474.
- [4] Zimmet P, Alberti KG, Kaufman F, Tajima N, Silink M, Arslanian S, *et al*. The metabolic syndrome in children and adolescents - an IDF consensus report. *Pediatr Diabetes* 2007;8(5):299–306. doi:10.1111/j.1399-5448.2007.00271.x, PMID:17850473.
- [5] Al-Hamad D, Raman V. Metabolic syndrome in children and adolescents. *Transl Pediatr* 2017;6(4):397–407. doi:10.21037/tp.2017.10.02, PMID:29184820.
- [6] Wongdokmai R, Shantavasinkul PC, Chanprasertyothin S, Panpunan P, Matchariyakul D, Sritara P, *et al*. The Involvement of Selenium in Type 2 Diabetes Development Related to Obesity and Low Grade Inflammation. *Diabetes Metab Syndr Obes* 2021;14:1669–1680. doi:10.2147/DMSO.S303146, PMID:33883916.
- [7] Soengas Fernández MD, Sotelo Pérez T, Velasco Pazos P, Cartea

- González ME. Antioxidant properties of Brassica vegetables. *Funct Plant Sci Biotechnol* 2011;5(2):43–55.
- [8] Dias JS. Nutritional quality and effect on disease prevention of vegetables. *Food Nutr Sci* 2019;10(04):369–402. doi:10.4236/fns.2019.104029.
- [9] Miean KH, Mohamed S. Flavonoid (myricetin, quercetin, kaempferol, luteolin, and apigenin) content of edible tropical plants. *J Agric Food Chem* 2001;49(6):3106–3112. doi:10.1021/jf000892m, PMID:11410016.
- [10] Keller KL, Reid A, MacDougall MC, Cassano H, Song JL, Deng L, *et al*. Sex differences in the effects of inherited bitter thiourea sensitivity on body weight in 4–6-year-old children. *Obesity (Silver Spring)* 2010;18(6):1194–1200. doi:10.1038/oby.2009.306, PMID:19779476.
- [11] Jani Mehta R, Mallan KM, Mihrshahi S, Mandalika S, Daniels LA. An exploratory study of associations between Australian-Indian mothers' use of controlling feeding practices, concerns and perceptions of children's weight and children's picky eating: Feeding practices of Australian-Indian mothers. *Nutr Diet* 2014;71(1):28–34. doi:10.1111/1747-0080.12046.
- [12] Connolly EL, Sim M, Travica N, Marx W, Beasy G, Lynch GS, *et al*. Glucosinolates From Cruciferous Vegetables and Their Potential Role in Chronic Disease: Investigating the Preclinical and Clinical Evidence. *Front Pharmacol* 2021;12:767975. doi:10.3389/fphar.2021.767975, PMID:34764875.
- [13] Taylor CM, Wernimont SM, Northstone K, Emmett PM. Picky/fussy eating in children: Review of definitions, assessment, prevalence and dietary intakes. *Appetite* 2015;95:349–359. doi:10.1016/j.appet.2015.07.026, PMID:26232139.
- [14] Aggarwal B, Jain V. Obesity in Children: Definition, Etiology and Approach. *Indian J Pediatr* 2018;85(6):463–471. doi:10.1007/s12098-017-2531-x, PMID:29177599.
- [15] Bowman SA, Gortmaker SL, Ebbeling CB, Pereira MA, Ludwig DS. Effects of fast-food consumption on energy intake and diet quality among children in a national household survey. *Pediatrics* 2004;113(1 Pt 1):112–118. doi:10.1542/peds.113.1.112, PMID:14702458.
- [16] Marino M, Puppo F, Del Bo' C, Vinelli V, Riso P, Porrini M, *et al*. A Systematic Review of Worldwide Consumption of Ultra-Processed Foods: Findings and Criticisms. *Nutrients* 2021;13(8):2778. doi:10.3390/nu13082778, PMID:34444936.
- [17] Biddle SJ, Garcia Bengoechea E, Wiesner G. Sedentary behaviour and adiposity in youth: a systematic review of reviews and analysis of causality. *Int J Behav Nutr Phys Act* 2017;14(1):43. doi:10.1186/s12966-017-0497-8, PMID:28351363.
- [18] Dubois L, Diasparra M, Bédard B, Kaprio J, Fontaine-Bisson B, Tremblay R, *et al*. Genetic and environmental influences on eating behaviors in 2.5- and 9-year-old children: a longitudinal twin study. *Int J Behav Nutr Phys Act* 2013;10:134. doi:10.1186/1479-5868-10-134, PMID:24313977.
- [19] Wittcopp C, Conroy R. Metabolic Syndrome in Children and Adolescents. *Pediatr Rev* 2016;37(5):193–202. doi:10.1542/pir.2014-0095, PMID:27139327.
- [20] Williams DJ, Edwards D, Hamernig I, Jian L, James AP, Johnson SK, *et al*. Vegetables containing phytochemicals with potential anti-obesity properties: A review. *Food Res Int* 2013;52(1):323–333. doi:10.1016/j.foodres.2013.03.015.
- [21] Zhen S, Ma Y, Zhao S, Yang X, Wen D. Dietary pattern is associated with obesity in Chinese children and adolescents: data from China Health and Nutrition Survey (CHNS). *Nutr J* 2018;17(1):68. doi:10.1186/s12937-018-0372-8, PMID:29996840.
- [22] Blekkenhorst LC, Sim M, Bondonno CP, Bondonno NP, Ward NC, Prince RL, *et al*. Cardiovascular Health Benefits of Specific Vegetable Types: A Narrative Review. *Nutrients* 2018;10(5):595. doi:10.3390/nu10050595, PMID:29751617.
- [23] Gan Y, Tong X, Li L, Cao S, Yin X, Gao C, *et al*. Consumption of fruit and vegetable and risk of coronary heart disease: a meta-analysis of prospective cohort studies. *Int J Cardiol* 2015;183:129–137. doi:10.1016/j.ijcard.2015.01.077, PMID:25662075.
- [24] Penczynski KJ, Herder C, Krupp D, Rienks J, Egert S, Wudy SA, *et al*. Flavonoid intake from fruit and vegetables during adolescence is prospectively associated with a favourable risk factor profile for type 2 diabetes in early adulthood. *Eur J Nutr* 2019;58(3):1159–1172. doi:10.1007/s00394-018-1631-3, PMID:29468461.
- [25] Hosseinpour-Niazi S, Bakhshi B, Betru E, Mirmiran P, Darand M, Azizi F. Prospective study of total and various types of vegetables and the risk of metabolic syndrome among children and adolescents. *World J Diabetes* 2019;10(6):362–375. doi:10.4239/wjdv.10.i6.362, PMID:31231459.
- [26] Silveira LS. Metabolic syndrome: Criteria for diagnosing in children and adolescents. *Endocrinol Metab Syndr* 2013;2(3):1–6. doi:10.4172/2161-1017.1000118.
- [27] Cook S, Weitzman M, Auinger P, Nguyen M, Dietz WH. Prevalence of a metabolic syndrome phenotype in adolescents: findings from the third National Health and Nutrition Examination Survey, 1988–1994. *Arch Pediatr Adolesc Med* 2003;157(8):821–827. doi:10.1001/archpedi.157.8.821, PMID:12912790.
- [28] Misra A, Khurana L. The metabolic syndrome in South Asians: epidemiology, determinants, and prevention. *Metab Syndr Relat Disord* 2009;7(6):497–514. doi:10.1089/met.2009.0024, PMID:19900153.
- [29] Misra A, Misra R, Wijesuriya M, Banerjee D. The metabolic syndrome in South Asians: continuing escalation & possible solutions. *Indian J Med Res* 2007;125(3):345–354. PMID:17496360.
- [30] Misra A, Khurana L, Vikram NK, Goel A, Wasir JS. Metabolic syndrome in children: current issues and South Asian perspective. *Nutrition* 2007;23(11-12):895–910. doi:10.1016/j.nut.2007.08.018, PMID:17936199.
- [31] Zhu Y, Zheng H, Zou Z, Jing J, Ma Y, Wang H, *et al*. Metabolic Syndrome and Related Factors in Chinese Children and Adolescents: Analysis from a Chinese National Study. *J Atheroscler Thromb* 2020;27(6):534–544. doi:10.5551/jat.50591, PMID:31827007.
- [32] Piernas C, Wang D, Du S, Zhang B, Wang Z, Su C, *et al*. Obesity, non-communicable disease (NCD) risk factors and dietary factors among Chinese school-aged children. *Asia Pac J Clin Nutr* 2016;25(4):826–840. doi:10.6133/apjcn.092015.37, PMID:27702726.
- [33] Wei X, Peng R, Cao J, Kang Y, Qu P, Liu Y, *et al*. Serum vitamin A status is associated with obesity and the metabolic syndrome among school-age children in Chongqing, China. *Asia Pac J Clin Nutr* 2016;25(3):563–570. doi:10.6133/apjcn.092015.03, PMID:27440692.
- [34] Lee MS, Wahlqvist ML, Yu HL, Pan WH. Hyperuricemia and metabolic syndrome in Taiwanese children. *Asia Pac J Clin Nutr* 2007;16(Suppl 2):594–600. PMID:17724000.
- [35] Siwarom S, Pirojsakul K, Aekplakorn W, Paksi W, Kessomboon P, Nee-lapaichit N, *et al*. Waist-to-Height Ratio Is a Good Predictor of Metabolic Syndrome in Adolescents: A Report From the Thai National Health Examination Survey V, 2014. *Asia Pac J Public Health* 2022;34(1):36–43. doi:10.1177/10105395211046474, PMID:34590882.
- [36] Rodrigues L, Espanca R, Costa AR, Antunes CM, Pomar C, Capela-Silva F, *et al*. Comparison of salivary proteome of children with different sensitivities for bitter and sweet tastes: association with body mass index. *Int J Obes (Lond)* 2019;43(4):701–712. doi:10.1038/s41366-018-0289-5, PMID:30568269.
- [37] Lim J, Pullicin AJ. Oral carbohydrate sensing: Beyond sweet taste. *Physiol Behav* 2019;202:14–25. doi:10.1016/j.physbeh.2019.01.021, PMID:30684587.
- [38] Deshaware S, Singhal R. Genetic variation in bitter taste receptor gene TAS2R38, PROP taster status and their association with body mass index and food preferences in Indian population. *Gene* 2017;627:363–368. doi:10.1016/j.gene.2017.06.047, PMID:28652185.
- [39] Forestell CA, Mennella JA. The Ontogeny of Taste Perception and Preference Throughout Childhood. In: Doty RL (ed). *Handbook of Olfaction and Gustation*. Hoboken, USA: John Wiley & Sons, Inc; 2015:795–828. doi:10.1002/9781118971758.ch36.
- [40] Vignini A, Borroni F, Sabbatinelli J, Pugnali S, Alia S, Taus M, *et al*. General Decrease of Taste Sensitivity Is Related to Increase of BMI: A Simple Method to Monitor Eating Behavior. *Dis Markers* 2019;2019:2978026. doi:10.1155/2019/2978026, PMID:31089392.
- [41] Cox DN, Hendrie GA, Carty D. Sensitivity, hedonics and preferences for basic tastes and fat amongst adults and children of differing weight status: A comprehensive review. *Food Qual Prefer* 2016;48:359–367. doi:10.1016/j.foodqual.2015.01.006.
- [42] Lanfer A, Bammann K, Knof K, Buchecker K, Russo P, Veidebaum T, *et al*. Predictors and correlates of taste preferences in European children: The IDEFICS study. *Food Qual Prefer* 2013;27(2):128–136.

- doi:10.1016/j.foodqual.2012.09.006.
- [43] Laureati M, Pagliarini E, Toschi TG, Monteleone E. Research challenges and methods to study food preferences in school-aged children: A review of the last 15 years. *Food Qual Prefer* 2015;46:92–102. doi:10.1016/j.foodqual.2015.07.010.
- [44] Weaver MR, Brittin HC. Food preferences of men and women by sensory evaluation versus questionnaire. *Fam Consum Sci Res J* 2001;29(3):288–301. doi:10.1177/1077727X01293006.
- [45] Davis C, Patte K, Levitan R, Reid C, Tweed S, Curtis C. From motivation to behaviour: a model of reward sensitivity, overeating, and food preferences in the risk profile for obesity. *Appetite* 2007;48(1):12–19. doi:10.1016/j.appet.2006.05.016, PMID:16875757.
- [46] Deglaire A, Méjean C, Castetbon K, Kesse-Guyot E, Hercberg S, Schlich P. Associations between weight status and liking scores for sweet, salt and fat according to the gender in adults (The Nutrinet-Santé study). *Eur J Clin Nutr* 2015;69(1):40–46. doi:10.1038/ejcn.2014.139, PMID:25074389.
- [47] Sia B, Low S, Foong W, Pramasivah M, Say YH. Demographic differences of preference, intake frequency and craving hedonic ratings of sweet foods among Malaysian subjects in Kuala Lumpur. *Malays J Med Health Sci* 2013;9(1):55–64.
- [48] Hill JO, Wyatt HR, Peters JC. Energy balance and obesity. *Circulation* 2012;126(1):126–132. doi:10.1161/CIRCULATIONAHA.111.087213, PMID:22753534.
- [49] Lim LS, Tang XH, Yang WY, Ong SH, Naumovski N, Jani R. Taste Sensitivity and Taste Preference among Malay Children Aged 7 to 12 Years in Kuala Lumpur-A Pilot Study. *Pediatr Rep* 2021;13(2):245–256. doi:10.3390/pediatric13020034, PMID:34069893.
- [50] Tepper BJ. Nutritional implications of genetic taste variation: the role of PROP sensitivity and other taste phenotypes. *Annu Rev Nutr* 2008;28:367–388. doi:10.1146/annurev.nutr.28.061807.155458, PMID:18407743.
- [51] Keller KL, Adise S. Variation in the Ability to Taste Bitter Thiourea Compounds: Implications for Food Acceptance, Dietary Intake, and Obesity Risk in Children. *Annu Rev Nutr* 2016;36:157–182. doi:10.1146/annurev-nutr-071715-050916, PMID:27070900.
- [52] Cartea ME, Lema M, Francisco M. Basic Information on Vegetable Brassica Crops. In: Sadowski J, Kole C (eds). *Genetics, Genomics and Breeding of Vegetable Brassicas*. Boca Raton: CRC Press; 2011:1–34. Available from: https://www.researchgate.net/publication/235679974_Basic_information_on_vegetables_Brassica_crops.
- [53] United Nations Children's Fund (UNICEF) (eds). *Fed to Fail? The Crisis of Children's Diets in Early Life*. 2021 Child Nutrition Report. New York: UNICEF; 2021.
- [54] Choudhury S, Headey DD, Masters WA. First foods: Diet quality among infants aged 6–23 months in 42 countries. *Food Policy* 2019;88:101762. doi:10.1016/j.foodpol.2019.101762, PMID:31853163.
- [55] Keats EC, Rappaport AI, Shah S, Oh C, Jain R, Bhutta ZA. The Dietary Intake and Practices of Adolescent Girls in Low- and Middle-Income Countries: A Systematic Review. *Nutrients* 2018;10(12):1978. doi:10.3390/nu10121978, PMID:30558128.
- [56] Cheng G, Duan R, Kranz S, Libuda L, Zhang L. Development of a Dietary Index to Assess Overall Diet Quality for Chinese School-Aged Children: The Chinese Children Dietary Index. *J Acad Nutr Diet* 2016;116(4):608–617. doi:10.1016/j.jand.2015.11.010, PMID:26825477.
- [57] Peltzer K, Pengpid S. Fruits and vegetables consumption and associated factors among in-school adolescents in five Southeast Asian countries. *Int J Environ Res Public Health* 2012;9(10):3575–3587. doi:10.3390/ijerph9103575, PMID:23202763.
- [58] Dhungana RR, Bista B, Pandey AR, de Courten M. Prevalence, clustering and sociodemographic distributions of non-communicable disease risk factors in Nepalese adolescents: secondary analysis of a nationwide school survey. *BMJ Open* 2019;9(5):e028263. doi:10.1136/bmjopen-2018-028263, PMID:31110109.
- [59] Institute of Public Health. *National Health and Morbidity Survey (NHMS) 2017: Key Findings from the Adolescent Health and Nutrition Surveys*. Kuala Lumpur (MY): National Institutes of Health, Ministry of Health Malaysia; 2018.
- [60] World Health Organization. *WHO European Childhood Obesity Surveillance Initiative (COSI) Report on the fourth round of data collection, 2015–2017*. Geneva: World Health Organization; 2021.
- [61] Diethelm K, Jankovic N, Moreno LA, Huybrechts I, De Henauw S, De Vriendt T, *et al.* Food intake of European adolescents in the light of different food-based dietary guidelines: results of the HELENA (Healthy Lifestyle in Europe by Nutrition in Adolescence) Study. *Public Health Nutr* 2012;15(3):386–398. doi:10.1017/S1368980011001935, PMID:21936969.
- [62] Zhu B, Liang Z, Zang Y, Zhu Z, Yang J. Diversity of glucosinolates among common brassicaceae vegetables in China. *Hortic Plant J* 2022. doi:10.1016/j.hpj.2022.08.006.
- [63] Issarakraisila M, Ma Q, Turner DW. Photosynthetic and growth responses of juvenile Chinese kale (*Brassica oleracea* var. *alboglabra*) and Caisin (*Brassica rapa* subsp. *parachinensis*) to waterlogging and water deficit. *Sci Hortic* 2007;111(2):107–113. doi:10.1016/j.scienta.2006.10.017.
- [64] Turreira-García N, Vilkkamaa AM, Byg A, Theilade I. Diversity, Knowledge, and use of leafy vegetables in Northern Thailand-maintenance and transmission of ethnobotanical knowledge during urbanisation. *Nat Hist Bull Siam Soc* 2017;62(1):85–105.
- [65] National Coordinating Committee on Food and Nutrition. *Malaysian Dietary Guidelines for Children and Adolescents*. Putrajaya: Ministry of Health Malaysia; 2013.
- [66] Izzah AN, Aminah A, Pauzi AM, Lee YH, Wan Rozita WM, *et al.* Patterns of fruits and vegetable consumption among adults of different ethnics in Selangor, Malaysia. *Int Food Res J* 2012;19(3):1095–1107.
- [67] Ali M, Wu SN, Wu MH. Evaluation of the net nutritive gain of policy interventions: An application to Taiwan household survey data. *Taiwan: Asian Vegetable Research and Development Center*; 2000.
- [68] Dias JC da S. Nutritional and Health Benefits of Brassica Vegetables and Exploitation of Innovative Markets Concepts for their Dietary Diversity. In: Chittoria RK (ed). *Emerging Trends in Disease and Health Research*. Book Publisher International; 2022:53–73.
- [69] Dias JC da S. Minerals and Nutraceuticals in Vegetables and Their Human Health Benefits. In: Tarro G, Sharma DC (eds). *Highlights on Medicine and Medical Science*. Book Publisher International; 2021:27–71. doi:10.9734/bpi/hmms/v12/10215D.
- [70] Wills RBH, Ranga A. Determination of carotenoids in Chinese vegetables. *Food Chem* 1996;56(4):451–455. doi:10.1016/0308-8146(95)00226-X.
- [71] Konings EJ, Roomans HH, Dorant E, Goldbohm RA, Saris WH, van den Brandt PA. Folate intake of the Dutch population according to newly established liquid chromatography data for foods. *Am J Clin Nutr* 2001;73(4):765–776. doi:10.1093/ajcn/73.4.765, PMID:11273852.
- [72] Melse-Boonstra A, Verhoeve P, Konings EJ, Van Dusseldorp M, Matser A, Hollman PC, *et al.* Influence of processing on total, monoglutamate and polyglutamate folate contents of leeks, cauliflower, and green beans. *J Agric Food Chem* 2002;50(12):3473–3478. doi:10.1021/jf0112318, PMID:12033813.
- [73] Puupponen-Pimiä R, Häkkinen ST, Aarni M, Suortti T, Lampi AM, Eurla M, *et al.* Blanching and long-term freezing affect various bioactive compounds of vegetables in different ways: Effect of blanching and freezing on vegetables. *J Sci Food Agric* 2003;83(14):1389–1402. doi:10.1002/jsfa.1589.
- [74] Hedges LJ, Lister CE. *Nutritional attributes of Brassica vegetables*. New Zealand: New Zealand Institute for Crop & Food Research; 2006:46.
- [75] Kamal RM, Abdull Razis AF, Mohd Sukri NS, Perimal EK, Ahmad H, Patrick R, *et al.* Beneficial Health Effects of Glucosinolates-Derived Isothiocyanates on Cardiovascular and Neurodegenerative Diseases. *Molecules* 2022;27(3):624. doi:10.3390/molecules27030624, PMID:35163897.
- [76] Drayna D. Human taste genetics. *Annu Rev Genomics Hum Genet* 2005;6:217–235. doi:10.1146/annurev.genom.6.080604.162340, PMID:16124860.
- [77] Garcia-Bailo B, Toguri C, Eny KM, El-Sohemy A. Genetic variation in taste and its influence on food selection. *OMICS* 2009;13(1):69–80. doi:10.1089/omi.2008.0031, PMID:18687042.
- [78] Duffy VB, Hayes JE, Davidson AC, Kidd JR, Kidd KK, Bartoshuk LM. Vegetable Intake in College-Aged Adults Is Explained by Oral Sensory Phenotypes and TAS2R38 Genotype. *Chemosens Percept* 2010;3(3-4):137–148. doi:10.1007/s12078-010-9079-8, PMID:21157576.
- [79] Sacerdote C, Guarrera S, Smith GD, Grioni S, Krogh V, Masala G, *et*

- al*. Lactase persistence and bitter taste response: instrumental variables and mendelian randomization in epidemiologic studies of dietary factors and cancer risk. *Am J Epidemiol* 2007;166(5):576–581. doi:10.1093/aje/kwm113, PMID:17596267.
- [80] Keller KL, Olsen A, Cravener TL, Bloom R, Chung WK, Deng L, *et al*. Bitter taste phenotype and body weight predict children's selection of sweet and savory foods at a palatable test-meal. *Appetite* 2014;77:113–121. doi:10.1016/j.appet.2014.02.019, PMID:24607656.
- [81] Gibson EL, Cooke L. Understanding Food Fussiness and Its Implications for Food Choice, Health, Weight and Interventions in Young Children: The Impact of Professor Jane Wardle. *Curr Obes Rep* 2017;6(1):46–56. doi:10.1007/s13679-017-0248-9, PMID:28205158.
- [82] Naidu BM, Mahmud SZ, Ambak R, Sallehuddin SM, Mutalip HA, Saari R, *et al*. Overweight among primary school-age children in Malaysia. *Asia Pac J Clin Nutr* 2013;22(3):408–415. doi:10.6133/apjcn.2013.22.3.18, PMID:23945411.
- [83] Baharudin A, Man CS, Ahmad MH, Wong NI, Salleh R, Megat Radzi MR, *et al*. Associated Factors to Prevalence of Childhood under Nutrition in Malaysia: Findings from the National Health and Morbidity Survey (NHMS 2016). *Health Sci J* 2019;13(1):627. doi:10.21767/1791-809X.1000627.
- [84] Scaglioni S, Arrizza C, Vecchi F, Tedeschi S. Determinants of children's eating behavior. *Am J Clin Nutr* 2011;94(6 Suppl):2006S–2011S. doi:10.3945/ajcn.110.001685, PMID:22089441.
- [85] Epstein LH, Gordy CC, Raynor HA, Beddome M, Kilanowski CK, Paluch R. Increasing fruit and vegetable intake and decreasing fat and sugar intake in families at risk for childhood obesity. *Obes Res* 2001;9(3):171–178. doi:10.1038/oby.2001.18, PMID:11323442.
- [86] Epstein LH, Paluch RA, Beecher MD, Roemmich JN. Increasing healthy eating vs. reducing high energy-dense foods to treat pediatric obesity. *Obesity (Silver Spring)* 2008;16(2):318–326. doi:10.1038/oby.2007.61, PMID:18239639.
- [87] Grunwald GK, Seagle HM, Peters JC, Hill JO. Quantifying and separating the effects of macronutrient composition and non-macronutrients on energy density. *Br J Nutr* 2001;86(2):265–276. doi:10.1079/bjn2001404, PMID:11502241.
- [88] Rolls BJ, Ello-Martin JA, Tohill BC. What can intervention studies tell us about the relationship between fruit and vegetable consumption and weight management? *Nutr Rev* 2004;62(1):1–17. doi:10.1111/j.1753-4887.2004.tb00001.x, PMID:14995052.
- [89] Marseglia L, Manti S, D'Angelo G, Nicotera A, Parisi E, Di Rosa G, *et al*. Oxidative stress in obesity: a critical component in human diseases. *Int J Mol Sci* 2014;16(1):378–400. doi:10.3390/ijms16010378, PMID:25548896.
- [90] González-Castejón M, Rodríguez-Casado A. Dietary phytochemicals and their potential effects on obesity: a review. *Pharmacol Res* 2011;64(5):438–455. doi:10.1016/j.phrs.2011.07.004, PMID:21798349.
- [91] Yeon JY, Kim HS, Sung MK. Diets rich in fruits and vegetables suppress blood biomarkers of metabolic stress in overweight women. *Prev Med* 2012;54(Suppl):S109–S115. doi:10.1016/j.ypmed.2011.12.026, PMID:22226662.