



Article information: <http://doi.org/10.14218/JCTH.2022.00312>

Highlights

1. Sarmentosin stimulated caspase-dependent autophagic apoptosis in HepG2 cells.
2. Sarmentosin induced autophagy via activating Nrf2 and regulating phosphorylation of mTOR in HepG2 cells.
3. Inhibition of caspase-dependent autophagic apoptosis via Sarmentosin was a potential treatment for HCC.

Original Article

Sarmentosin Induces Autophagy-dependent Apoptosis via Activation of Nrf2 in Hepatocellular Carcinoma

Zhitao Jiang^{1#}, Liyuan Gao^{1#}, Chundi Liu¹, Jianchun Wang¹, Yi Han¹ and Jinhua Pan^{2##}*

¹Department of Pharmacy, Zhangjiagang Hospital of Traditional Chinese Medicine Affiliated to Nanjing University of Chinese Medicine, Zhangjiagang, Jiangsu, China; ²School of Pharmacy, Nanjing University of Chinese Medicine, Nanjing, Jiangsu, China

***Correspondence to:** Jinhua Pan, Nanjing University of Chinese Medicine, Nanjing, Jiangsu, China. ORCID: <https://orcid.org/0000-0002-0574-192X>. Tel/Fax: +86-512-56380623, E-mail: panjinhua@163.com