





Review Article

# Immune Regulatory Networks and Therapy of $\gamma\delta$ T Cells in Liver Cancer: Recent Trends and Advancements



Kun-Li Yin<sup>1#</sup>, Kai-Jian Chu<sup>2#</sup>, Ming Li<sup>1</sup>, Yu-Xin Duan<sup>1</sup>, Yan-Xi Yu<sup>1</sup>, Mei-Qing Kang<sup>1</sup>, Da Fu<sup>3\*</sup>  and Rui Liao<sup>1\*</sup> 

<sup>1</sup>Department of Hepatobiliary Surgery, the First Affiliated Hospital of Chongqing Medical University, Chongqing, China; <sup>2</sup>Biliary Surgical Department I, the Eastern Hepatobiliary Surgical Hospital, Naval Medical University, Shanghai, China; <sup>3</sup>General Surgery, Institute of Pancreatic Diseases, Ruijin Hospital, Shanghai Jiaotong University School of Medicine, Shanghai, China

Received: 10 August 2023 | Revised: 20 November 2023 | Accepted: 19 December 2023 | Published online: 08 January, 2024

## Abstract

The roles of  $\gamma\delta$  T cells in liver cancer, especially in the potential function of immunotherapy due to their direct cytotoxic effects on tumor cells and secretion of important cytokines and chemokines, have aroused research interest. This review briefly describes the basic characteristics of  $\gamma\delta$  T cells, focusing on their diverse effects on liver cancer. In particular, different subtypes of  $\gamma\delta$  T cells have diverse or even opposite effects on liver cancer. We provide a detailed description of the immune regulatory network of  $\gamma\delta$  T cells in liver cancer from two aspects: immune components and nonimmune components. The interactions between various components in this immune regulatory network are dynamic and pluralistic, ultimately determining the biological effects of  $\gamma\delta$  T cells in liver cancer. We also integrate the current knowledge of  $\gamma\delta$  T-cell immunotherapy for liver cancer treatment, emphasizing the potential of these cells in liver cancer immunotherapy.

**Citation of this article:** Yin KL, Chu KJ, Li M, Duan YX, Yu YX, Kang MQ, *et al.* Immune Regulatory Networks and Therapy of  $\gamma\delta$  T Cells in Liver Cancer: Recent Trends and Advancements. *J Clin Transl Hepatol* 2023. doi: 10.14218/JCTH.2023.00355.

## Introduction

Liver cancer, as a globally notorious malignant tumor, causes hundreds of thousands of deaths and poses a serious socioeconomic burden.<sup>1,2</sup> The pathogenesis of liver cancer is unclear and is inseparable from its complex tumor microen-

vironment (TME).<sup>3</sup> The protumor/antitumor factors or cellular/noncellular components form a multicellular network in the TME, collectively determining the malignant biological behaviors of liver cancer cells.<sup>4-7</sup> Since patients with liver cancer are often diagnosed in the middle to late stages, most patients lose the opportunity for surgical radical resection and face a poor prognosis.<sup>8</sup> Notably, immunotherapy has significantly improved the treatment of liver cancer.<sup>9-11</sup> Cellular immunotherapy, involving natural killer cells (NKs),<sup>12,13</sup> chimeric antigen receptor T cells (CAR-T cells),<sup>14,15</sup> and  $\gamma\delta$  T cells, is emerging as a star in the field of immunotherapy, with its efficacy confirmed by various of mechanistic studies and clinical research.

$\gamma\delta$  T cells belong to the innate but nonconventional lymphocyte family, constituting approximately 5% (on average) of total T cells in peripheral blood.<sup>16-18</sup>  $\gamma\delta$  T cells, comprising 15-25% of the total liver T cells, are 5-10 times more abundant in the liver than in other tissues and organs.<sup>19</sup> Moreover, hepatic  $\gamma\delta$  T cells are highly localized in the liver, exhibiting a more active and mature phenotype with high CD44 and low CD62 markers.<sup>20</sup> In the thymus, "double negative" (DN) thymocytes, the primary immature precursors of  $\gamma\delta$  T cells, undergo rearrangement after  $\alpha$ -selection,  $\beta$ -selection,  $\gamma$ -selection and  $\delta$ -selection. Following TCR  $\gamma\delta$  signal stimulation,<sup>21</sup> TCR  $\gamma\delta$ + DN progenitors are directed into the  $\gamma\delta$  T-cell lineage and subsequently differentiate and mature into  $\gamma\delta$  T cells.<sup>22,23</sup> Additionally, the gene expression profile reveals that the gene signatures of  $\gamma\delta$  T cells are equivalent to those of a mixture of  $\alpha\beta$  T cells and NK cells, endowing  $\gamma\delta$  T cells with the characteristics of both cell types.<sup>24</sup> Compared with  $\alpha\beta$  T cells,  $\gamma\delta$  T cells typically exhibit different immune phenotypes. Besides the common expression of CD2, CD3, CD5, and CD7,  $\gamma\delta$  T cells are usually negative for CD4 and CD8, with occasional CD8 positivity.<sup>25</sup> Although they both express CD3, the expression level is higher in  $\gamma\delta$  T cells than in  $\alpha\beta$  T cells.<sup>26</sup> The most notable distinction from  $\alpha\beta$  T cells is that  $\gamma\delta$  T cells are not restricted by major histocompatibility complex (MHC) molecules,<sup>27</sup> enabling them to be activated within minutes to detect molecular signals stimulated by infection or cancer and subsequently produce a large number of pro-inflammatory cytokines and chemokines.<sup>21,28</sup>

Much research attention is promptly focused on understanding the relationship between  $\gamma\delta$  T cells and tumors, as well as the therapeutic prospects of  $\gamma\delta$  T cells in antitumor activities. Here, we summarize the functional plasticity of  $\gamma\delta$  T cells and their dual role in liver cancer. We also discuss the

**Keywords:** Liver cancer;  $\gamma\delta$  T cells; Immunotherapy; Tumor microenvironment.

**Abbreviations:** APCs, antigen-presenting cells; ADCC, antibody-dependent cell-mediated cytotoxicity; CAR-T cells, chimeric antigen receptor T cells; CCR, C-C chemokine receptor; CCL, C-C motif chemokine ligand; DCs, dendritic cells; HSCs, hepatic stellate cells; MDSCs, myeloid-derived suppressor cells; NKs, natural killer cells; NKR1, natural killer cell receptors; TME, tumor microenvironment; ZOL, zoledronate.

\*Contributed equally to this work.

**Correspondence to:** Da Fu, General Surgery, Institute of Pancreatic Diseases, Ruijin Hospital, Shanghai Jiaotong University School of Medicine, Shanghai 200025, China. ORCID: <https://orcid.org/0000-0002-0878-2575>. Tel: +86-15921527578, Fax: +86-21-34186000, E-mail: [fu800da900@126.com](mailto:fu800da900@126.com); Rui Liao, Department of Hepatobiliary Surgery, the First Affiliated Hospital of Chongqing Medical University, Chongqing 400016, China. ORCID: <https://orcid.org/0000-0002-0057-2792>. Tel: +86-15922963763, Fax: +86-23-89011016, E-mail: [liao99@163.com](mailto:liao99@163.com)

immune regulatory networks of  $\gamma\delta$  T cells involved in the microenvironment of liver cancer,  $\gamma\delta$  T-cell-based immunotherapy in liver cancer and perspectives on its development and use in the future.

## Subtypes of $\gamma\delta$ T cells and pathophysiological roles

### Classification of $\gamma\delta$ T Cells

Human  $\gamma\delta$  T cells are usually divided into  $V\delta 1+$ ,  $V\delta 2+$ , and  $V\delta 3+\gamma\delta$  T cells according to the difference in the arrangement of the  $\gamma$  chain and  $\delta$  chain.<sup>29</sup>  $V\delta 1+\gamma\delta$  T cells are mainly distributed in the epithelium and mucosa, accounting for approximately 5–10% of  $\gamma\delta$  T cells.<sup>30,31</sup>  $V\delta 2+\gamma\delta$  T cells are mainly present in peripheral blood, constituting the majority of the total number of  $\gamma\delta$  T cells. In particular,  $\gamma\delta$  T cells expressing the  $V\gamma 9V\delta 2$  TCR account for 50–95% of total  $\gamma\delta$  T cells and are also the main cells responsible for the antitumor effects.<sup>32</sup>  $V\delta 3+$   $\gamma\delta$  T cells are mainly present in the liver and small intestine epithelium. There is relatively little research on the role of  $V\delta 3+$   $\gamma\delta$  T cells in tumors, but there is still research potential.<sup>33,34</sup> Furthermore,  $V\delta 1+$   $\gamma\delta$  T cells and  $V\delta 2+$   $\gamma\delta$  T cells can be further classified by the expression of different CD molecules and the skewed expression patterns of the memory markers CD45RA and CD27.  $V\delta 1+$   $\gamma\delta$  T cells from healthy individuals mainly represent a CD45RA+CD27+ phenotype,<sup>35</sup> with high expression of CD56.<sup>36</sup> However,  $V\delta 1+\gamma\delta$  T cells in tumor tissue tend to exhibit a CD45RA-CD27- phenotype.<sup>37,38</sup>  $V\delta 2+$   $\gamma\delta$  T cells mainly exhibit a CD45RA- CD27+ phenotype, with CCR5 and CD161 generally overexpressed, allowing them to quickly migrate to inflammatory tissue.<sup>36</sup> Accordingly,  $\gamma\delta$  T cells display subset-specific features that need to be emphasized, as these features increase the understanding of the true role of  $\gamma\delta$  T cells in different diseases, especially their role in cancer.

### Pathophysiological roles of $\gamma\delta$ T cells

$\gamma\delta$  T cells exert an extensive array of biological effects. Their main characteristics include cytotoxic activity against target cells and the secretion of different types of cytokines and chemokines.  $\gamma\delta$  T cells can also interact with other cells, such as  $\alpha\beta$  T cells, dendritic cells (DCs), B cells, NKs and macrophages. Through these functional activities,  $\gamma\delta$  T cells exhibit anti-infection and antitumor effects and play a role in immune regulation and damage repair.

Numerous studies have revealed that  $\gamma\delta$  T cells exert their antitumor effect mainly through the production of cytokines (such as interferon  $\gamma$  (IFN  $\gamma$ ) and TNF- $\alpha$ ), triggering  $\gamma\delta$  TCR recognition molecules, NK cell receptors (NKR), and antibody-dependent cell-mediated cytotoxicity (ADCC). The recognition of  $\gamma\delta$  TCR depends on interactions with specific phosphoantigens,<sup>39–41</sup> such as butyrophilin 3 A1 (BTN3A1),<sup>42</sup> and subsequently activating  $V\gamma 9V\delta 2$  T cells. BTN2A1 is essential for BTN3A1-dependent  $V\gamma 9V\delta 2$  T-cell activation against cancer cells, regulating  $V\gamma 9V\delta 2$  T-cell binding to the TCR.<sup>43</sup> Representative molecules of the NK recognition mechanism include NKG2D,<sup>44</sup> NKp30,<sup>45</sup> and NKp44.<sup>45</sup> Various cytokines and chemokines assist  $\gamma\delta$  T cells in achieving cytotoxic functions while acting as regulators in the immune microenvironment. Some scientific evidence indicates that  $\gamma\delta$  T cells exhibit cytotoxic activity in liver cancer<sup>46–48</sup> and are under the regulation of cytokines such as IL-2 and IL-21.<sup>49</sup> Additionally, IL-17 and IL-2, the most common cytokines secreted by  $\gamma\delta$  T cells, have been identified as biomarkers of poor prognosis in malignant tumors, related to the maintenance and promotion of the inflammatory environment.<sup>50,51</sup>  $\gamma\delta$  T cells are the main providers of IFN  $\gamma$  in tumor immunity, and IFN  $\gamma$  plays

an crucial role in controlling tumor development.<sup>52,53</sup> Currently, it is acknowledged that the chemokines produced by  $\gamma\delta$  T cells include C-C chemokine receptor (CCR)2,<sup>54</sup> CCR5,<sup>55</sup> CCR6,<sup>56</sup> CCR7,<sup>57</sup> and CCR9,<sup>55</sup> which play an auxiliary role in  $\gamma\delta$  T cell anti-infection and antitumor functions. The recruitment and activation of  $\gamma\delta$  T cells depend on the reactivity of different chemokine signals. Chemokines induce  $\gamma\delta$  T cells to accurately migrate to the desired site, and  $\gamma\delta$  T cells and other cells produce chemokines, achieving this goal together.


Elucidating the interactions between  $\gamma\delta$  T cells and other immune cells will increase the understanding of  $\gamma\delta$  T cells.  $\gamma\delta$  T cells have been confirmed to function as antigen-presenting cells (APCs), similar to DCs, effectively processing and presenting antigens to CD4+ T cells and CD8+ T cells to initiate immune responses.<sup>58</sup> Furthermore,  $\gamma\delta$  T cells provide strong stimulation signals for the differentiation and maturation of DCs through the Fas-Fas L pathway.<sup>59,60</sup> Another study found that  $\gamma\delta$  T cells and DCs mutually promote maturation, and the increase in DC-induced apoptosis induces the expression of  $\gamma\delta$  T-cell ligands, thereby activating  $\gamma\delta$  T cells.<sup>59</sup> The proliferation and differentiation of  $\alpha\beta$  T cells are still induced by  $\gamma\delta$  T cells,<sup>61</sup> but in the tumor microenvironment,  $\gamma\delta$  T cells show an inhibitory effect on the activation of  $\alpha\beta$  T cells and support the occurrence of pancreatic tumors.<sup>62</sup> It has been proven that  $V\delta 2+$  and  $V\delta 3+\gamma\delta$  T cells affect the differentiation, antibody secretion, and cytokine production of B cells, starting from the moment that B cells are released from the bone marrow.<sup>63,64</sup>  $\gamma\delta$  T cells are associated with changes in the number of macrophages, and they also promote M2 macrophage polarization by secreting IL-17A, thus contributing to the clearance of infected cells.<sup>65</sup> Additionally, the number of peripheral  $\gamma\delta$  T cells is negatively correlated with the accumulation of neutrophils and M1 macrophages. Peripheral  $\gamma\delta$  T cells limit the expansion and recruitment of neutrophils to alleviate inflammation.<sup>66,67</sup> In a liver ischemia-reperfusion injury model,  $\gamma\delta$  TCR and IL-17a contribute to the increase in neutrophil numbers and exacerbate liver injury.<sup>68</sup> It is thus clear that the relationships between  $\gamma\delta$  T cells and other immune cells are extensive and far-reaching, shaping the body's vast immune regulatory network. The subtypes of  $\gamma\delta$  T cells and the pathophysiological roles of  $\gamma\delta$  T cells in tumors are summarized in Figure 1.

### A dual role of $\gamma\delta$ T cells in liver cancer

In the TME of liver cancer, the diversity of  $\gamma\delta$  T cell subpopulations in anti-tumor immune functions, such as the dynamic interactions between tumor promotion and inhibition, has received widespread attention, with special emphasis on the differences between  $V\delta 1+$  and  $V\delta 2+$   $\gamma\delta$  T cells. The functional paradigm of  $\gamma\delta$  T cells and their secreted factors was defined in tumor immunity through an in-depth understanding of  $\gamma\delta$  T cell subpopulations, enabling these subpopulations to exert anti-tumor effects in the tumor microenvironment. However, the  $\gamma\delta$  T cell subpopulations and their corresponding tumor effects are still being refined. With the development of more experimental methods, the dual role and mechanism of  $\gamma\delta$  T cells in liver cancer will be elucidated in greater detail and accuracy, becoming the key to explaining immune tolerance in liver cancer. A dual role of  $\gamma\delta$  T cells in liver cancer is summarized in Figure 2.

### Antitumor immunosurveillance in liver cancer

Studies using  $\gamma\delta$  T cells, particularly  $V\gamma 9V\delta 2$  T cells, to treat liver cancer have shown positive results. A high proportion of  $V\gamma 9V\delta 2$  T cells is associated with longer HCC patient survival times.<sup>69,70</sup> NKRs expressed by  $\gamma\delta$  T cells, especially the NK-




**Fig. 1. The subtypes of  $\gamma\delta$  T cells and the pathophysiological roles of  $\gamma\delta$  T cells in tumors.** Human  $\gamma\delta$  T cells are usually divided into  $V\delta 1+$ ,  $V\delta 2+$ , and  $V\delta 3+$   $\gamma\delta$  T cells according to the arrangement of the  $\gamma$  chain and  $\delta$  chain.  $V\delta 1+$   $\gamma\delta$  T cells and  $V\delta 2+$   $\gamma\delta$  T cells can be further classified by the expression of the memory markers CD45RA and CD27. The pathophysiological roles of  $\gamma\delta$  T cells include their cytotoxic activity against target cells, the secretion of different types of cytokines and chemokines and their interaction with other cells, such as  $\alpha\beta$  T cells, dendritic cells (DCs), B cells, natural killer cells (NKs), and macrophages. DCs, dendritic cells; NKs, natural killer cells. IFN- $\gamma$ , interferon  $\gamma$ ; CCR, C-C chemokine receptor; TNF- $\alpha$ , tumor necrosis factor- $\alpha$ ; IL, interleukin; TCR, T cell receptor.

G2D receptor, are the main cell type that combats cancer.<sup>71</sup> High expression of the NKG2D receptor is beneficial for enhancing the cytotoxicity of  $\gamma\delta$  T cells against HCC cell lines.<sup>72</sup> The ULBP family, which binds to NKRs and is expressed by tumors, is involved in the recognition of tumor antigens by  $V\gamma 9V\delta 2$  T cells, and its expression level determines the susceptibility of  $V\gamma 9V\delta 2$  T-cell lysis.<sup>73,74</sup> High ULBP1 expression is positively correlated with the degree of severity of HCC.<sup>75</sup> In patients with HCC, ULBP1 deficiency is an independent risk factor for early recurrence and poor prognosis.<sup>72</sup> After binding to NKG2D, the release of IFN- $\gamma$  and TNF- $\alpha$  is induced to promote  $\gamma\delta$  T-cell cytolytic activity.<sup>76</sup>  $\gamma\delta$  T cells are important sources of IFN- $\gamma$  and TNF- $\alpha$ , which are known to inhibit tumor growth by specifically inducing apoptosis and inhibiting angiogenesis.<sup>77,78</sup>  $\gamma\delta$  T cells expressing DNAX accessory molecule-1 (DNAM-1) and CD96 can more efficiently recognize Nectin-like-5 expressed on HCC cells and enhance the lysis of HCC cells, with a concomitant increase in IFN- $\gamma$  production.<sup>79</sup> In addition, research has found that CD96+  $\gamma\delta$  T cells are released from the liver and circulate in the bloodstream, having an inhibitory effect on the progression of other tumors.<sup>80</sup> Several reports have shown that the killing ability of  $\gamma\delta$  T cells against cancer cells is related to external

factors. A co-culture system of zoledronate (ZOL),<sup>81</sup> alendronate (ALD),<sup>82</sup> or artesunate<sup>83</sup> with  $\gamma\delta$  T cells can effectively activate  $\gamma\delta$  T cells and increase tumor susceptibility. The antitumor properties of  $\gamma\delta$  T cells have great clinical application value. Further in-depth research is still needed to determine how  $\gamma\delta$  T cells can increase their own efficacy and enhance their recognition of HCC cells.

#### Tumor-promoting tolerance in liver cancer

Some studies have revealed the protumor effects of  $\gamma\delta$  T cells. HCC-infiltrating  $\gamma\delta$  T cells are mainly composed of IL-17-producing  $V\delta 1$   $\gamma\delta$  T cells,<sup>84</sup> which play a role in promoting tumor proliferation,<sup>85</sup> angiogenesis,<sup>86</sup> metastasis<sup>87</sup> and immunotherapy resistance.<sup>88</sup>  $V\delta 1$   $\gamma\delta$  T cells are considered tissue-resident cells that are significantly enriched in the liver.<sup>69</sup> Tissue residence produces a local immune tolerance effect that can be maintained for a long time, which is becoming a key feature of antitumor protection.<sup>89</sup> Further research has found that the  $V\delta 1+$  T-cell population, which highly expresses LAG3, negatively regulates the antitumor function of T cells through immune suppression.<sup>90</sup> In the HCC mouse model, IL-17A was secreted mainly by  $V\gamma 4$   $\gamma\delta$  T cells, so re-



**Fig. 2. A dual role of  $\gamma\delta$  T cells in liver cancer.** Different subtypes of  $\gamma\delta$  T cells have diverse or even opposite effects on liver cancer. IFN- $\gamma$ , interferon  $\gamma$ ; TGF- $\beta$ , tumor growth factor- $\beta$ ; TNF- $\alpha$ , tumor necrosis factor- $\alpha$ ; IL, interleukin; LAG3, lymphocyte activation gene-3; PD-1, programmed death -1; CD96, cluster of differentiation-96; ULBP1, ul16-binding protein; NKGD2, natural killer group member D.

plenishing V $\gamma$ 4  $\gamma\delta$  T cells promoted tumor growth by recruiting myeloid-derived suppressor cells (MDSCs) in a CXCL5/CXCR2-dependent manner and further inhibiting CD8+ T cells to enhance immune suppression.<sup>91</sup> More recently, among HCC patients, the cytotoxicity of  $\gamma\delta$  T cells was significantly reduced, which is related to the expression of PD-1 in  $\gamma\delta$  T cells being significantly upregulated. Coincubation of  $\gamma\delta$  T cells leads to an increased proportion of PD-L1+ HCC cells, which is highly likely to enhance the immune escape of tumor cells.<sup>49</sup> Interestingly, V $\gamma$ 9V $\delta$ 2 T cells, also representing immunosuppressive properties, deviate into Th1-, Th17-, or Treg-like phenotypes depending on the environment and secrete immunosuppressive factors (IL-4, IL-17 or IL-10 and TGF- $\beta$ ) to inhibit T-cell proliferation and indirectly suppress tumor immunity.<sup>92,93</sup> Thus, the tumor-promoting effect of  $\gamma\delta$  T cells on liver cancer is comprehensive. Weakening or inhibiting the antitumor effect of  $\gamma\delta$  T cells is both a challenge and an opportunity for liver cancer research.

### Immune regulatory networks of $\gamma\delta$ T cells involved in the microenvironment of liver cancer

#### Nonimmune component regulatory networks

The immune regulatory network of  $\gamma\delta$  T cells in liver cancer is extremely sophisticated and multivariant, of which non-

immune components are an important factor. Environmental factors, such as reactive oxygen species production,<sup>94</sup> oxygen tension,<sup>95</sup> and lipoprotein levels,<sup>95</sup> have been proven to impact the expression of IFN  $\gamma$  and/or NKR in  $\gamma\delta$  T cells. He *et al.* used single-cell sequencing to explain the metabolic changes of HCC-infiltrating  $\gamma\delta$  T cells which are significantly inhibited in oxygen metabolism, lipid metabolism, and amino acid synthesis, and tilt towards glutamine-rich metabolism, providing nutrients for tumor cell metabolism.<sup>90</sup> Combined with a genome-wide CRISPR screening of target cancer cells, another study found that the triggering of  $\gamma\delta$  T cells by BTN3A and BTN2A1 depended on AMP-activated protein kinase (AMPK), and demonstrated that  $\gamma\delta$  T cells are regulated by multiple layers of BTN3A, such as transcription, post-translational modifications, and membrane transport, further deepening our understanding of  $\gamma\delta$  T cell stress monitoring.<sup>96</sup> The homeostasis of IL-17A-producing  $\gamma\delta$  T cells retained in the liver is regulated by palmitic acid and CD1d, a necessary lipid antigen.<sup>97</sup> Moreover, apoptosis, ferroptosis, and pyroptosis significantly induce an immunosuppressive microenvironment in HCC, in which an imbalance in  $\gamma\delta$  T cells and an increase in the Vd1+/Vd2+ ratio are markers of poor prognosis in HCC patients.<sup>98</sup> Several reports have shown that the tumor-killing ability of  $\gamma\delta$  T cells is related to external factors. Artesunate enhances the antitumor function of  $\gamma\delta$  T cells by upregulating the expression of Fas on HepG2 cells while re-



ducing the secretion of TGF- $\beta$  by HepG2 cells, thus reversing the immune escape of HepG2 cells.<sup>83</sup> The co-cultivation of zoledronate and HCC cell lines also resulted in an increase in the number and killing ability of  $\gamma\delta$  T cells.<sup>99</sup> Overexpression of miR-382 in normal liver cells increases the sensitivity of  $\gamma\delta$  T cells to HCC, thereby promoting HCC cell lysis.<sup>100</sup> Macrophage-stimulating protein (MSP) and peptide HP1 stimulate V $\gamma$ 9V $\delta$ 2 T cells, which have high expression levels of IFN- $\gamma$  and TNF- $\alpha$ , to exert a Th1-biased immune response for inhibiting HCC.<sup>46</sup> Other cellular components in the liver, such as hepatic stellate cells, activate TLR3 under the mediation of exosomes, thereby increasing  $\gamma\delta$  T cell-mediated production of IL-17A, which exacerbates liver fibrosis.<sup>101</sup> Evidently, the importance of  $\gamma\delta$ T cell and cellular component interactions in elucidating the mechanism of liver cancer has begun to be recognized.<sup>102</sup>

### **Immune cellular regulatory networks**

Crosstalk between  $\gamma\delta$  T cells and other immune cells, such as neutrophils, macrophages, T cells, and B cells, is widespread and has been identified in many malignant tumors. The immune balance in liver cancer is the result of  $\gamma\delta$  T cell self-regulation and their interactions with other immune cells. Tumor-infiltrating  $\gamma\delta$  T cells respond to signals from microorganisms and tumors, activate CD4+ and CD8+ T cells,<sup>103</sup> and regulate the number and function of CD4, CD8, and NK cells.<sup>90</sup> The activation of CD8+ T cells by  $\gamma\delta$  T cells may be achieved through the scavenger receptor CD36, resulting in tumor antigen-specific CD8+ T-cell responses.<sup>104</sup> The regulation between different immunocytes is often bidirectional, but the effect of CD4+, CD8+, and NK cells on  $\gamma\delta$  T cells has not yet been revealed. The elucidation of these effects will help clarify the mechanism of  $\gamma\delta$  T cells in liver cancer and even other diseases. IL-17A produced by HCC-infiltrating V $\gamma$ 4  $\gamma\delta$  T cells enhances the infiltration of MDSCs, leading to the inhibition of the CD8+ T-cell response and the promotion of tumor growth.<sup>91</sup> Additionally, MDSCs also harbor the capacity to modulate the generation of IFN- $\gamma$  and the antitumor effect of V $\delta$ 2  $\gamma\delta$  T cells in the tumor microenvironment.<sup>105</sup> Moreover, TEM results have shown that there is a negative correlation between the number of  $\gamma\delta$  T cells and the number of Treg cells in HCC, possibly because CD4+CD25+ regulatory T cells directly suppress the cytotoxic function and IFN- $\gamma$  secretion of  $\gamma\delta$  T cells in a TGF- $\beta$ - and IL-10-dependent manner.<sup>106</sup> On the other hand,  $\gamma\delta$  T cells can differentiate into  $\gamma\delta$  Treg cells and play an immunosuppressive role in cancer, as tumor-derived  $\gamma\delta$  Treg cells promote T cell and DC senescence through TLR8 signaling. This results in the inhibition of innate and adaptive immunity to maintain tumor suppression.<sup>107,108</sup> To date, there is limited research evidence for the interaction between  $\gamma\delta$  T cells and other immune cells in liver cancer, which can help elucidate the occurrence and development of the liver. The relationships between  $\gamma\delta$  T cells and other immune or nonimmune cells in liver cancer are worth further exploration.

### **Immune molecular regulatory networks**

Immune molecules are both executors and regulators of  $\gamma\delta$  T cell function. Relying on various immune molecules,  $\gamma\delta$  T cells successfully interact with other cells to construct a meticulous and efficient regulatory network in the liver cancer microenvironment. The high expression of CCR1/CCR5 makes  $\gamma\delta$  T cells recruited by C-C motif chemokine ligand 4 (CCL4)/CCL5 into HCC tissue, exerting protective effects on CD8+ T cells and strengthening the body's antitumor ability.<sup>109</sup> Another study pointed out that the high expression

of CCR5 is a selective feature of V $\delta$ 2/V $\gamma$ 9  $\gamma\delta$  T cells, which is consistent with the shift in antitumor ability.<sup>110</sup> Blockade of the CCL2/CCR2 axis has been considered a novel protective factor during the recruitment of V $\delta$ 1 T cells to HCC lesions.<sup>54,111</sup> Furthermore, mutual crosstalk between chemokines and cytokines should be emphasized, which serves as a feedback mechanism that determines the final performance of  $\gamma\delta$  T cells in liver cancer.<sup>112,113</sup> The binding of CCR6 and CCL20 significantly aggregates IL-17- and IL-22-expressing  $\gamma\delta$  T cells in damaged liver cells, which promotes the apoptosis of hepatic stellate cells (HSCs) involving Fas ligands, preventing the liver from entering an inflammatory and fibrotic state.<sup>114</sup>


For the interleukin family, TNF- $\alpha$  and INF- $\gamma$  are both produced by and act upon  $\gamma\delta$  T cells. For example, the decrease in the cytotoxicity of  $\gamma\delta$  T cells in the blood circulation of HCC patients may be explained by the fact that the levels of some pro-inflammatory cytokines (IL-2 and IL-21) are reduced and PD-1 is upregulated.<sup>49</sup> Moreover, tumor-infiltrating V $\gamma$ 1  $\gamma\delta$  T cells produce IL-4, which significantly reduces the expression levels of NKG2D, perforin-1, and IFN- $\gamma$  in V $\gamma$ 4  $\gamma\delta$  T cells, diminishing cytotoxicity and reducing antitumor function.<sup>115</sup> IL-35, an immunosuppressive cytokine, is produced by Treg cells in the hepatic microenvironment, and its overexpression promotes the depletion of  $\gamma\delta$  T cells and impairs their antitumor function in HCC.<sup>116</sup> However, the influence of PD-1 expression on  $\gamma\delta$  T cells is not well defined. In hematological tumors, the blockade of PD-1 does not directly affect the cell-dependent lysis ability of  $\gamma\delta$  T cells, but PD-1+  $\gamma\delta$  T cells produce more IFN- $\gamma$  to enhance their own lethality.<sup>117</sup> Similarly, IL-17A production from  $\gamma\delta$  T cells can be inhibited by high expression of PD-1 while reducing organ inflammation damage and immune resistance.<sup>88,118</sup> Other studies contradict the above viewpoints, suggesting that high numbers of PD-1+  $\gamma\delta$  T cells are correlated with a poor prognosis for patients with acute leukemia.<sup>119</sup> In HCC cell lines, the cytotoxicity of PD-1+  $\gamma\delta$  T cells is weakened.<sup>49</sup> Deeper exploration of the relationship between  $\gamma\delta$  T cells and PD-1 expression will lead to a more comprehensive understanding of immunotherapy. Thus, the immunoregulatory balance of  $\gamma\delta$  T cells involved in liver cancer is not yet completely understood, but the joint forces of multiple cytokines, chemokines, and immunocytes determine the overall performance of  $\gamma\delta$  T cells in liver cancer. Brief summary of immune regulatory networks of  $\gamma\delta$  T cells involved in the microenvironment of liver cancer is summarized in Figure 3.

### **$\gamma\delta$ T-cell-based immunotherapy in liver cancer**

$\gamma\delta$  T-cell-based immunotherapy in liver cancer is being vigorously developed and has achieved surprising results. According to current research results, there are three mainstream directions for  $\gamma\delta$  T-cell-based immunotherapy, namely, synergistic effects of other antibodies, chimeric antigen receptor-based  $\gamma\delta$  T cells, and  $\gamma\delta$  T-cell transplantation. The practical application of  $\gamma\delta$  T cell-centered immunotherapy in liver cancer is both promising and challenging with tremendous room for exploration. Studies of potential  $\gamma\delta$  T-cell-based immunotherapy in liver cancer is summarized in Table 1.<sup>70,120–127</sup>

#### **Synergistic effects of other antibodies**

The cytotoxicity of  $\gamma\delta$  T cells in human liver sinuses depends on the signaling of phosphoantigens, NK receptors, and immune checkpoint molecules. Nitrogen-containing bisphosphonates have been widely used in tumor treatment, with outstanding therapeutic achievements.<sup>128–132</sup>



**Fig. 3. Brief summary of immune regulatory networks of  $\gamma\delta$  T cells involved in the microenvironment of liver cancer.** The immune regulatory network of  $\gamma\delta$  T cells in liver cancer is extremely sophisticated and multivariant. Immune regulatory networks of  $\gamma\delta$  T cells are involved in the immune microenvironment of liver cancer through immune molecules, immune cells and nonimmune components. CD, cluster of differentiation; NK cell, natural-killer cell; MDSC, Myeloid-derived suppressor cell; Treg cell, T regulatory cell; CCR, C-C chemokine receptor; IL, interleukin; IFN- $\gamma$ , interferon  $\gamma$ ; TNF- $\alpha$ , tumor necrosis factor- $\alpha$ .

Similarly, in HCC, the addition of zoledronate markedly optimizes  $\gamma\delta$  T-cell-mediated immunotherapy and directly inhibits tumor proliferation.<sup>120,133</sup> Moreover, the addition of EpCAM-specific monoclonal antibodies can prominently enhance the lysis effect of  $\gamma\delta$  T cells on hepatoblastoma.<sup>121</sup> Another study also revealed a thought-provoking phenomenon in which chemotherapy accelerates the immune senescence and dysfunction of V $\delta$ 2  $\gamma\delta$  T cells in patients with metastatic liver cancer.<sup>122</sup> Therefore, the combination of  $\gamma\delta$  T cells and other drugs or antibodies is a double-edged sword that requires more clinical practice to provide more favorable evidence.

#### **Chimeric antigen receptor (CAR) T-cell immunotherapy**

CAR-T-cell immunotherapy is a precise and efficient tumor treatment method, while CAR- $\gamma\delta$  T cells possess the advantages of both treatments, pushing tumor immunotherapy to a new level. This strategy has been applied in clinical practice in hematological tumors and has great potential in solid tumors.<sup>123,134–137</sup> V $\delta$ 1  $\gamma\delta$  T cells engineered with glypican-3-specific CAR and soluble IL-15 efficiently and robustly control tumor growth in HCC and exhibit strong tumor tissue aggregation.<sup>124</sup> Although promising, some technical problems related to CAR- $\gamma\delta$  T cells still need to be addressed, such

as antigen recognition and design, gene delivery technology, and cell proliferation.

#### **$\gamma\delta$ T-cell transplantation.**

As HCC progresses, the function of  $\gamma\delta$  T cells is lost, and the V $\delta$ 2  $\gamma\delta$  T-cell population is heavily consumed.<sup>90</sup> At the same time, the expression of immunosuppressive receptors in the V $\delta$ 1  $\gamma\delta$  T-cell population is upregulated, and the antitumor ability of V $\delta$ 2  $\gamma\delta$  T cells is enhanced.<sup>138</sup> Therefore, allogeneic V $\delta$ 2  $\gamma\delta$  T cells serve as a promising supplement in the treatment of liver cancer. In a study of V $\gamma$ 9V $\delta$ 2 T-cell adoptive therapy, expanded cells also displayed robust antitumor efficacy against HCC and improved immune efficacy functions, including antitumor ability, proliferation, and differentiation.<sup>70</sup> In corresponding clinical trials, the prolonged survival time of patients confirmed the safety and effectiveness of allogeneic V $\gamma$ 9V $\delta$ 2 T-cell immunotherapy.<sup>70,125,128</sup> Four clinical trials related to liver cancer have been registered so far (<https://www.clinicaltrials.gov/> ID: NCT02425735, NCT03183219, NCT04518774, NCT05628545). In this clinical trial of advanced liver cancer (NCT03183219), of the 8 liver cancer patients, 7 survived for more than 10 months, and 3 survived for more than 30 months.<sup>70</sup> Moreover, local regional therapy combined with allogeneic  $\gamma\delta$  T-cell adoptive transplantation is also very safe and effective in the treatment of advanced

**Table 1. Studies of potential  $\gamma\delta$  T-cell-based immunotherapy in liver cancer**

Treatment	Tumor type	Effector cells	Mechanism	Outcomes	Reference
Zoledronic acid	SK-HEP-1 and H22 cells	–	ZOL optimizes $\gamma\delta$ T-cell-mediated immunotherapy and inhibits growth of HCC cells	Inhibited tumor proliferation	120
Zoledronate	Hepatocellular carcinoma and colorectal carcinoma with hepatic metastases	V $\gamma$ 9V $\delta$ 2 T cells	The cytotoxic level of V $\gamma$ 9V $\delta$ 2 T cells against freshly autologous tumor cells isolated from patients could be significantly increased by pretreating the tumor cells with zoledronate	Improved cytotoxicity	123
Zoledronate and IL-2	Hepatocellular carcinoma	$\gamma\delta$ T cells	Zoledronate with IL-2 may efficiently expand $\gamma\delta$ T cells sourced from the peripheral blood of patients with HCC	Increased the quantity of the $\gamma\delta$ T cells	127
EpCAM-specific monoclonal antibodies	Hepatoblastoma	$\gamma\delta$ T cells	Tumor cell lysis by $\gamma\delta$ T cells can be dramatically augmented	Improved cytotoxicity	121
Chemotherapy	Metastatic liver cancer	V $\delta$ 2+ T cells	V $\delta$ 2+ T cells are coupled with impairments in quantity, cytotoxicity and production of TNF- $\alpha$ and IFN- $\gamma$	Decreased cytotoxicity	122
Glypican-3 (GPC-3)-specific chimeric antigen receptor (CAR) and soluble IL-15	Hepatocellular carcinoma	CAR V $\delta$ 1 T cells	GPC-3.CAR/sIL-15 V $\delta$ 1 T cells displayed robust in vitro and vivo proliferation, cytokine production, cytotoxic activity and suppression of tumor growth	Delayed tumour growth	124
The expanded $\gamma\delta$ T cells from healthy donors	Late-stage liver cancer	V $\gamma$ 9V $\delta$ 2 T cells	The expanded $\gamma\delta$ T cells possessed significantly improved immune effector functions, including proliferation, differentiation, and cancer cell killing, both in vitro and in the humanized mouse model, with the safety and efficacy in clinical trial	Improved antitumor efficacy and prolonged survival time of patients	70
The expanded $\gamma\delta$ T cells from healthy donors	Cholangiocarcinoma	V $\gamma$ 9V $\delta$ 2 T cell	Allogenic $\gamma\delta$ T cell treatments positively regulated peripheral immune functions of the patient, depleted tumor activity, improved quality of life, and prolonged his life span	Improved antitumor efficacy	125
Locoregional Therapy Combined with Adoptive Transfer of Allogeneic $\gamma\delta$ T Cells	Hepatocellular Carcinoma and Intrahepatic Cholangiocarcinoma	$\gamma\delta$ T cells	The novel combination of locoregional ablation with adoptive transfer of allogeneic $\gamma\delta$ T cells was safe, and patients with HCC in the combined treatment group had a longer OS.	Prolonged survival time of patients	126

HCC and intrahepatic cholangiocarcinoma.<sup>126</sup>

For liver cancer patients, the optimization of  $\gamma\delta$  T-cell immunotherapy emphasizes individualization. The quality of  $\gamma\delta$  T-cell amplification is related to clinical and pathological characteristics such as tumor size, quantity, and serum AFP levels.<sup>127</sup> During  $\gamma\delta$  T cell immunotherapy, the physiological and biochemical indicators of patients should be dynamically monitored to maximize the advantages of  $\gamma\delta$  T cell immunotherapy.<sup>127</sup>

### Conclusions and future perspectives

An increasing number of studies have confirmed the unique properties of  $\gamma\delta$  T cells and their plasticity in the field of cancer research. Different subtypes of  $\gamma\delta$  T cells and their secreted cytokines have different protumor or antitumor effects. The enormous immune regulatory network formed around  $\gamma\delta$  T cells, including immune and nonimmune components in the microenvironment of liver cancer, regulates the

differentiation and performance of  $\gamma\delta$  T cells and determines the ultimate occurrence and development of liver cancer.  $\gamma\delta$  T-cell-based immunotherapy has gradually become a highly promising treatment protocol for liver cancer, accompanied by opportunities and challenges. However, the exploration of  $\gamma\delta$  T cells in liver cancer is still incomplete, and more systematic and accurate explorations are still needed in many aspects. First, the heterogeneity of  $\gamma\delta$  T cells needs to be emphasized during research, as the functions of different subpopulations are completely different. Second, breakthroughs and innovations in technology and methods, such as the application of high-throughput technologies and the development of stable transfer models of  $\gamma\delta$  T cells, can broaden the horizons of  $\gamma\delta$  T cell research. Third, research on how to thoroughly induce the activation of  $\gamma\delta$  T cells in clinical applications and develop specific engineered  $\gamma\delta$  T cells targeting tumor antigens still requires in-depth explorations of tumor-related mechanisms. Fourth, combination strategies with other forms of treatment, such as surgery and targeted therapy, should also be

the focus of future clinical investigation. Although research on  $\gamma\delta$  T cells is still in its infancy due to some technological limitations,  $\gamma\delta$  T cells, as attractive antitumor candidates, undoubtedly increase the understanding of liver cancer, and the application of  $\gamma\delta$  T cells in the field of liver cancer has great potential and a promising future.

### Acknowledgments

Thanks 'HOME for Researchers' and 'Figdraw' for the graphic assistance.

### Funding

This study was supported partly by grants from National Natural Science Foundation (81972214, 82373348), Shanghai Natural Science Foundation (21140903500), the Natural Science Foundation of Chongqing (No. CSTB2022NSCQ-MSX0112); Science and Health Joint Research Project of Chongqing Municipality (2020GDRC013); Program for Youth Innovation in Future Medicine, Chongqing Medical University (W0087).

### Conflict of interest

RL has been an editorial board member of *Journal of Clinical and Translational Hepatology* since 2022. The other authors have no conflict of interests related to this publication

### Author contributions

Study concept and design (KLY, KJC, DF, RL), drafting of the manuscript (KJC, ML, YXD, YXY, MQK), critical revision of the manuscript for important intellectual content (KJC, ML, YXD, YXY, MQK), administrative or material support (KJC, DF, RL), and study supervision (KLY, KJC, DF, RL), critical funding (DF, RL). All authors have made a significant contribution to this study and have approved the final manuscript.

### References

- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018;68(6):394-424. doi:10.3322/caac.21492, PMID:30207593.
- Vogel A, Meyer T, Sapisochin G, Salem R, Saborowski A. Hepatocellular carcinoma. *Lancet* 2022;400(10360):1345-1362. doi:10.1016/S0140-6736(22)01200-4, PMID:36084663.
- Donne R, Lujambio A. The liver cancer immune microenvironment: Therapeutic implications for hepatocellular carcinoma. *Hepatology* 2023;77(5):1773-1796. doi:10.1002/hep.32740, PMID:35989535.
- Sas Z, Cendrowicz E, Weinhäuser I, Rygiel TP. Tumor Microenvironment of Hepatocellular Carcinoma: Challenges and Opportunities for New Treatment Options. *Int J Mol Sci* 2022;23(7):3778. doi:10.3390/ijms23073778, PMID:35409139.
- Giannelli G, Koudelkova P, Dituri F, Mikulits W. Role of epithelial to mesenchymal transition in hepatocellular carcinoma. *J Hepatol* 2016;65(4):798-808. doi:10.1016/j.jhep.2016.05.007, PMID:27212245.
- Lv Y, Wang Z, Yuan K. Role of Noncoding RNAs in the Tumor Immune Microenvironment of Hepatocellular Carcinoma. *J Clin Transl Hepatol* 2023;11(3):682-694. doi:10.14218/JCTH.2022.00412, PMID:3696988.
- Berasain C, Castillo J, Perugorria MJ, Latasa MU, Prieto J, Avila MA. Inflammation and liver cancer: new molecular links. *Ann N Y Acad Sci* 2009;1155:206-221. doi:10.1111/j.1749-6632.2009.03704.x, PMID:19250206.
- Villanueva A. Hepatocellular Carcinoma. *N Engl J Med* 2019;380(15):1450-1462. doi:10.1056/NEJMra1713263, PMID:30970190.
- Colli LM, Machiela MJ, Zhang H, Myers TA, Jessop L, Delattre O, *et al*. Landscape of Combination Immunotherapy and Targeted Therapy to Improve Cancer Management. *Cancer Res* 2017;77(13):3666-3671. doi:10.1158/0008-5472.CAN-16-3338, PMID:28446466.
- Zhou C, Weng J, Gao Y, Liu C, Zhu X, Zhou Q, *et al*. A Novel mRNA Signature Related to Immunity to Predict Survival and Immunotherapy Response in Hepatocellular Carcinoma. *J Clin Transl Hepatol* 2022;10(5):925-938. doi:10.14218/JCTH.2021.00283, PMID:36304510.
- Duan Y, Zhang H, Tan T, Ye W, Yin K, Yu Y, *et al*. The immune response of hepatocellular carcinoma after locoregional and systemic therapies:

- The available combination option for immunotherapy. *Biosci Trends* 2023. doi:10.5582/bst.2023.01275, PMID:37981319.
- Kumar N, Khakoo SI. Hepatocellular carcinoma: Prospects for natural killer cell immunotherapy. *HLA* 2018;92(1):3-11. doi:10.1111/tan.13275, PMID:29667374.
  - Chu J, Gao F, Yan M, Zhao S, Yan Z, Shi B, *et al*. Natural killer cells: a promising immunotherapy for cancer. *J Transl Med* 2022;20(1):240. doi:10.1186/s12967-022-03437-0, PMID:35606854.
  - June CH, O'Connor RS, Kawalekar OU, Ghassemi S, Milone MC. CAR T cell immunotherapy for human cancer. *Science* 2018;359(6382):1361-1365. doi:10.1126/science.aar6711, PMID:29567707.
  - Titov A, Zmievskaya E, Ganeeva I, Valiullina A, Petukhov A, Rakhmatullina A, *et al*. Adoptive Immunotherapy beyond CAR T-Cells. *Cancers (Basel)* 2021;13(4):743. doi:10.3390/cancers13040743, PMID:33670139.
  - Brenner MB, McLean J, Dialynas DP, Strominger JL, Smith JA, Owen FL, *et al*. Identification of a putative second T-cell receptor. *Nature* 1986;322(6075):145-149. doi:10.1038/322145a0, PMID:3755221.
  - Groh V, Porcelli S, Fabbri M, Lanier LL, Picker LJ, Anderson T, *et al*. Human lymphocytes bearing T cell receptor gamma/delta are phenotypically diverse and evenly distributed throughout the lymphoid system. *J Exp Med* 1989;169(4):1277-1294. doi:10.1084/jem.169.4.1277, PMID:2564416.
  - Carding SR, Egan PJ. Gammadelta T cells: functional plasticity and heterogeneity. *Nat Rev Immunol* 2002;2(5):336-345. doi:10.1038/nri797, PMID:12033739.
  - Racanelli V, Rehmann B. The liver as an immunological organ. *Hepatology* 2006;43(2 Suppl 1):S54-S62. doi:10.1002/hep.21060, PMID:16447271.
  - Hunter S, Willcox CR, Davey MS, Kasatskaya SA, Jeffery HC, Chudakov DM, *et al*. Human liver infiltrating  $\gamma\delta$  T cells are composed of clonally expanded circulating and tissue-resident populations. *J Hepatol* 2018;69(3):654-665. doi:10.1016/j.jhep.2018.05.007, PMID:29758330.
  - Fiala GJ, Gomes AQ, Silva-Santos B. From thymus to periphery: Molecular basis of effector  $\gamma\delta$ -T cell differentiation. *Immunol Rev* 2020;298(1):47-60. doi:10.1111/imr.12918, PMID:33191519.
  - Michie AM, Zúñiga-Pflücker JC. Regulation of thymocyte differentiation: pre-TCR signals and beta-selection. *Semin Immunol* 2002;14(5):311-323. doi:10.1016/s1044-5323(02)00664-7, PMID:12220932.
  - Ferreira LM. Gammadelta T cells: innately adaptive immune cells? *Int Rev Immunol* 2013;32(3):223-248. doi:10.3109/08830185.2013.783831, PMID:23617235.
  - Pont F, Familiades J, Déjean S, Fruchon S, Cendron D, Poupot M, *et al*. The gene expression profile of phosphoantigen-specific human  $\gamma\delta$  T lymphocytes is a blend of  $\alpha\beta$  T-cell and NK-cell signatures. *Eur J Immunol* 2012;42(1):228-240. doi:10.1002/eji.201141870, PMID:21968650.
  - Roden AC, Morice WG, Hanson CA. Immunophenotypic attributes of benign peripheral blood gammadelta T cells and conditions associated with their increase. *Arch Pathol Lab Med* 2008;132(11):1774-1780. doi:10.5858/132.11.1774, PMID:18976014.
  - Nicolas L, Monneret G, Debard AL, Blesius A, Gutowski MC, Salles G, *et al*. Human gammadelta T cells express a higher TCR/CD3 complex density than alphabeta T cells. *Clin Immunol* 2001;98(3):358-363. doi:10.1006/clim.2000.4978, PMID:11237559.
  - Willcox CR, Mohammed F, Willcox BE. The distinct MHC-unrestricted immunobiology of innate-like and adaptive-like human  $\gamma\delta$  T cell subsets-Nature's CAR-T cells. *Immunol Rev* 2020;298(1):25-46. doi:10.1111/imr.12928, PMID:33084045.
  - Melandri D, Zlatareva I, Chaleil RAG, Dart RJ, Chancellor A, Nussbaumer O, *et al*. The  $\gamma\delta$ TCR combines innate immunity with adaptive immunity by utilizing spatially distinct regions for agonist selection and antigen responsiveness. *Nat Immunol* 2018;19(12):1352-1365. doi:10.1038/s41590-018-0253-5, PMID:30420626.
  - Kazen AR, Adams EJ. Evolution of the V, D, and J gene segments used in the primate gammadelta T-cell receptor reveals a dichotomy of conservation and diversity. *Proc Natl Acad Sci U S A* 2011;108(29):E332-E340. doi:10.1073/pnas.1105105108, PMID:21730193.
  - Maeurer MJ, Martin D, Walter W, Liu K, Zitvogel L, Haluszczak K, *et al*. Human intestinal Vdelta1+ lymphocytes recognize tumor cells of epithelial origin. *J Exp Med* 1996;183(4):1681-1696. doi:10.1084/jem.183.4.1681, PMID:8666926.
  - Chowers Y, Holtmeier W, Harwood J, Morzycka-Wroblewska E, Kagnoff MF. The V delta 1 T cell receptor repertoire in human small intestine and colon. *J Exp Med* 1994;180(1):183-190. doi:10.1084/jem.180.1.183, PMID:8006582.
  - Pauza CD, Cairo C. Evolution and function of the TCR Vgamma9 chain repertoire: It's good to be public. *Cell Immunol* 2015;296(1):22-30. doi:10.1016/j.cellimm.2015.02.010, PMID:25769734.
  - Mangan BA, Dunne MR, O'Reilly VP, Dunne PJ, Exley MA, O'Shea D, *et al*. Cutting edge: CD1d restriction and Th1/Th2/Th17 cytokine secretion by human V $\delta$ 3 T cells. *J Immunol* 2013;191(1):30-34. doi:10.4049/jimmunol.1300121, PMID:23740951.
  - Zhao Y, Niu C, Cui J. Gamma-delta ( $\gamma\delta$ ) T cells: friend or foe in cancer development? *J Transl Med* 2018;16(1):3. doi:10.1186/s12967-017-1378-2, PMID:29316940.
  - Davey MS, Willcox CR, Joyce SP, Ladell K, Kasatskaya SA, McLaren JE, *et al*. Clonal selection in the human V $\delta$ 1 T cell repertoire indicates  $\gamma\delta$  TCR-dependent adaptive immune surveillance. *Nat Commun* 2017;8:14760. doi:10.1038/ncomms14760, PMID:28248310.
  - Sanz M, Mann BT, Ryan PL, Bosque A, Pennington DJ, Hackstein H, *et al*. Deep characterization of human  $\gamma\delta$  T cell subsets defines shared and lineage-specific traits. *Front Immunol* 2023;14:1148988. doi:10.3389/fimmu.2023.1148988, PMID:37063856.
  - Meraviglia S, Lo Presti E, Tosolini M, La Mendola C, Orlando V, Todaro M,



- et al*. Distinctive features of tumor-infiltrating  $\gamma\delta$  T lymphocytes in human colorectal cancer. *Oncoimmunology* 2017;6(10):e1347742. doi:10.1080/2162402X.2017.1347742, PMID:29123962.
- [38] Wu Y, Biswas D, Usaite I, Angelova M, Boeing S, Karasaki T, *et al*. A local human V $\delta$ 1 T cell population is associated with survival in nonsmall-cell lung cancer. *Nat Cancer* 2022;3(6):696–709. doi:10.1038/s43018-022-00376-z, PMID:35637401.
- [39] Tanaka Y, Morita CT, Tanaka Y, Nieves E, Brenner MB, Bloom BR. Natural and synthetic non-peptide antigens recognized by human gamma delta T cells. *Nature* 1995;375(6527):155–158. doi:10.1038/3751555a0, PMID:7753173.
- [40] Hintz M, Reichenberg A, Altincicek B, Bahr U, Gschwind RM, Kollas AK, *et al*. Identification of (E)-4-hydroxy-3-methyl-but-2-enyl pyrophosphate as a major activator for human gammadelta T cells in *Escherichia coli*. *FEBS Lett* 2001;509(2):317–322. doi:10.1016/S0014-5793(01)03191-X, PMID:11741609.
- [41] Sandstrom A, Peigné CM, Léger A, Crooks JE, Konczak F, Gesnel MC, *et al*. The intracellular B30.2 domain of butyrophilin 3A1 binds phosphoantigens to mediate activation of human V $\gamma$ 9V $\delta$ 2 T cells. *Immunity* 2014;40(4):490–500. doi:10.1016/j.immuni.2014.03.003, PMID:24703779.
- [42] Payne KK, Mine JA, Biswas S, Chaurio RA, Perales-Puchalt A, Anadon CM, *et al*. BTN3A1 governs antitumor responses by coordinating  $\alpha\beta$  and  $\gamma\delta$  T cells. *Science* 2020;369(6506):942–949. doi:10.1126/science.aay2767, PMID:32820120.
- [43] Cano CE, Pasero C, De Gassart A, Kerneur C, Gabriac M, Fullana M, *et al*. BTN2A1, an immune checkpoint targeting V $\gamma$ 9V $\delta$ 2 T cell cytotoxicity against malignant cells. *Cell Rep* 2021;36(2):109359. doi:10.1016/j.celrep.2021.109359, PMID:34260935.
- [44] Simões AE, Di Lorenzo B, Silva-Santos B. Molecular Determinants of Target Cell Recognition by Human  $\gamma\delta$  T Cells. *Front Immunol* 2018;9:929. doi:10.3389/fimmu.2018.00929, PMID:29755480.
- [45] Silva-Santos B, Strid J. Working in “NK Mode”: Natural Killer Group 2 Member D and Natural Cytotoxicity Receptors in Stress-Surveillance by  $\gamma\delta$  T Cells. *Front Immunol* 2018;9:851. doi:10.3389/fimmu.2018.00851, PMID:29740448.
- [46] Xi X, Guo Y, Zhu M, Qiu F, Lei F, Li G, *et al*. Identification of new potential antigen recognized by  $\gamma\delta$ T cells in hepatocellular carcinoma. *Cancer Immunol Immunother* 2021;70(7):1917–1927. doi:10.1007/s00262-020-02826-y, PMID:33399933.
- [47] Hoh A, Dewerth A, Vogt F, Wenz J, Baeuerle PA, Warmann SW, *et al*. The activity of  $\gamma\delta$  T cells against paediatric liver tumour cells and spheroids in cell culture. *Liver Int* 2013;33(1):127–136. doi:10.1111/liv.12011, PMID:23088518.
- [48] Ren H, Li W, Liu X, Zhao N.  $\gamma\delta$  T cells: The potential role in liver disease and implications for cancer immunotherapy. *J Leukoc Biol* 2022;112(6):1663–1668. doi:10.1002/JLB.5MR0822-733RRR, PMID:36098208.
- [49] Jiang H, Yang Z, Song Z, Green M, Song H, Shao Q.  $\gamma\delta$  T cells in hepatocellular carcinoma patients present cytotoxic activity but are reduced in potency due to IL-2 and IL-21 pathways. *Int Immunopharmacol* 2019;70:167–173. doi:10.1016/j.intimp.2019.02.019, PMID:30802679.
- [50] Liu B, He X, Wang Y, Huang JW, Zheng YB, Li Y, *et al*. Bibliometric Analysis of  $\gamma\delta$  T Cells as Immune Regulators in Cancer Prognosis. *Front Immunol* 2022;13:874640. doi:10.3389/fimmu.2022.874640, PMID:35493488.
- [51] Li T, Zhou Y, Sun X, Bian Y, Wang K, Guo Q, *et al*. Interleukin-2 maintains the survival of interleukin-17 $^+$  gamma/delta T cells in inflammation and autoimmune diseases. *Int Immunopharmacol* 2020;86:106721. doi:10.1016/j.intimp.2020.106721, PMID:32615450.
- [52] Castro F, Cardoso AP, Gonçalves RM, Serre K, Oliveira MJ. Interferon-Gamma at the Crossroads of Tumor Immune Surveillance or Evasion. *Front Immunol* 2018;9:847. doi:10.3389/fimmu.2018.00847, PMID:29780381.
- [53] Cheon H, Wang Y, Wightman SM, Jackson MW, Stark GR. How cancer cells make and respond to interferon-I. *Trends Cancer* 2023;9(1):83–92. doi:10.1016/j.trecan.2022.09.003, PMID:36216730.
- [54] Lança T, Costa MF, Gonçalves-Sousa N, Rei M, Grosso AR, Penido C, *et al*. Protective role of the inflammatory CCR2/CCL2 chemokine pathway through recruitment of type 1 cytotoxic  $\gamma\delta$  T lymphocytes to tumor beds. *J Immunol* 2013;190(12):6673–6680. doi:10.4049/jimmunol.1300434, PMID:23686489.
- [55] Penido C, Costa MF, Souza MC, Costa KA, Candéa AL, Benjamim CF, *et al*. Involvement of CC chemokines in gammadelta T lymphocyte trafficking during allergic inflammation: the role of CCL2/CCR2 pathway. *Int Immunol* 2008;20(1):129–139. doi:10.1093/intimm/20.1, PMID:18056919.
- [56] Wunderlich CM, Ackermann PJ, Ostermann AL, Adams-Quack P, Vogt MC, Tran ML, *et al*. Obesity exacerbates colitis-associated cancer via IL-6-regulated macrophage polarisation and CCL-20/CCR-6-mediated lymphocyte recruitment. *Nat Commun* 2018;9(1):1646. doi:10.1038/s41467-018-03773-0, PMID:29695802.
- [57] Reinhardt A, Ravens S, Fleige H, Haas JD, Oberdörfer L, Łyszkiewicz M, *et al*. CCR7-mediated migration in the thymus controls  $\gamma\delta$  T-cell development. *Eur J Immunol* 2014;44(5):1320–1329. doi:10.1002/eji.201344330, PMID:24500801.
- [58] Moser B, Brandes M. Gammadelta T cells: an alternative type of professional APC. *Trends Immunol* 2006;27(3):112–118. doi:10.1016/j.it.2006.01.002, PMID:16442347.
- [59] Collins C, Wolfe J, Roessner K, Shi C, Sigal LH, Budd RC. Lyme arthritis synovial gammadelta T cells instruct dendritic cells via fas ligand. *J Immunol* 2005;175(9):5656–5665. doi:10.4049/jimmunol.175.9.5656, PMID:16237055.
- [60] Zhu X, Zeng Z, Qiu D, Chen J. V $\gamma$ 9V $\delta$ 2 T cells inhibit immature dendritic cell transdifferentiation into osteoclasts through downregulation of RANK, c-Fos and ATP6V0D2. *Int J Mol Med* 2018;42(4):2071–2079. doi:10.3892/ijmm.2018.3791, PMID:30066839.
- [61] Meuter S, Eberl M, Moser B. Prolonged antigen survival and cytosolic export in cross-presenting human gammadelta T cells. *Proc Natl Acad Sci U S A* 2010;107(19):8730–8735. doi:10.1073/pnas.1002769107, PMID:20413723.
- [62] Daley D, Zambirinis CP, Seifert L, Akkad N, Mohan N, Werba G, *et al*.  $\gamma\delta$  T Cells Support Pancreatic Oncogenesis by Restraining  $\alpha\beta$  T Cell Activation. *Cell* 2016;166(6):1485–1499.e15. doi:10.1016/j.cell.2016.07.046, PMID:27569912.
- [63] Rampoldi F, Ullrich L, Prinz I. Revisiting the Interaction of  $\gamma\delta$  T-Cells and B-Cells. *Cells* 2020;9(3):743. doi:10.3390/cells9030743, PMID:32197382.
- [64] Petrasca A, Melo AM, Breen EP, Doherty DG. Human V $\delta$ 3 $^+$   $\gamma\delta$  T cells induce maturation and IgM secretion by B cells. *Immunol Lett* 2018;196:126–134. doi:10.1016/j.imlet.2018.02.002, PMID:29438730.
- [65] Mathews JA, Kasahara DI, Ribeiro L, Wurmbrand AP, Ninin FM, Shore SA.  $\gamma\delta$  T Cells Are Required for M2 Macrophage Polarization and Resolution of Ozone-Induced Pulmonary Inflammation in Mice. *PLoS One* 2015;10(7):e0131236. doi:10.1371/journal.pone.0131236, PMID:26135595.
- [66] Nguyen CT, Furuya H, Das D, Marusina AI, Merleev AA, Ravindran R, *et al*. Peripheral  $\gamma\delta$  T Cells Regulate Neutrophil Expansion and Recruitment in Experimental Psoriatic Arthritis. *Arthritis Rheumatol* 2022;74(9):1524–1534. doi:10.1002/art.42124, PMID:35320625.
- [67] Bouchareychas L, Grössinger EM, Kang M, Adamopoulos IE.  $\gamma\delta$ TCR regulates production of interleukin-27 by neutrophils and attenuates inflammatory arthritis. *Sci Rep* 2018;8(1):7590. doi:10.1038/s41598-018-25988-3, PMID:29765156.
- [68] Al Mogrampi S, Boumpourea C, Afalonati H, Lagou M, Angelopoulou K, Anastakis D, *et al*. Inhibition of  $\gamma\delta$ -TCR or IL17a Reduces T-Cell and Neutrophil Infiltration after Ischemia/Reperfusion Injury in Mouse Liver. *J Clin Med* 2023;12(5):1751. doi:10.3390/jcm12051751, PMID:36902538.
- [69] Zakeri N, Hall A, Swadling L, Pallett LJ, Schmidt NM, Diniz MO, *et al*. Characterisation and induction of tissue-resident gamma delta T-cells to target hepatocellular carcinoma. *Nat Commun* 2022;13(1):1372. doi:10.1038/s41467-022-29012-1, PMID:35296658.
- [70] Xu Y, Xiang Z, Alnaggar M, Kouakanou L, Li J, He J, *et al*. Allogeneic V $\gamma$ 9V $\delta$ 2 T-cell immunotherapy exhibits promising clinical safety and prolongs the survival of patients with late-stage lung or liver cancer. *Cell Mol Immunol* 2021;18(2):427–439. doi:10.1038/s41423-020-0515-7, PMID:32939032.
- [71] Kang Y, Han M, Kim M, Hwang HJ, Ahn BC, Tak E, *et al*. Cytotoxicity of Human Hepatic Intrahepatic Gamma/Delta T Cells Depends on Phosphoantigen and NK Receptor Signaling. *Anticancer Res* 2023;43(1):63–73. doi:10.21873/anticancer.16135, PMID:36585162.
- [72] Kamimura H, Yamagiwa S, Tsuchiya A, Takamura M, Matsuda Y, Ohkoshi S, *et al*. Reduced NKG2D ligand expression in hepatocellular carcinoma correlates with early recurrence. *J Hepatol* 2012;56(2):381–388. doi:10.1016/j.jhep.2011.06.017, PMID:21756848.
- [73] Kong Y, Cao W, Xi X, Ma C, Cui L, He W. The NKG2D ligand ULBP4 binds to TCRgamma9/delta2 and induces cytotoxicity to tumor cells through both TCRgamma9/delta2 and NKG2D. *Blood* 2009;114(2):310–317. doi:10.1182/blood-2008-12-196287, PMID:19436053.
- [74] Lança T, Correia DV, Moita CF, Raquel H, Neves-Costa A, Ferreira C, *et al*. The MHC class Ib protein ULBP1 is a nonredundant determinant of leukemia/lymphoma susceptibility to gammadelta T-cell cytotoxicity. *Blood* 2010;115(12):2407–2411. doi:10.1182/blood-2009-08-237123, PMID:20101024.
- [75] Cadoux M, Caruso S, Pham S, Gougelet A, Pophillat C, Riou R, *et al*. Expression of NKG2D ligands is downregulated by  $\beta$ -catenin signalling and associates with HCC aggressiveness. *J Hepatol* 2021;74(6):1386–1397. doi:10.1016/j.jhep.2021.01.017, PMID:33484773.
- [76] Rincon-Orozco B, Kunzmann V, Wrobel P, Kabelitz D, Steinle A, Herrmann T. Activation of V gamma 9V delta 2 T cells by NKG2D. *J Immunol* 2005;175(4):2144–2151. doi:10.4049/jimmunol.175.4.2144, PMID:16081780.
- [77] Li P, Du Q, Cao Z, Guo Z, Evankovich J, Yan W, *et al*. Interferon- $\gamma$  induces autophagy with growth inhibition and cell death in human hepatocellular carcinoma (HCC) cells through interferon-regulatory factor-1 (IRF-1). *Cancer Lett* 2012;314(2):213–222. doi:10.1016/j.canlet.2011.09.031, PMID:22056812.
- [78] Jang MK, Kim HS, Chung YH. Clinical aspects of tumor necrosis factor- $\alpha$  signaling in hepatocellular carcinoma. *Curr Pharm Des* 2014;20(17):2799–2808. doi:10.2174/13816128113199990587, PMID:23944370.
- [79] Toutirais O, Cabillic F, Le Fric G, Salot S, Loyer P, Le Gallo M, *et al*. DNAX accessory molecule-1 (CD226) promotes human hepatocellular carcinoma cell lysis by Vgamma9Vdelta2 T cells. *Eur J Immunol* 2009;39(5):1361–1368. doi:10.1002/eji.200838409, PMID:19404979.
- [80] Bruni E, Cimino MM, Donadon M, Carriero R, Terzoli S, Piazza R, *et al*. Intrahepatic CD69(+)V $\delta$ 1 T cells re-circulate in the blood of patients with metastatic colorectal cancer and limit tumor progression. *J Immunother Cancer* 2022;10(7):e004579. doi:10.1136/jitc-2022-004579, PMID:35863820.
- [81] Bertaina A, Zorzoli A, Petretto A, Barbarito G, Inglese E, Merli P, *et al*. Zoledronic acid boosts  $\gamma\delta$  T-cell activity in children receiving  $\alpha\beta$ (+) T and CD19(+) cell-depleted grafts from an HLA-haplo-identical donor. *Oncoimmunology* 2017;6(2):e1216291. doi:10.1080/2162402X.2016.1216291, PMID:28344861.
- [82] Hodgins NO, Al-Jamal WT, Wang JT, Parente-Pereira AC, Liu M, Maher J, *et al*. In vitro potency, in vitro and in vivo efficacy of liposomal alendronate in combination with  $\gamma\delta$  T cell immunotherapy in mice. *J Control Release* 2016;241:229–241. doi:10.1016/j.jconrel.2016.09.023, PMID:27664328.
- [83] Qian P, Zhang YW, Zhou ZH, Liu JQ, Yue SY, Guo XL, *et al*. Artesunate enhances  $\gamma\delta$  T-cell-mediated antitumor activity through augmenting  $\gamma\delta$  T-cell function and reversing immune escape of HepG2 cells. *Immunopharmacol*

- Immunotoxicol 2018;40(2):107–116. doi:10.1080/08923973.2017.1386212, PMID:29405080.
- [84] Patil RS, Shah SU, Shrikhande SV, Goel M, Dikshit RP, Chiplunkar SV. IL17 producing  $\gamma\delta$ T cells induce angiogenesis and are associated with poor survival in gallbladder cancer patients. *Int J Cancer* 2016;139(4):869–881. doi:10.1002/ijc.30134, PMID:27062572.
- [85] Gu FM, Li QL, Gao Q, Jiang JH, Zhu K, Huang XY, *et al*. IL-17 induces AKT-dependent IL-6/JAK2/STAT3 activation and tumor progression in hepatocellular carcinoma. *Mol Cancer* 2011;10:150. doi:10.1186/1476-4598-10-150, PMID:22171994.
- [86] Wakita D, Sumida K, Iwakura Y, Nishikawa H, Ohkuri T, Chamoto K, *et al*. Tumor-infiltrating IL-17-producing gammadelta T cells support the progression of tumor by promoting angiogenesis. *Eur J Immunol* 2010;40(7):1927–1937. doi:10.1002/eji.200940157, PMID:20397212.
- [87] Coffelt SB, Kersten K, Doornebal CW, Weiden J, Vrijland K, Hau CS, *et al*. IL-17-producing  $\gamma\delta$  T cells and neutrophils conspire to promote breast cancer metastasis. *Nature* 2015;522(7556):345–348. doi:10.1038/nature14282, PMID:25822788.
- [88] Edwards SC, Hedley A, Hoevenaer WHM, Wiesheu R, Glauner T, Kilbey A, *et al*. PD-1 and TIM-3 differentially regulate subsets of mouse IL-17A-producing  $\gamma\delta$  T cells. *J Exp Med* 2023;220(2):e20211431. doi:10.1084/jem.20211431, PMID:36480166.
- [89] Barsch M, Salié H, Schlaack AE, Zhang Z, Hess M, Mayer LS, *et al*. T-cell exhaustion and residency dynamics inform clinical outcomes in hepatocellular carcinoma. *J Hepatol* 2022;77(2):397–409. doi:10.1016/j.jhep.2022.02.032, PMID:35367533.
- [90] He W, Hu Y, Chen D, Li Y, Ye D, Zhao Q, *et al*. Hepatocellular carcinoma-infiltrating  $\gamma\delta$  T cells are functionally defected and allogenic V $\delta$ 2(+)  $\gamma\delta$  T cell can be a promising complement. *Clin Transl Med* 2022;12(4):e800. doi:10.1002/ctm2.800, PMID:35390227.
- [91] Ma S, Cheng Q, Cai Y, Gong H, Wu Y, Yu X, *et al*. IL-17A produced by  $\gamma\delta$  T cells promotes tumor growth in hepatocellular carcinoma. *Cancer Res* 2014;74(7):1969–1982. doi:10.1158/0008-5472.CAN-13-2534, PMID:24525743.
- [92] Kühl AA, Pawlowski NN, Grollich K, Blessenohl M, Westermann J, Zeitz M, *et al*. Human peripheral gammadelta T cells possess regulatory potential. *Immunology* 2009;128(4):580–588. doi:10.1111/j.1365-2567.2009.03162.x, PMID:19807790.
- [93] Lafont V, Sanchez F, Laprevotte E, Michaud HA, Gros L, Eliaou JF, *et al*. Plasticity of  $\gamma\delta$  T Cells: Impact on the Anti-Tumor Response. *Front Immunol* 2014;5:622. doi:10.3389/fimmu.2014.00622, PMID:25538706.
- [94] Sabbione F, Gabelloni ML, Ernst G, Gori MS, Salamone G, Oleastro M, *et al*. Neutrophils suppress  $\gamma\delta$  T-cell function. *Eur J Immunol* 2014;44(3):819–830. doi:10.1002/eji.201343664, PMID:24271816.
- [95] Li L, Cao B, Liang X, Lu S, Luo H, Wang Z, *et al*. Microenvironmental oxygen pressure orchestrates an anti- and pro-tumoral  $\gamma\delta$  T cell equilibrium via tumor-derived exosomes. *Oncogene* 2019;38(15):2830–2843. doi:10.1038/s41388-018-0627-z, PMID:30546089.
- [96] Mamedov MR, Vedova S, Freimer JW, Sahu AD, Ramesh A, Arce MM, *et al*. CRISPR screens decode cancer cell pathways that trigger  $\gamma\delta$  T cell detection. *Nature* 2023;621(7977):188–195. doi:10.1038/s41586-023-06482-x, PMID:37648854.
- [97] Torres-Hernandez A, Wang W, Nikiforov Y, Tejada K, Torres L, Kalabin A, *et al*.  $\gamma\delta$  T Cells Promote Steatohepatitis by Orchestrating Innate and Adaptive Immune Programming. *Hepatology* 2020;71(2):477–494. doi:10.1002/hep.30952, PMID:31529720.
- [98] Hu Y, Chen D, Hong M, Liu J, Li Y, Hao J, *et al*. Apoptosis, Pyroptosis, and Ferroptosis Conspiringly Induce Immunosuppressive Hepatocellular Carcinoma Microenvironment and  $\gamma\delta$  T-Cell Imbalance. *Front Immunol* 2022;13:845974. doi:10.3389/fimmu.2022.845974, PMID:35444645.
- [99] Sugai S, Yoshikawa T, Iwama T, Tsuchiya N, Ueda N, Fujinami N, *et al*. Hepatocellular carcinoma cell sensitivity to V $\gamma$ 9V $\delta$ 2 T lymphocyte-mediated killing is increased by zoledronate. *Int J Oncol* 2016;48(5):1794–1804. doi:10.3892/ijo.2016.3403, PMID:26936487.
- [100] Chen Z, Zheng Z, Feng L, Huo Z, Huang L, Fu M, *et al*. Overexpression of miR-382 Sensitizes Hepatocellular Carcinoma Cells to  $\gamma\delta$  T Cells by Inhibiting the Expression of c-FLIP. *Mol Ther Oncolytics* 2020;18:467–475. doi:10.1016/j.omto.2020.07.012, PMID:32953981.
- [101] Seo W, Eun HS, Kim SY, Yi HS, Lee YS, Park SH, *et al*. Exosome-mediated activation of toll-like receptor 3 in stellate cells stimulates interleukin-17 production by  $\gamma\delta$  T cells in liver fibrosis. *Hepatology* 2016;64(2):616–631. doi:10.1002/hep.28644, PMID:27178735.
- [102] Zhou BY, Gong JH, Cai XY, Wang JX, Luo F, Jiang N, *et al*. An imbalance between stellate cells and  $\gamma\delta$ T cells contributes to hepatocellular carcinoma aggressiveness and recurrence. *Hepatol Int* 2019;13(5):631–640. doi:10.1007/s12072-019-09969-w, PMID:31342250.
- [103] Brandes M, Willmann K, Moser B. Professional antigen-presentation function by human gammadelta T Cells. *Science* 2005;309(5732):264–268. doi:10.1126/science.1110267, PMID:15933162.
- [104] Muto M, Baghdadi M, Maekawa R, Wada H, Seino K. Myeloid molecular characteristics of human  $\gamma\delta$  T cells support their acquisition of tumor antigen-presenting capacity. *Cancer Immunol Immunother* 2015;64(8):941–949. doi:10.1007/s00262-015-1700-x, PMID:25904200.
- [105] Sacchi A, Tumino N, Sabatini A, Cimino E, Casetti R, Bordoni V, *et al*. Myeloid-Derived Suppressor Cells Specifically Suppress IFN- $\gamma$  Production and Antitumor Cytotoxic Activity of V $\delta$ 2 T Cells. *Front Immunol* 2018;9:1271. doi:10.3389/fimmu.2018.01271, PMID:29928279.
- [106] Yi Y, He HW, Wang JX, Cai XY, Li YW, Zhou J, *et al*. The functional impairment of HCC-infiltrating  $\gamma\delta$  T cells, partially mediated by regulatory T cells in a TGF $\beta$ - and IL-10-dependent manner. *J Hepatol* 2013;58(5):977–983. doi:10.1016/j.jhep.2012.12.015, PMID:23262246.
- [107] Ye J, Ma C, Hsueh EC, Eickhoff CS, Zhang Y, Varvares MA, *et al*. Tumor-derived  $\gamma\delta$  regulatory T cells suppress innate and adaptive immunity through the induction of immunosenescence. *J Immunol* 2013;190(5):2403–2414. doi:10.4049/jimmunol.1202369, PMID:23355732.
- [108] Peters C, Kabelitz D, Wesch D. Regulatory functions of  $\gamma\delta$  T cells. *Cell Mol Life Sci* 2018;75(12):2125–2135. doi:10.1007/s00018-018-2788-x, PMID:29520421.
- [109] Zhao N, Dang H, Ma L, Martin SP, Forgues M, Ylaja K, *et al*. Intratumoral  $\gamma\delta$  T-Cell Infiltrates, Chemokine (C-C Motif) Ligand 4/Chemokine (C-C Motif) Ligand 5 Protein Expression and Survival in Patients With Hepatocellular Carcinoma. *Hepatology* 2021;73(3):1045–1060. doi:10.1002/hep.31412, PMID:32502310.
- [110] Glatzel A, Wesch D, Schiemann F, Brandt E, Janssen O, Kabelitz D. Patterns of chemokine receptor expression on peripheral blood gamma delta T lymphocytes: strong expression of CCR5 is a selective feature of V delta 2/V gamma 9 gamma delta T cells. *J Immunol* 2002;168(10):4920–4929. doi:10.4049/jimmunol.168.10.4920, PMID:11994442.
- [111] Li X, Yao W, Yuan Y, Chen P, Li B, Li J, *et al*. Targeting of tumour-infiltrating macrophages via CCL2/CCR2 signalling as a therapeutic strategy against hepatocellular carcinoma. *Gut* 2017;66(1):157–167. doi:10.1136/gutjnl-2015-310514, PMID:26452628.
- [112] Ajeubor MN, Jin Y, Gremillion GL, Strieter RM, Chen Q, Adegboyega PA. GammadeltaT cells initiate acute inflammation and injury in adenovirus-infected liver via cytokine-chemokine cross talk. *J Virol* 2008;82(19):9564–9576. doi:10.1128/JVI.00927-08, PMID:18667515.
- [113] Russo RC, Savino B, Mirolo M, Buracchi C, Germano G, Anselmo A, *et al*. The atypical chemokine receptor ACKR2 drives pulmonary fibrosis by tuning influx of CCR2+ and CCR5+ IFN $\gamma$ -producing  $\gamma\delta$ T cells in mice. *Am J Physiol Lung Cell Mol Physiol* 2018;314(6):L1010–L1025. doi:10.1152/ajplung.00233.2017, PMID:29469612.
- [114] Hammerich L, Bangen JM, Govaere O, Zimmermann HW, Gassler N, Huss S, *et al*. Chemokine receptor CCR6-dependent accumulation of  $\gamma\delta$  T cells in injured liver restricts hepatic inflammation and fibrosis. *Hepatology* 2014;59(2):630–642. doi:10.1002/hep.26697, PMID:23959575.
- [115] Hao J, Dong S, Xia S, He W, Jia H, Zhang S, *et al*. Regulatory role of V $\gamma$ 1  $\gamma\delta$  T cells in tumor immunity through IL-4 production. *J Immunol* 2011;187(10):4979–4986. doi:10.4049/jimmunol.1101389, PMID:21987661.
- [116] Ren H, Liu X, Xu Q, Li W, Zhao N. Interleukin-35 expression promotes hepatocellular carcinogenesis by inducing  $\gamma\delta$  T-cell exhaustion. *Genomics* 2023;115(3):110639. doi:10.1016/j.ygeno.2023.110639, PMID:37187255.
- [117] Hoeres T, Holzmann E, Smetak M, Birkmann J, Wilhelm M. PD-1 signaling modulates interferon- $\gamma$  production by Gamma Delta ( $\gamma\delta$ ) T-Cells in response to leukemia. *Oncoimmunology* 2019;8(3):1550618. doi:10.1080/2162402X.2018.1550618, PMID:30723581.
- [118] Sheng Y, Chen K, Jiang W, Wu Z, Zhang W, Jing H, *et al*. PD-1 restrains IL-17A production from  $\gamma\delta$  T cells to modulate acute radiation-induced lung injury. *Transl Lung Cancer Res* 2021;10(2):685–698. doi:10.21037/tlcr-20-838, PMID:33718014.
- [119] Zheng J, Qiu D, Jiang X, Zhao Y, Zhao H, Wu X, *et al*. Increased PD-1(+)Foxp3(+)  $\gamma\delta$  T cells associate with poor overall survival for patients with acute myeloid leukemia. *Front Oncol* 2022;12:1007565. doi:10.3389/fonc.2022.1007565, PMID:36591503.
- [120] Liu Q, Tao YH, Bai RZ, Chang SJ, Hua D. Zoledronic acid inhibits growth of hepatocellular carcinoma cells in vitro and in vivo. *Chin Med J (Engl)* 2013;126(8):1486–1490. doi:10.3760/cma.j.issn.0366-6999.20122385, PMID:23595382.
- [121] Armeanu-Ebinger S, Hoh A, Wenz J, Fuchs J. Targeting EpCAM (CD326) for immunotherapy in hepatoblastoma. *Oncoimmunology* 2013;2(1):e22620. doi:10.4161/onci.22620, PMID:23482411.
- [122] Bruni E, Cazzetta V, Donadon M, Cimino M, Torzilli G, Spata G, *et al*. Chemotherapy accelerates immune-senescence and functional impairments of V $\delta$ 2(pos) T cells in elderly patients affected by liver metastatic colorectal cancer. *J Immunother Cancer* 2019;7(1):347. doi:10.1186/s40425-019-0825-4, PMID:31829255.
- [123] Capsomidis A, Benthall G, Van Acker HH, Fisher J, Kramer AM, Abeln Z, *et al*. Chimeric Antigen Receptor-Engineered Human Gamma Delta T Cells: Enhanced Cytotoxicity with Retention of Cross Presentation. *Mol Ther* 2018;26(2):354–365. doi:10.1016/j.yjmt.2017.12.001, PMID:29310916.
- [124] Makkouk A, Yang XC, Barca T, Lucas A, Turkoz M, Wong JTS, *et al*. Off-the-shelf  $\delta$ 1 gamma delta T cells engineered with glypican-3 (GPC-3)-specific chimeric antigen receptor (CAR) and soluble IL-15 display robust antitumor efficacy against hepatocellular carcinoma. *J Immunother Cancer* 2021;9(12):e003441. doi:10.1136/jitc-2021-003441, PMID:34916256.
- [125] Varon S, Skaper SD, Barbin G, Selak I, Manthorpe M. Low molecular weight agents support survival of cultured neurons from the central nervous system. *J Neurosci* 1984;4(3):654–658. doi:10.1186/s40425-019-0501-8, PMID:6368763.
- [126] Zhang T, Chen J, Niu L, Liu Y, Ye G, Jiang M, *et al*. Clinical Safety and Efficacy of Locoregional Therapy Combined with Adoptive Transfer of Allogeneic  $\gamma\delta$  T Cells for Advanced Hepatocellular Carcinoma and Intrahepatic Cholangiocarcinoma. *J Vasc Interv Radiol* 2022;33(1):19–27.e3. doi:10.1016/j.jvir.2021.09.012, PMID:34600129.
- [127] Tian W, Ma J, Shi R, Ren C, He J, Zhao H.  $\gamma\delta$  T cell-mediated individualized immunotherapy for hepatocellular carcinoma considering clinicopathological characteristics and immunosuppressive factors. *Oncol Lett* 2018;15(4):5433–5442. doi:10.3892/ol.2018.8026, PMID:29552184.
- [128] Cabillit F, Toutirais O, Lavoué V, de La Pintièrre CT, Daniel P, Rioux-Leclerc N, *et al*. Aminobisphosphonate-pretreated dendritic cells trigger successful Vgamma9Vdelta2 T cell amplification for immunotherapy in advanced cancer patients. *Cancer Immunol Immunother* 2010;59(11):1611–1619.

- doi:10.1007/s00262-010-0887-0, PMID:20582413.
- [129] Fusco V, Campisi G, Bedogni A. One changing and challenging scenario: the treatment of cancer patients with bone metastases by bisphosphonates and denosumab, the cost-benefit evaluation of different options, and the risk of medication-related osteonecrosis of the jaw (MRONJ). *Support Care Cancer* 2022;30(9):7047–7051. doi:10.1007/s00520-022-06982-y, PMID:35312858.
- [130] Tuesley KM, Webb PM, Protani MM, Spilsbury K, Pearson SA, Coory MD, *et al*. Nitrogen-based Bisphosphonate Use and Ovarian Cancer Risk in Women Aged 50 Years and Older. *J Natl Cancer Inst* 2022;114(6):878–884. doi:10.1093/jnci/djac050, PMID:35262727.
- [131] Wei Z, Pan B, Jia D, Yu Y. Long-term safety and efficacy of bisphosphonate therapy in advanced lung cancer with bone metastasis. *Future Oncol* 2022;18(18):2257–2267. doi:10.2217/fon-2022-0098, PMID:35414201.
- [132] Sorscher S. Duration and Dose of Adjuvant Zoledronic Acid for Treatment of Early Breast Cancer. *JAMA Oncol* 2022;8(1):171. doi:10.1001/jamaoncol.2021.5765, PMID:34792535.
- [133] Bouet-Toussaint F, Cabillic F, Toutirais O, Le Gallo M, Thomas de la Pintièrre C, Daniel P, *et al*. Vgamma9Vdelta2 T cell-mediated recognition of human solid tumors. Potential for immunotherapy of hepatocellular and colorectal carcinomas. *Cancer Immunol Immunother* 2008;57(4):531–539. doi:10.1007/s00262-007-0391-3, PMID:17764010.
- [134] Ferry GM, Agbuduwe C, Forrester M, Dunlop S, Chester K, Fisher J, *et al*. A Simple and Robust Single-Step Method for CAR-V $\delta$ 1  $\gamma\delta$ T Cell Expansion and Transduction for Cancer Immunotherapy. *Front Immunol* 2022;13:863155. doi:10.3389/fimmu.2022.863155, PMID:35711450.
- [135] Rozenbaum M, Meir A, Aharoni Y, Itzhaki O, Schachter J, Bank I, *et al*. Gamma-Delta CAR-T Cells Show CAR-Directed and Independent Activity Against Leukemia. *Front Immunol* 2020;11:1347. doi:10.3389/fimmu.2020.01347, PMID:32714329.
- [136] Ganapathy T, Radhakrishnan R, Sakshi S, Martin S. CAR  $\gamma\delta$  T cells for cancer immunotherapy. Is the field more yellow than green? *Cancer Immunol Immunother* 2023;72(2):277–286. doi:10.1007/s00262-022-03260-y, PMID:35960333.
- [137] Nishimoto KP, Barca T, Azameera A, Makkouk A, Romero JM, Bai L, *et al*. Allogeneic CD20-targeted  $\gamma\delta$  T cells exhibit innate and adaptive antitumor activities in preclinical B-cell lymphoma models. *Clin Transl Immunology* 2022;11(2):e1373. doi:10.1002/cti2.1373, PMID:35136603.
- [138] Sun R, Li J, Lin X, Yang Y, Liu B, Lan T, *et al*. Peripheral immune characteristics of hepatitis B virus-related hepatocellular carcinoma. *Front Immunol* 2023;14:1079495. doi:10.3389/fimmu.2023.1079495, PMID:37077908.