



Review Article



The Role of Mitochondrial Complexes in Liver Diseases

Hai An*

Academy of Chinese Medical Sciences, Zhejiang Chinese Medical University, Hangzhou, Zhejiang, China

Received: April 28, 2025 | Revised: August 14, 2025 | Accepted: September 05, 2025 | Published online: October 10, 2025

Abstract

Mitochondrial respiratory complexes (Complexes I–V) and their assembly into respiratory supercomplexes (SCs) are fundamental to liver bioenergetics, redox homeostasis, and metabolic adaptability. Disruption of these systems contributes to major liver diseases, including non-alcoholic fatty liver disease, alcoholic liver disease, drug-induced liver injury, viral hepatitis, and hepatocellular carcinoma, by impairing adenosine triphosphate synthesis, increasing oxidative stress, and altering metabolic pathways. Recent advances have clarified the structural-functional interdependence of individual complexes within SCs, revealing their dynamic remodeling in response to physiological stress and pathological injury. These insights open opportunities for clinical translation, such as targeting SC stability with pharmacological agents, nutritional strategies, or gene therapy, and employing mitochondrial transplantation in cases of severe mitochondrial failure. Precision medicine approaches, incorporating multi-omics profiling and patient-derived models, may enable individualized interventions and early detection using SC integrity as a biomarker. By linking molecular mechanisms to therapeutic strategies, this review underscores the potential of mitochondrial-targeted interventions to improve outcomes in patients with liver disease.

Citation of this article: An H. The Role of Mitochondrial Complexes in Liver Diseases. *J Clin Transl Hepatol* 2025. doi: 10.14218/JCTH.2025.00194.

Introduction

The liver, a vital organ responsible for metabolism and detoxification, is highly susceptible to damage from prolonged exposure to harmful substances such as alcohol, drugs, and toxins.¹ These agents induce oxidative stress by increasing intracellular reactive oxygen species (ROS) levels, which subsequently damage cell membranes, proteins, and DNA, ultimately leading to hepatocyte apoptosis or necrosis.² Mitochondrial dysfunction is a key factor in the onset and progression of liver injury. As the central hub of intracellular energy metabolism, mitochondria, when impaired, can cause reduced adenosine triphosphate (ATP) production,

exacerbated oxidative stress, and increased apoptosis. This disorder may be associated with mitochondrial DNA (mtDNA) mutations, protein synthesis defects, or decreased membrane potential.³ Emerging evidence shows that mitochondrial complexes often assemble into respiratory supercomplexes (SCs), which enhance electron transfer, reduce ROS generation, and stabilize individual complexes.⁴ Disruption of SCs has been increasingly recognized in liver diseases, highlighting the importance of mitochondrial structural integrity. An in-depth investigation of these mechanisms can provide insights into the pathogenesis of liver diseases and guide the development of novel therapeutic strategies.

Functions and biological significance of mitochondrial complexes

Mitochondrial complexes reside within the inner mitochondrial membrane and comprise five core complexes (I–V). These complexes function synergistically to transfer electrons from nicotinamide adenine dinucleotide (NAD) and flavin adenine dinucleotide (FAD) to oxygen, ultimately producing ATP to meet cellular energy demands. Each complex consists of specific subunits and carries out distinct redox reactions (Table 1).^{5–27}

Mitochondrial complex I

Mitochondrial complex I (NADH dehydrogenase) is the first enzyme complex in the mitochondrial electron transport chain (ETC) and the largest and most structurally intricate component of the chain. It facilitates electron transfer from NADH to coenzyme Q (CoQ) while concurrently pumping four protons into the mitochondrial intermembrane space, thereby establishing a proton gradient across the inner mitochondrial membrane. This gradient drives ATP synthase (complex V), a critical step in ATP production.^{28,29} In liver diseases, reduced complex I activity disrupts the NADH/NAD⁺ ratio, leading to redox imbalance, oxidative stress, and lipid peroxidation, which ultimately exacerbates hepatocyte injury.³⁰ Furthermore, complex I dysfunction undermines the integrity of the mitochondrial respiratory chain, reduces ATP synthesis, and aggravates metabolic dysregulation, intensifying liver damage.³¹ Complex I dysfunction is a key contributor to multiple liver diseases. In alcoholic liver disease and drug-induced liver injury (DILI), reduced complex I activity impairs mitochondrial electron transfer, increases ROS accumulation, and intensifies lipid peroxidation, leading to hepatocyte injury.^{32,33} In non-alcoholic fatty liver disease (NAFLD), complex I dysfunction is strongly linked to impaired fatty acid β -oxidation and mitochondrial dysfunction, serving as a critical driver of disease progression.^{34,35} In cirrhosis,

Keywords: Liver disease; Mitochondrial complex; Energy metabolism; Adenosine triphosphate; ATP; ATP production; Electron transfer chain.

***Correspondence to:** Hai An, Academy of Chinese Medical Sciences, Zhejiang Chinese Medical University, 260 Baichuan Street, Fuyang District, Hangzhou, Zhejiang 311402, China. ORCID: <https://orcid.org/000-0002-6634-0855>. Tel: +86-15619270193, E-mail: hai_zcmu@zcmu.edu.cn.

Table 1. Overview of mitochondrial complexes I–V in liver diseases

Complex	Associated liver diseases	Clinical relevance / Therapeutic strategy
Complex I	ALD, ⁵ NAFLD, ⁶ cirrhosis, HCC ⁷	NAD ⁺ precursors, ⁸ CoQ10, ⁹ antioxidants, metformin
Complex II	HCC, ¹⁰ metabolic liver disease ¹¹	SDHIs (inhibitors or modulators), ¹² metabolic control of succinate ¹³
Complex III	NAFLD, ¹⁴ IRI, ¹⁵ DILI ¹⁶	CoQ10 supplementation, NAD ⁺ boosting, ¹⁷ ROS scavengers ¹⁸
Complex IV	Fibrosis, ¹⁹ viral hepatitis, ²⁰ IRI ²¹	Oxygen delivery modulation, ²² copper cofactors, ²³ NO donors ²⁴
Complex V	Hepatitis, ²⁵ ALD ²⁶	Creatine, pyruvate supplementation, mitoprotective peptides ²⁷

ALD, alcoholic liver disease; HCC, hepatocellular carcinoma; NAFLD, non-alcoholic fatty liver disease; NAD, nicotinamide adenine dinucleotide; CoQ, coenzyme Q; SDHIs, succinate dehydrogenase inhibitors; IRI, ischemia-reperfusion injury; DILI, drug induced-liver injury; ROS, reactive oxygen species; NO, nitric oxide.

reduced complex I activity worsens oxidative stress, further compromising hepatocyte function and accelerating disease progression.³⁶ In hepatocellular carcinoma, complex I dysfunction may induce redox imbalance, promoting tumor cell proliferation and metastasis.³⁷ Viral infections may also directly or indirectly affect complex I function. For example, hepatitis C virus infection has been shown to cause abnormal mitochondrial function, including decreased complex I activity and increased ROS production.³⁸ These findings highlight the essential role of mitochondrial complex I in the initiation and progression of liver diseases.

Mitochondrial complex II

Mitochondrial complex II (succinate dehydrogenase, SDH) is a nuclear-encoded enzyme that bridges the tricarboxylic acid (TCA) cycle and the ETC.³⁹ Unlike other ETC complexes, complex II does not translocate protons across the mitochondrial membrane. Instead, it catalyzes the oxidation of succinate to fumarate, reducing FAD to FADH₂ and transferring electrons to CoQ.⁴⁰ This provides an alternative pathway for electrons to bypass complex I and enter the ETC, ensuring continuity of energy metabolism.⁴¹ As a critical link between the TCA cycle and the ETC, complex II plays a key role in maintaining mitochondrial metabolic stability and cellular bioenergetic balance, particularly when NADH oxidation is impaired or its supply is insufficient. By mediating succinate metabolism, complex II integrates the TCA cycle, fatty acid oxidation, and the ETC, ensuring efficient utilization of carbon substrates and sustaining ATP synthesis.⁴² However, long-term or excessive use of SDH inhibitors may lead to abnormal accumulation of succinate, disrupting mitochondrial oxidative metabolism. This effect is especially pronounced in metabolically susceptible individuals and may result in severe adverse consequences.⁴⁰ Notably, while insufficient SDH activity can impair mitochondrial function, excessive SDH activity may also induce metabolic imbalances. Therefore, appropriate regulation of complex II activity may represent a potential strategy for treating metabolic disorders.⁴³ Dysfunction of complex II is linked not only to hereditary mitochondrial diseases but also to the pathogenesis of cancer, neurodegenerative disorders, and metabolic syndromes. Research indicates that SDH functions as a tumor suppressor, whereas its metabolite, succinate, may act as an oncometabolite promoting tumor progression under specific circumstances.^{44,45} In HCC tumor specimens, succinate dehydrogenase subunit B (SDHB) expression is reduced by 40%.⁴⁶ Reduced SDHB expression disrupts mitochondrial ETC function, leading to the Warburg effect. This metabolic reprogramming promotes oncogenic processes and strengthens the malignant phenotype of HCC cells.⁴⁷ Such a metabolic shift may represent an adaptive strategy employed by tumor cells to evade immune detection and enhance resistance to apoptosis.⁴⁸

Mitochondrial complex III

Mitochondrial complex III (cytochrome bc₁ complex) is a crucial component of the ETC, responsible for transferring electrons from CoQ to cytochrome c while actively pumping protons into the intermembrane space. This process generates the proton gradient essential for ATP synthesis.^{49,50} When complex III activity decreases, electron transfer efficiency declines, leading to a significant increase in oxidative stress and worsening mitochondrial damage.⁵¹ Given that the liver is one of the most metabolically active organs, its mitochondria are highly sensitive to oxidative stress. Impaired complex III function results in excessive mitochondrial ROS generation, which damages cellular membranes, proteins, and DNA, ultimately causing hepatocyte injury and inflammation.⁵² Additionally, complex III dysfunction is closely associated with energy metabolism disorders in liver diseases. Patients often exhibit mitochondrial dysfunction, leading to reduced ATP synthesis, and a further decline in complex III activity exacerbates energy insufficiency, disrupting normal hepatocyte physiological functions.⁵³ Strategies to enhance mitochondrial function, such as supplementation with NAD⁺ precursors (e.g., nicotinamide riboside) or CoQ, can help maintain complex III activity and improve liver metabolic function.⁵⁴ Notably, complex III dysfunction not only affects energy metabolism but also worsens liver injury by promoting apoptosis. Reduced complex III activity increases electron leakage, resulting in cytochrome c release, activation of caspase cascades, and ultimately triggering apoptosis. In liver diseases, apoptosis is a key driver of hepatocyte injury, fibrosis, and disease progression.⁵⁵ Furthermore, complex III dysfunction may contribute to the development and progression of liver tumors. Studies indicate that decreased complex III activity is closely related to tumor cell proliferation, invasion, and metastasis, potentially promoting liver cancer development by regulating metabolic reprogramming, redox balance, and signal transduction pathways.⁵⁶

Mitochondrial complex IV

Mitochondrial complex IV (cytochrome c oxidase, COX) serves as the final enzyme in the ETC, facilitating electron transfer to oxygen to form water. Simultaneously, it pumps protons into the intermembrane space, sustaining the proton gradient required for ATP synthesis.⁵⁷ COX consists of multiple subunits, among which the catalytic subunit containing copper ions plays a crucial role in electron transfer.⁵⁸ COX activity is pivotal in regulating mitochondrial oxidative phosphorylation (OXPHOS) and cellular energy homeostasis, and its dysfunction is implicated in multiple liver pathologies. A decrease in COX activity can impair OXPHOS in hepatocytes, reduce ATP synthesis, disrupt intracellular energy metabolism, and exacerbate liver injury.^{59,60} During liver fibrosis, reduced COX activity may lead to abnormal extracellular matrix deposition, accelerating pathological progression.⁶¹ Elevated

oxidative stress is closely associated with COX dysfunction. Excessive ROS accumulation not only damages hepatocytes but may further disrupt mitochondrial membrane potential, leading to mitochondrial dysfunction.⁶² In liver cancer, abnormal COX activity may enable cancer cells to adapt to hypoxic environments, alter metabolic pathways, and consequently influence tumor growth, invasion, and metastasis.⁶³ In hepatic ischemia-reperfusion injury (IRI), decreased COX activity exacerbates mitochondrial damage, rendering hepatocytes more susceptible to apoptosis or necrosis and worsening tissue injury.⁶⁴ In drug-induced DILI, doxorubicin can significantly inhibit complex IV activity, disrupting cytochrome c redox balance, promoting activation of apoptosis-related markers such as caspase 3, and ultimately triggering cellular energy metabolism disorders and apoptosis.⁶⁵

Mitochondrial complex V

Mitochondrial complex V, or ATP synthase, represents the final enzyme of the OXPHOS system. It harnesses the proton gradient generated by the ETC to drive ATP synthesis from adenosine diphosphate and inorganic phosphate, fulfilling cellular energy demands.⁶⁶ In numerous liver diseases, a decline in complex V activity disrupts ATP synthesis, compromising cellular energy supply and exacerbating hepatocyte dysfunction. In patients with chronic hepatitis or liver cirrhosis, reduced ATP production weakens hepatocyte resilience, making cells more susceptible to pathological damage.^{61,67} Moreover, decreased ATP levels reduce the cell's antioxidant capacity, impairing free radical scavenging and exacerbating oxidative stress, further impairing mitochondrial function and aggravating liver damage.⁶⁸ Complex V dysfunction also closely relates to apoptosis. Reduced complex V activity can alter mitochondrial membrane potential, activating apoptotic signaling pathways, such as the caspase cascade, ultimately inducing programmed cell death in hepatocytes.⁶⁹ This mechanism is particularly prominent in liver cancer and cirrhosis, suggesting that abnormal complex V activity contributes significantly to disease onset and progression. Additionally, complex V dysfunction may interfere with autophagy, an essential mechanism for maintaining cellular homeostasis. Autophagy plays a crucial role in clearing damaged mitochondria and regulating energy balance. When complex V activity is impaired, autophagic flux may be inhibited, leading to the accumulation of damaged organelles and further exacerbating hepatocyte dysfunction.⁷⁰

Structural and functional interdependence of mitochondrial complexes

Mitochondrial OXPHOS has traditionally been understood as a linear sequence of electron transfers through five separate complexes (I–V) embedded in the inner mitochondrial membrane. However, emerging structural and biochemical evidence has revised this view, showing that ETC complexes do not function as isolated units but rather as organized respiratory SCs, often referred to as “respirasomes”.⁷¹ These supramolecular assemblies, mainly consisting of complex I, complex III, and complex IV, facilitate more efficient electron transport, limit ROS production, and stabilize individual complex subunits.⁷²

Structure and composition of respiratory SCs

Respiratory SCs are dynamic assemblies primarily composed of complex I, complex III dimer (III₂), and complex IV in various stoichiometries. The most common mammalian respirasome identified through cryo-electron microscopy is the com-




Fig. 1. The structure of mitochondrial supercomplexes. CI, complex I; CIII₂, complex III dimer; CIV, complex IV.

plex I–III₂–IV.⁷³ These assemblies are stabilized by specific lipids, especially cardiolipin, and accessory proteins such as COX7A2L (SCAF1), which anchor complex IV to complex III₂ and modulate SC stability (Fig. 1).⁷⁴ The formation of SCs optimizes spatial proximity among ETC components, facilitating direct substrate channeling between complexes, minimizing electron leakage, and preventing excess ROS formation, particularly important in metabolically active organs such as the liver. In liver diseases, including IRI and NAFLD, disruption of SC assembly has been reported, further implicating mitochondrial architecture in disease pathogenesis and highlighting potential targets for therapeutic stabilization.^{75,76}

Significance and dependency of respiratory SCs in liver disease

In the liver, the exceptionally high metabolic demand requires efficient electron transport and ATP synthesis. Respiratory SCs enhance mitochondrial efficiency by facilitating electron flux, optimizing proton pumping, and minimizing ROS leakage. Evidence from mouse hepatocyte studies demonstrates that SC destabilization is closely associated with mitochondrial dysfunction, ATP depletion, and elevated ROS production, processes central to the pathogenesis of hepatic disorders.^{77,78} In pathological conditions such as NASH and DILI, oxidative damage or depletion of cardiolipin can compromise SC integrity, leading to the disassembly of complex I and a subsequent rise in oxidative stress.^{79,80} These structural disruptions are further associated with metabolic inflexibility and impaired adaptive responses to nutrient overload or oxidative insults, hallmark features of metabolic liver diseases.

The stability and functional integrity of individual ETC complexes are highly dependent on their incorporation into SCs. Complex I, for example, is inherently unstable in isolation, with degradation markedly accelerated when not embedded within an SC. Conversely, SC assembly promotes the biogenesis and stabilization of complexes III and IV, highlighting a mutual structural dependency that may function as a mitochondrial quality control mechanism.^{81,82} This intricate interdependence means that dysfunction in one complex, such as mutations in complex III, can trigger secondary destabilization of complex I or IV, creating a cascade of mitochondrial defects. Such interconnected failure mechanisms may account for the non-linear and disproportionate phenotypic manifestations observed in certain mitochondrial pathologies and liver diseases.

Table 2. The liver model, complex effects, and intervention strategies applied in research

Liver disease model	Cell/Animal type	Targeted complex
NAFLD ¹⁰⁴	HFD-fed mice; AML12 cells	I, III, IV
DILI ^{38,105}	APAP-treated mice; HepG2 cells	I, III
IRI ¹⁰⁶	Ischemia-reperfusion rat model	III, IV
HCC ¹⁰⁷	HepG2, mouse xenografts	II, V
Cholestatic liver disease ^{108,109}	Bile duct ligation mice; HepaRG cells	I, III, V
Cirrhosis ^{110,111}	CCl ₄ model in rats	I, V

NAFLD, non-alcoholic fatty liver disease; HFD, high-fat diet; I-V, complex I-V; DILI, drug induced-liver injury; APAP, acetaminophen; IRI, ischemia-reperfusion injury; HCC, hepatocellular carcinoma; CCl₄, carbon tetrachloride.

Supercomplex dynamics, disruption, and therapeutic potential

The architecture of respiratory SCs is highly dynamic, undergoing modulation in response to physiological conditions such as nutritional state, oxygen availability, and metabolic stress.⁸³ In the early stages of liver disease, including simple steatosis, hepatocytes may reorganize SCs as a compensatory adaptation to maintain ATP production. However, persistent insults, such as chronic injury or lipid overload, progressively disrupt SC structure, leading to impaired respiratory efficiency and mitochondrial dysfunction.^{84,85} For instance, in fatty liver, prolonged ethanol exposure impairs cardiolipin remodeling enzymes such as Tafazzin, resulting in SC disintegration and mitochondrial fragmentation.^{86,87} Similarly, in NASH models, high-fat diets hinder SC assembly, exacerbating hepatic inflammation and fibrosis.⁸⁸

Understanding the structural-functional coupling within SCs opens new avenues for therapeutic intervention, as stabilizing these assemblies offers a strategy to restore mitochondrial function without directly modifying individual complex activities. Potential approaches include: cardiolipin-stabilizing agents such as the SS-31 peptide, which maintain SC cohesion and reduce ROS production;⁸⁹ nutritional interventions, including ketogenic diets and omega-3 fatty acids, which enhance SC assembly;^{90,91} and gene therapies aimed at restoring essential SC assembly factors such as SCAF1 in genetically susceptible individuals.⁹² Furthermore, SC integrity holds potential as a biomarker for mitochondrial health, enabling early detection of mitochondrial dysfunction in chronic liver diseases. Cutting-edge imaging and proteomics technologies now permit accurate quantification of SC abundance and configuration in liver biopsies and patient-derived organoids, bridging mechanistic insights and clinical translation.^{93,94}

Mitochondrial complex alterations in liver injury

The pathophysiological progression of hepatic injury is closely linked to dynamic alterations in mitochondrial complex functionality. During early injury phases, transient enhancement of mitochondrial complex activity serves as a compensatory response to heightened metabolic demands. However, this adaptive mechanism progressively deteriorates with disease advancement, ultimately leading to bioenergetic failure through impaired ATP synthesis and disrupted cellular energy regulation.^{95,96}

The cascade of mitochondrial complex dysfunction during hepatic injury unfolds through multiple mechanisms. Primarily, depolarization of the mitochondrial membrane potential emerges as an early sentinel event, disrupting electrochemical gradients and directly compromising ETC efficiency and

OXPHOS capacity.⁹⁷ Concurrently, marked attenuation of respiratory chain activity, particularly in complexes I and III, is linked to oxidative post-translational modifications of protein subunits, impaired cofactor biogenesis, and progressive mtDNA mutation accumulation.^{56,98} As injury progresses, the resulting oxidative stress within the mitochondrial matrix exerts a triple pathogenic impact: excessive ROS generation overwhelms endogenous antioxidant defenses, induces phospholipid membrane peroxidation and protein carbonylation, and directly destabilizes mitochondrial genomic integrity.³⁵ At irreversible stages, maladaptive rewiring of mitochondrial quality control manifests through hyperactivated mitophagic flux and dysregulated apoptotic signaling, collectively driving cristae collapse and depletion of bioenergetic reserves.^{99,100} Ultimately, exponential escalation of mtDNA mutational burden establishes a self-reinforcing pathological cycle, in which disrupted respiratory SC assembly and defective synthesis of 13 ETC-essential polypeptides collectively dismantle the mitochondrial bioenergetic network.^{101–103} These interlocking mechanisms, from molecular perturbations to organelle-level failure, orchestrate a systematic progression toward mitochondrial functional collapse, exemplifying the hierarchical integration of redox imbalance and systemic energetic catastrophe.

Collectively, hepatic injury is fundamentally characterized by progressive deterioration of mitochondrial complex structure and function. These multidimensional alterations exhibit dual significance: first, the extent of mitochondrial damage directly correlates with disease severity; second, it exacerbates maladaptive progression by compromising energy transduction efficiency in the OXPHOS system and amplifying stress signaling cascades, such as mitochondrial-derived ROS. Crucially, mitochondrial complex dysfunction not only serves as an early diagnostic biomarker but also acts as a pathogenic catalyst by disrupting respiratory SC assembly and promoting mtDNA mutation accumulation, establishing a self-perpetuating “damage–dysfunction–reinforced damage” positive feedback loop that mechanistically links bioenergetic failure to hepatic deterioration.

From mechanisms to therapy: Targeting mitochondrial complexes in liver disease

Mitochondrial dysfunction has emerged as a central mechanism underlying various forms of liver injury, including NAFLD, DILI, IRI, HCC, and cirrhosis (Table 2).^{38,104–111} Among mitochondrial components, complexes I–V serve not only as key bioenergetic modules but also as regulators of redox signaling, apoptosis, and cellular adaptation. Therefore, therapeutic targeting of mitochondrial complexes represents a promising avenue for disease modification. In recent years, strategies ranging from pharmacological modulation and

metabolic supplementation to organelle transplantation and gene therapy have been explored. Translating these interventions from mechanistic insight to clinical practice requires careful evaluation of their feasibility, delivery methods, and compatibility with individualized medicine.

Mitochondrial transfer therapy: A novel frontier

Mitochondrial transfer therapy, defined as the exogenous transplantation of functional mitochondria into damaged cells, has emerged as a compelling therapeutic strategy to restore bioenergetic function and cellular viability.^{112–114} In the liver, where energy metabolism is tightly coupled with detoxification, protein synthesis, and immune regulation, mitochondrial transfer may provide rapid bioenergetic rescue in acute injury settings such as IRI, acetaminophen overdose, or fulminant hepatic failure.

Several experimental models have demonstrated the efficacy of mitochondrial transfer in reversing hepatic mitochondrial dysfunction. For instance, injection of intact mitochondria isolated from healthy hepatocytes into damaged liver tissues restored ATP production, reduced oxidative stress, and prevented hepatocyte death.^{115,116} In a mouse model of acetaminophen-induced liver injury, mitochondrial transplantation significantly improved survival and reduced serum alanine aminotransferase/aspartate aminotransferase levels, demonstrating hepatoprotective effects.¹¹⁷ Mechanistically, the internalized mitochondria were shown to integrate with host mitochondrial networks, restore membrane potential, and reinitiate mitochondrial biogenesis via PGC-1 α and TFAM activation.^{118–120}

Despite promising preclinical outcomes, several translational challenges remain. Firstly, the delivery route must ensure mitochondrial viability and tissue-specific targeting. While intravenous administration has shown effectiveness in animal models, its safety and efficacy in humans require clinical validation.^{121–124} Secondly, donor mitochondrial source, immunogenicity, and quality control must be standardized to minimize rejection or off-target effects.^{125,126} Thirdly, optimal dosing regimens and timing post-injury are yet to be defined.¹²⁷ Currently, early-phase clinical trials evaluating autologous mitochondrial transfer in cardiac and renal ischemia are ongoing, potentially laying the groundwork for future hepatology applications.^{128,129}

Pharmacological and precision medicine for mitochondrial dysfunction

Modulating the activity of ETC complexes through pharmacological agents or nutritional supplements is an active area of investigation, with growing evidence supporting their role in mitigating mitochondrial dysfunction in liver diseases.^{59,130} NAD⁺ precursors, such as nicotinamide riboside and nicotinamide mononucleotide, have shown potential in restoring complex I function and enhancing OXPHOS, particularly in NAFLD and aging-related hepatic disorders. Ongoing clinical trials (NCT04571008) are evaluating their effects on liver fat content and mitochondrial health in humans.¹³¹ Coenzyme Q10, an essential electron carrier between complexes I/II and III, has been clinically assessed for its hepatoprotective properties, demonstrating improvements in liver function markers and oxidative stress parameters in NAFLD patients. However, optimal dosing regimens and long-term safety remain under investigation.¹³² Other promising modulators include resveratrol, which activates the SIRT1/PGC-1 α axis to promote mitochondrial biogenesis and alleviate hepatic steatosis, and creatine, which stabilizes ATP buffering and supports complex V function.^{133,134}

Despite these advances, clinical efficacy is often limited by poor bioavailability, insufficient tissue specificity, and inter-individual variability.^{135–137} Overcoming these barriers will require the development of advanced drug delivery systems, including liver-targeted nanoparticles for mitochondrial delivery, mitochondria-penetrating peptides capable of traversing the inner membrane, and prodrugs selectively activated within the oxidative environment of damaged hepatocytes.

Given the heterogeneity of liver diseases, precision medicine approaches are critical for effective and individualized interventions.¹³⁸ Mitochondrial complex defects in NAFLD may differ fundamentally from those in HCC or cholestatic liver diseases, necessitating targeted therapeutic strategies.¹³⁹ Integrating multi-omics profiling, including transcriptomics, metabolomics, and proteomics, enables identification of complex-specific dysfunction signatures and stratification of patient subgroups most likely to benefit from targeted therapy. In parallel, genomic sequencing and bioinformatics tools can uncover mutations or polymorphisms in ETC subunits, such as SDHB in HCC or ATP5B in metabolic liver disease, which may serve as biomarkers for therapeutic selection and prognosis prediction. Patient-derived organoids and induced pluripotent stem cell-based liver models offer translational platforms for preclinical testing of mitochondrial-targeted interventions in a personalized manner. Ultimately, clinical feasibility will depend on practical considerations, including the ability to titrate dosing based on mitochondrial function, ensuring long-term safety, particularly regarding off-target ROS generation, developing combination regimens that address both mitochondrial dysfunction and inflammation, and achieving cost-effectiveness and scalability, especially in resource-limited settings.

Emerging insights into mitochondrial complexes in liver disease

Recent advances in mitochondrial biology have significantly enhanced our understanding of mitochondrial complex dysregulation in hepatic pathophysiology. Accumulating evidence demonstrates that dynamic alterations in mitochondrial complex activity strongly correlate with histological grading of liver injury. Suppression of mitochondrial complex function during hepatic damage directly compromises ATP biosynthesis, establishing a causal relationship between bioenergetic failure and disrupted cellular metabolism. Notably, in NAFLD, impairment of mitochondrial complexes induces redox imbalance through defective electron transport, creating a self-perpetuating cycle of oxidative damage and hepatocyte degeneration.¹⁴⁰

Critical pathophysiological insights emerge from investigations into mitochondrial enzyme stoichiometry. Quantitative proteomic analyses reveal marked dysregulation of core respiratory components, particularly complexes I and IV, across various liver pathologies. In HCC, downregulation of complex I and IV subunits correlate with diminished ETC efficiency, creating a metabolic environment that promotes tumor proliferation and metastatic progression.^{141,142}

The redox environment of the mitochondrial ETC has emerged as a key determinant of disease pathogenesis. Hepatic disorders are consistently associated with disruption of redox homeostasis within cristae compartments, significantly affecting ETC function. During fibrotic remodeling, disruption of redox balance across mitochondrial complexes drives excessive superoxide leakage, triggering ROS-mediated amplification of parenchymal injury via lipid peroxidation cascades.¹⁴³

Of particular therapeutic relevance are the regulatory

networks governing mitochondrial complex dynamics. Multiple evolutionarily conserved pathways, including PI3K/Akt, mTOR, and AMPK, have been identified as master regulators of respiratory complex biogenesis and turnover. In HCC models, constitutive activation of PI3K/Akt signaling induces proteasomal degradation of complex I subunits through phosphorylation-dependent ubiquitination, thereby reprogramming cellular metabolism to support neoplastic growth and invasion.^{144–146}

Therapeutic prospects of targeting mitochondrial complexes

Targeting mitochondrial complexes has emerged as a promising therapeutic strategy for liver diseases. As central components of the ETC, mitochondrial complexes demonstrate functional impairments that are mechanistically associated with hepatic pathogenesis. Interventions directed at these complexes can correct metabolic abnormalities at their biochemical source, thereby restoring hepatic functionality. For instance, CRISPR/Cas9-mediated gene editing enables precise repair of pathogenic mtDNA mutations, effectively reversing complex dysfunction and alleviating disease symptoms.^{147,148}

Recent progress in molecular biology has accelerated the development of mitochondrial complex-targeted therapies. Specific antioxidants targeting complex I components (NADH dehydrogenase and CoQ) effectively neutralize ROS in hepatocytes, attenuating oxidative damage.^{149–153} Pharmacological agents such as rofecoxib enhance complex I activity, demonstrating therapeutic potential for metabolic liver disorders.^{154,155} Inhibitors of complex III reduce cellular redox potential, suppressing tumor progression,¹⁵⁶ while moderate complex III inhibition decreases oxidative stress in hepatic injury models.^{157,158} Similarly, complex IV-targeted antioxidants mitigate oxidative hepatocyte damage, promoting functional recovery.¹⁵⁹

This therapeutic approach demonstrates significant clinical promise through its multifaceted advantages. The foremost strength lies in exceptional molecular specificity achieved via direct targeting of mitochondrial complexes, effectively circumventing off-target effects on healthy cellular components. Complementing this precision, the inherent functional diversity among different complexes enables tailored therapeutic strategies aligned with individual pathophysiological profiles. Furthermore, comparative pharmacological analyses reveal superior safety margins over conventional treatments, particularly in minimizing systemic toxicity. However, translational implementation faces several critical challenges. Chief among these is interindividual heterogeneity in complex expression patterns, necessitating sophisticated pharmacogenomic approaches for dose optimization.^{160–162} Additionally, formulation challenges persist due to suboptimal pharmacokinetic properties, including reduced bioavailability and chemical instability. Of particular concern is emerging evidence of adaptive metabolic reprogramming in chronic disease states, which may undermine long-term therapeutic efficacy. Despite these challenges, targeted modulation of mitochondrial complexes remains a promising area of hepatology research, with the potential to redefine current treatment paradigms by integrating mechanistic precision and metabolic restoration.

Mitochondrial complexes are crucial for sustaining hepatocyte energy metabolism, oxidative balance, and overall cellular viability. Dysfunction of these complexes disrupts ATP synthesis, triggers apoptosis and autophagy dysregulation, and accelerates liver disease progression. Therefore, investi-

gating the mechanisms of mitochondrial complex dysfunction and their relationship with liver diseases will deepen our understanding of liver pathology and facilitate the development of targeted therapeutic approaches.

Conclusions

Emerging evidence underscores mitochondrial complexes and respiratory SCs as central regulators of hepatic bioenergetics, redox balance, and metabolic adaptability. Their structural-functional interdependence ensures efficient electron transport and ATP synthesis while maintaining resilience against oxidative and metabolic stress. Disruption of complex integrity, as observed in NAFLD, alcoholic liver disease, and other liver pathologies, compromises energy metabolism and amplifies oxidative damage, thereby accelerating disease progression. Therapeutically, strategies that stabilize SC architecture, enhance specific complex activity, or restore essential assembly factors hold promise for mitigating mitochondrial dysfunction. Integrating these approaches within precision medicine frameworks could enable targeted, patient-specific interventions, offering a pathway toward improved outcomes in diverse liver diseases.

Funding

This research was funded by the Research Project of Zhejiang Chinese Medical University (grant number 2024JKJNTZ01) and the Chinese Medicine Science Foundation of Zhejiang Chinese Medical University (grant number 2023J05).

Conflict of interest

The author has no conflicts of interest related to this publication.

Author contributions

HA is the sole author of the manuscript and approved the final version of the manuscript.

References

- [1] Devarbhani H, Asrani SK, Arab JP, Nartey YA, Pose E, Kamath PS. Global burden of liver disease: 2023 update. *J Hepatol* 2023;79(2):516–537. doi:10.1016/j.jhep.2023.03.017, PMID:36990226.
- [2] LeFort KR, Rungratanawanich W, Song BJ. Contributing roles of mitochondrial dysfunction and hepatocyte apoptosis in liver diseases through oxidative stress, post-translational modifications, inflammation, and intestinal barrier dysfunction. *Cell Mol Life Sci* 2024;81(1):34. doi:10.1007/s00018-023-05061-7, PMID:38214802.
- [3] Zhang X, Tachibana S, Wang H, Hisada M, Williams GM, Gao B, *et al.* Interleukin-6 is an important mediator for mitochondrial DNA repair after alcoholic liver injury in mice. *Hepatology* 2010;52(6):2137–2147. doi:10.1002/hep.23909, PMID:20931558.
- [4] Goicoechea L, Conde de la Rosa L, Torres S, García-Ruiz C, Fernández-Checa JC. Mitochondrial cholesterol: Metabolism and impact on redox biology and disease. *Redox Biol* 2023;61:102643. doi:10.1016/j.redox.2023.102643, PMID:36857930.
- [5] Wu G, Yang J, Lv H, Jing W, Zhou J, Feng Y, *et al.* Taurine prevents ethanol-induced apoptosis mediated by mitochondrial or death receptor pathways in liver cells. *Amino Acids* 2018;50(7):863–875. doi:10.1007/s00726-018-2561-3, PMID:29626300.
- [6] Zhao MG, Sheng XP, Huang YP, Wang YT, Jiang CH, Zhang J, *et al.* Triterpenic acids-enriched fraction from *Cyclocarya paliurus* attenuates non-alcoholic fatty liver disease via improving oxidative stress and mitochondrial dysfunction. *Biomed Pharmacother* 2018;104:229–239. doi:10.1016/j.biopha.2018.03.170, PMID:29775890.
- [7] Kim TS, Lee M, Park M, Kim SY, Shim MS, Lee CY, *et al.* Metformin and Dichloroacetate Suppress Proliferation of Liver Cancer Cells by Inhibiting mTOR Complex 1. *Int J Mol Sci* 2021;22(18):10027. doi:10.3390/ijms221810027, PMID:34576192.
- [8] Davuluri G, Allaway A, Thapaliya S, Rennison JH, Singh D, Kumar A, *et al.* Hyperammonaemia-induced skeletal muscle mitochondrial dysfunction results in cataplerosis and oxidative stress. *J Physiol* 2016;594(24):7341–

7360. doi:10.1113/JP272796, PMID:27558544.
- [9] Khoo NKH, Fazzari M, Chartoumpekis DV, Li L, Guimaraes DA, Arteel GE, *et al.* Electrophilic nitro-oleic acid reverses obesity-induced hepatic steatosis. *Redox Biol* 2019;22:101132. doi:10.1016/j.redox.2019.101132, PMID:30769284.
- [10] Li J, Liang N, Long X, Zhao J, Yang J, Du X, *et al.* SDHC-related deficiency of SDH complex activity promotes growth and metastasis of hepatocellular carcinoma via ROS/NFκB signaling. *Cancer Lett* 2019;461:44–55. doi:10.1016/j.canlet.2019.07.001, PMID:31278950.
- [11] Luo H, Wang Q, Yang F, Liu R, Gao Q, Cheng B, *et al.* Signaling metabolite succinylacetone activates HIF-1α and promotes angiogenesis in GSTZ1-deficient hepatocellular carcinoma. *JCI Insight* 2023;8(23):e164968. doi:10.1172/jci.insight.164968, PMID:37906252.
- [12] Rizza S, Montagna C, Cardaci S, Maiani E, Di Giacomo G, Sanchez-Quiles V, *et al.* S-nitrosylation of the Mitochondrial Chaperone TRAP1 Sensitizes Hepatocellular Carcinoma Cells to Inhibitors of Succinate Dehydrogenase. *Cancer Res* 2016;76(14):4170–4182. doi:10.1158/0008-5472.Can-15-2637, PMID:27216192.
- [13] Xu Y, Yang Y, Yu X, Han L, Liu S, Zhang N, *et al.* Integrated functional genomics-identified LYRM4 promotes fumarate accumulation and hepatocellular carcinoma progression. *Arch Biochem Biophys* 2025;770:110448. doi:10.1016/j.abb.2025.110448, PMID:40320061.
- [14] Staňková P, Kučera O, Peterová E, Lotková H, Maseko TE, Nožičková K, *et al.* Adaptation of Mitochondrial Substrate Flux in a Mouse Model of Nonalcoholic Fatty Liver Disease. *Int J Mol Sci* 2020;21(3):1101. doi:10.3390/ijms21031101, PMID:32046101.
- [15] Abe J, Vujic A, Prag HA, Murphy MP, Krieg T. Malonate given at reperfusion prevents post-myocardial infarction heart failure by decreasing ischemia/reperfusion injury. *Basic Res Cardiol* 2024;119(4):691–697. doi:10.1007/s00395-024-01063-z, PMID:38864895.
- [16] Weng Z, Zhou P, Salminen WF, Yang X, Harrill AH, Cao Z, *et al.* Green tea epigallocatechin gallate binds to and inhibits respiratory complexes in swelling but not normal rat hepatic mitochondria. *Biochem Biophys Res Commun* 2014;443(3):1097–1104. doi:10.1016/j.bbrc.2013.12.110, PMID:24384371.
- [17] Li YH, Choi DH, Lee EH, Seo SR, Lee S, Cho EH. Sirtuin 3 (SIRT3) Regulates α-Smooth Muscle Actin (α-SMA) Production through the Succinate Dehydrogenase-G Protein-coupled Receptor 91 (GPR91) Pathway in Hepatic Stellate Cells. *J Biol Chem* 2016;291(19):10277–10292. doi:10.1074/jbc.M115.692244, PMID:26912655.
- [18] Chi X, Yu D, Li P, Lu Q, Jiang W, Hao K. The protection effects of (1E,6E)-1,7-diphenylhepta-1,6-diene-3,5-dione, a curcumin analogue, against operative liver injury in rats. *Eur J Pharm Sci* 2017;100:94–101. doi:10.1016/j.ejps.2016.12.042, PMID:28057548.
- [19] Yang H, Xuefeng Y, Shandong W, Jianhua X. COX-2 in liver fibrosis. *Clin Chim Acta* 2020;506:196–203. doi:10.1016/j.cca.2020.03.024, PMID:32184095.
- [20] Li J, Hu R, Liu X, Peng L, Yi J, Zhong X, *et al.* S100A9/RAGE pathway regulation of mitophagy and the effect of JianPi LiShi YangGan formula in acute-on-chronic liver failure. *J Ethnopharmacol* 2025;348:119887. doi:10.1016/j.jep.2025.119887, PMID:40294664.
- [21] Fuentes-Agudo M, Luque-Tévar M, Cucarella C, Brea R, Boscá L, Quintana-Cabrera R, *et al.* COX-2 Expression in Hepatocytes Improves Mitochondrial Function after Hepatic Ischemia-Reperfusion Injury. *Antioxidants (Basel)* 2022;11(9):1724. doi:10.3390/antiox11091724, PMID:36139798.
- [22] Pham L, Arroum T, Wan J, Pavelich L, Bell J, Morse PT, *et al.* Regulation of mitochondrial oxidative phosphorylation through tight control of cytochrome c oxidase in health and disease - Implications for ischemia/reperfusion injury, inflammatory diseases, diabetes, and cancer. *Redox Biol* 2024;78:103426. doi:10.1016/j.redox.2024.103426, PMID:39566165.
- [23] Morse PT, Goebel DJ, Wan J, Tuck S, Hakim L, Hüttemann CL, *et al.* Cytochrome c oxidase-modulatory near-infrared light penetration into the human brain: Implications for the noninvasive treatment of ischemia/reperfusion injury. *IUBMB Life* 2021;73(3):554–567. doi:10.1002/iub.2405, PMID:33166061.
- [24] Zhao Z, Zhang C, Lin J, Zheng L, Li H, Qi X, *et al.* COX-2/sEH Dual Inhibitor PTUPB Alleviates CCI (4) -Induced Liver Fibrosis and Portal Hypertension. *Front Med (Lausanne)* 2021;8:761517. doi:10.3389/fmed.2021.761517, PMID:35004731.
- [25] Shiriaeva AP, Baidiuk EV, Arkad'eva AV, Okovityi SV, Morozov VI, Sakuta GA. [Hepatocyte mitochondrion respiratory chain in rats with experimental toxic hepatitis]. *Tsitologiya* 2007;49(2):125–132. PMID:17432597.
- [26] Bhopale KK, Amer SM, Kaphalia L, Soman KV, Wiktorowicz JE, Shakeel Ansari GA, *et al.* Proteomic Profiling of Liver and Plasma in Chronic Ethanol Feeding Model of Hepatic Alcohol Dehydrogenase-Deficient Mice. *Alcohol Clin Exp Res* 2017;41(10):1675–1685. doi:10.1111/acer.13470, PMID:28792616.
- [27] Rao Y, Lu YT, Li C, Song QQ, Xu YH, Xu Z, *et al.* Bouchardatine analogue alleviates non-alcoholic fatty liver disease/non-alcoholic steatohepatitis in high-fat fed mice by inhibiting ATP synthase activity. *Br J Pharmacol* 2019;176(16):2877–2893. doi:10.1111/bph.14713, PMID:31113010.
- [28] Maclean AE, Bridges HR, Silva MF, Ding S, Ovcariakova J, Hirst J, *et al.* Complexome profile of Toxoplasma gondii mitochondria identifies divergent subunits of respiratory chain complexes including new subunits of cytochrome bc1 complex. *PLoS Pathog* 2021;17(3):e1009301. doi:10.1371/journal.ppat.1009301, PMID:33651838.
- [29] Sazanov LA. Respiratory complex I: mechanistic and structural insights provided by the crystal structure of the hydrophilic domain. *Biochemistry* 2007;46(9):2275–2288. doi:10.1021/bi602508x, PMID:17274631.
- [30] Hirst J. Mitochondrial complex I. *Annu Rev Biochem* 2013;82:551–575. doi:10.1146/annurev-biochem-070511-103700, PMID:23527692.
- [31] Xiong W, Yuan Z, Wang T, Wu S, Xiong Y, Yao Y, *et al.* Quercitrin Attenuates Acetaminophen-Induced Acute Liver Injury by Maintaining Mitochondrial Complex I Activity. *Front Pharmacol* 2021;12:586010. doi:10.3389/fphar.2021.586010, PMID:34025394.
- [32] Matyas C, Varga ZV, Mukhopadhyay P, Palocz J, Lajtos T, Erdelyi K, *et al.* Chronic plus binge ethanol feeding induces myocardial oxidative stress, mitochondrial and cardiovascular dysfunction, and steatosis. *Am J Physiol Heart Circ Physiol* 2016;310(11):H1658–H1670. doi:10.1152/ajp-heart.00214.2016, PMID:27106042.
- [33] Cho T, Wang X, Uetrecht J. Rotenone Increases Isoniazid Toxicity but Does Not Cause Significant Liver Injury: Implications for the Hypothesis that Inhibition of the Mitochondrial Electron Transport Chain Is a Common Mechanism of Idiosyncratic Drug-Induced Liver Injury. *Chem Res Toxicol* 2019;32(7):1423–1431. doi:10.1021/acs.chemrestox.9b00116, PMID:31251588.
- [34] Ding N, Wang K, Jiang H, Yang M, Zhang L, Fan X, *et al.* AGK regulates the progression to NASH by affecting mitochondria complex I function. *Theranostics* 2022;12(7):3237–3250. doi:10.7150/thno.69826, PMID:35547757.
- [35] Paradies G, Paradies V, Ruggiero FM, Petrosillo G. Oxidative stress, cardiolipin and mitochondrial dysfunction in nonalcoholic fatty liver disease. *World J Gastroenterol* 2014;20(39):14205–14218. doi:10.3748/wjg.v20.i39.14205, PMID:25339807.
- [36] Chávez E, Lozano-Rosas MG, Domínguez-López M, Velasco-Loyden G, Rodríguez-Aguilera JR, José-Núñez C, *et al.* Functional, Metabolic, and Dynamic Mitochondrial Changes in the Rat Cirrhotic-Hepatocellular Carcinoma Model and the Protective Effect of IFC-305. *J Pharmacol Exp Ther* 2017;361(2):292–302. doi:10.1124/jpet.116.239301, PMID:28209723.
- [37] Xiao MH, Lin YF, Xie PP, Chen HX, Deng JW, Zhang W, *et al.* Downregulation of a mitochondrial micropeptide, MPM, promotes hepatoma metastasis by enhancing mitochondrial complex I activity. *Mol Ther* 2022;30(2):714–725. doi:10.1016/j.ymthe.2021.08.032, PMID:34478872.
- [38] Imaizumi N, Kwang Lee K, Zhang C, Boelsterli UA. Mechanisms of cell death pathway activation following drug-induced inhibition of mitochondrial complex I. *Redox Biol* 2015;4:279–288. doi:10.1016/j.redox.2015.01.005, PMID:25625582.
- [39] Lehninger AL, Nelson DL, Cox MM. . Lehninger principles of biochemistry. 3rd ed. New York: Worth Publishers; 2000.
- [40] Ragab EM, El Gamal DM, Mohamed TM, Khamis AA. Study of the inhibitory effects of chrysin and its nanoparticles on mitochondrial complex II subunit activities in normal mouse liver and human fibroblasts. *J Genet Eng Biotechnol* 2022;20(1):15. doi:10.1186/s43141-021-00286-0, PMID:35089446.
- [41] Sun F, Huo X, Zhai Y, Wang A, Xu J, Su D, *et al.* Crystal structure of mitochondrial respiratory membrane protein complex II. *Cell* 2005;121(7):1043–1057. doi:10.1016/j.cell.2005.05.025, PMID:15989954.
- [42] Gnaiger E. Correction: Complex II ambiguities-FADH2 in the electron transfer system. *J Biol Chem* 2024;300(12):108009. doi:10.1016/j.jbc.2024.108009, PMID:39662301.
- [43] Chouchani ET, Pell VR, Gaude E, Aksentijevic D, Sundier SY, Robb EL, *et al.* Ischaemic accumulation of succinate controls reperfusion injury through mitochondrial ROS. *Nature* 2014;515(7527):431–435. doi:10.1038/nature13909, PMID:25383517.
- [44] Bouillaud F. Inhibition of Succinate Dehydrogenase by Pesticides (SDHIs) and Energy Metabolism. *Int J Mol Sci* 2023;24(4):4045. doi:10.3390/ijms24044045, PMID:36835457.
- [45] Dalla Pozza E, Dando I, Pacchiana R, Liboi E, Scupoli MT, Donadelli M, *et al.* Regulation of succinate dehydrogenase and role of succinate in cancer. *Semin Cell Dev Biol* 2020;98:4–14. doi:10.1016/j.semcdb.2019.04.013, PMID:31039394.
- [46] Shimizu T, Inoue K, Hachiya H, Shibuya N, Shimoda M, Kubota K. Frequent alteration of the protein synthesis of enzymes for glucose metabolism in hepatocellular carcinomas. *J Gastroenterol* 2014;49(9):1324–1332. doi:10.1007/s00535-013-0895-x, PMID:24203292.
- [47] Tseng PL, Wu WH, Hu TH, Chen CH, Cheng HC, Li CF, *et al.* Decreased succinate dehydrogenase B in human hepatocellular carcinoma accelerates tumor malignancy by inducing the Warburg effect. *Sci Rep* 2018;8(1):3081. doi:10.1038/s41598-018-21361-6, PMID:29449614.
- [48] Kroemer G. Mitochondria in cancer. *Oncogene* 2006;25(34):4630–4632. doi:10.1038/sj.onc.1209589, PMID:16892077.
- [49] Banerjee R, Purhonen J, Kallijärvi J. The mitochondrial coenzyme Q junction and complex III: biochemistry and pathophysiology. *FEBS J* 2022;289(22):6936–6958. doi:10.1111/febs.16164, PMID:34428349.
- [50] Lenaz G, Fato R, Formigini G, Genova ML. The role of Coenzyme Q in mitochondrial electron transport. *Mitochondrion* 2007;7(Suppl):S8–33. doi:10.1016/j.mito.2007.03.009, PMID:17485246.
- [51] Guarás A, Perales-Clemente E, Calvo E, Acín-Pérez R, Loureiro-Lopez M, Pujol C, *et al.* The CoQH2/CoQ Ratio Serves as a Sensor of Respiratory Chain Efficiency. *Cell Rep* 2016;15(1):197–209. doi:10.1016/j.celrep.2016.03.009, PMID:27052170.
- [52] Lin J, Puigserver P, Donovan J, Tarr P, Spiegelman BM. Peroxisome proliferator-activated receptor gamma coactivator 1βeta (PGC-1βeta), a novel PGC-1-related transcription coactivator associated with host cell factor. *J Biol Chem* 2002;277(3):1645–1648. doi:10.1074/jbc.C100631200, PMID:11733490.
- [53] Belosludtseva NV, Starinets VS, Semenova AA, Igoshkina AD, Dubinin MV, Belosludtsev KN. S-15176 Difumarate Salt Can Impair Mitochondrial Function through Inhibition of the Respiratory Complex III and Permeabilization of the Inner Mitochondrial Membrane. *Biology (Basel)* 2022;11(3):380. doi:10.3390/biology11030380, PMID:35336754.
- [54] Katsyuba E, Romani M, Hofer D, Auwerx J. NAD(+) homeostasis in health

- and disease. *Nat Metab* 2020;2(1):9–31. doi:10.1038/s42255-019-0161-5, PMID:32694684.
- [55] Lopez-Cruzan M, Herman B. Loss of caspase-2 accelerates age-dependent alterations in mitochondrial production of reactive oxygen species. *BioGerontology* 2013;14(2):121–130. doi:10.1007/s10522-013-9415-x, PMID:23504374.
- [56] Yang Q, Wang L, Liu J, Cao W, Pan Q, Li M. Targeting the complex I and III of mitochondrial electron transport chain as a potentially viable option in liver cancer management. *Cell Death Discov* 2021;7(1):293. doi:10.1038/s41420-021-00675-x, PMID:34650055.
- [57] Kadenbach B. Complex IV - The regulatory center of mitochondrial oxidative phosphorylation. *Mitochondrion* 2021;58:296–302. doi:10.1016/j.mito.2020.10.004, PMID:33069909.
- [58] Srinivasan S, Avadhani NG. Cytochrome c oxidase dysfunction in oxidative stress. *Free Radic Biol Med* 2012;53(6):1252–1263. doi:10.1016/j.freeradbiomed.2012.07.021, PMID:22841758.
- [59] Kumar A, Davuluri G, Welch N, Kim A, Gangadhariah M, Allaway A, *et al.* Oxidative stress mediates ethanol-induced skeletal muscle mitochondrial dysfunction and dysregulates protein synthesis and autophagy. *Free Radic Biol Med* 2019;145:284–299. doi:10.1016/j.freeradbiomed.2019.09.031, PMID:31574345.
- [60] Sarti P, Aresse M, Bacchi A, Barone MC, Forte E, Mastronicola D, *et al.* Nitric oxide and mitochondrial complex IV. *IUBMB Life* 2003;55(10–11):605–611. doi:10.1080/15216540310001628726, PMID:14711006.
- [61] Müller-Höcker J, Aust D, Rohrbach H, Napiwotzky J, Reith A, Link TA, *et al.* Defects of the respiratory chain in the normal human liver and in cirrhosis during aging. *Hepatology* 1997;26(3):709–719. doi:10.1002/hep.510260324, PMID:9303502.
- [62] Zhang J, Kay MK, Park MH, Meruvu S, Powell C, Choudhury M. LncRNA DLEU2 regulates sirtuins and mitochondrial respiratory chain complex IV: a novel pathway in obesity and offspring's health. *Int J Obes (Lond)* 2022;46(5):969–976. doi:10.1038/s41366-022-01075-6, PMID:35058572.
- [63] Sang Y, Liu JY, Wang FY, Luo XY, Chen ZQ, Zhuang SM, *et al.* Mitochondrial micropeptide STMP1 promotes G1/S transition by enhancing mitochondrial complex IV activity. *Mol Ther* 2022;30(8):2844–2855. doi:10.1016/j.ymthe.2022.04.012, PMID:35450818.
- [64] Zazueta C, Buelna-Chontal M, Macías-López A, Román-Anguiano NG, González-Pacheco H, Pavón N, *et al.* Cytidine-5'-Diphosphocholine Protects the Liver From Ischemia/Reperfusion Injury Preserving Mitochondrial Function and Reducing Oxidative Stress. *Liver Transpl* 2018;24(8):1070–1083. doi:10.1002/lt.25179, PMID:29679463.
- [65] Zhang P, Lu H, Wu Y, Lu D, Li C, Yang X, *et al.* COX5A Alleviates Doxorubicin-Induced Cardiotoxicity by Suppressing Oxidative Stress, Mitochondrial Dysfunction and Cardiomyocyte Apoptosis. *Int J Mol Sci* 2023;24(12):10400. doi:10.3390/ijms241210400, PMID:37373547.
- [66] Fernandez-Vizcarra E, Zeviani M. Mitochondrial disorders of the OXPHOS system. *FEBS Lett* 2021;595(8):1062–1106. doi:10.1002/1873-3468.13995, PMID:33159691.
- [67] Hu X, Que W, Hirano H, Wang Z, Nozawa N, Ishii T, *et al.* 5-Aminolevulinic acid/sodium ferrous citrate enhanced the antitumor effects of programmed cell death-ligand 1 blockade by regulation of exhausted T cell metabolism in a melanoma model. *Cancer Sci* 2021;112(7):2652–2663. doi:10.1111/cas.14930, PMID:33934440.
- [68] Zhao RZ, Jiang S, Zhang L, Yu ZB. Mitochondrial electron transport chain, ROS generation and uncoupling (Review). *Int J Mol Med* 2019;44(1):3–15. doi:10.3892/ijmm.2019.4188, PMID:31115493.
- [69] Ren Z, Chen S, Seo JE, Guo X, Li D, Ning B, *et al.* Mitochondrial dysfunction and apoptosis underlie the hepatotoxicity of perhexiline. *Toxicol In Vitro* 2020;69:104987. doi:10.1016/j.tiv.2020.104987, PMID:32861758.
- [70] Lee K, Haddad A, Osme A, Kim C, Borzou A, Ilchenko S, *et al.* Hepatic Mitochondrial Defects in a Nonalcoholic Fatty Liver Disease Mouse Model Are Associated with Increased Degradation of Oxidative Phosphorylation Subunits. *Mol Cell Proteomics* 2018;17(12):2371–2386. doi:10.1074/mcp.RA118.000961, PMID:30171159.
- [71] Eldeeb MH, Camacho Lopez LJ, Fontanesi F. Mitochondrial respiratory supercomplexes of the yeast *Saccharomyces cerevisiae*. *IUBMB Life* 2024;76(8):485–504. doi:10.1002/iub.2817, PMID:38529880.
- [72] Javadov S, Jiang S, Chapa-Dubocq XR, Khuchua Z, Camara AK. Mitochondrial respiratory supercomplexes in mammalian cells: structural versus functional role. *J Mol Med (Berl)* 2021;99(1):57–73. doi:10.1007/s00109-020-02004-8, PMID:33201259.
- [73] Zheng W, Chai P, Zhu J, Zhang K. High-resolution in situ structures of mammalian respiratory supercomplexes. *Nature* 2024;631(8019):232–239. doi:10.1038/s41586-024-07488-9, PMID:38811722.
- [74] Zheng W, Chai P, Zhu J, Zhang K. High-resolution In-situ Structures of Mammalian Mitochondrial Respiratory Supercomplexes in Reaction within Native Mitochondria. *bioRxiv* 2024. doi:10.1101/2024.04.02.587796, PMID:38617346.
- [75] Ma X, McKeen T, Zhang J, Ding WX. Role and Mechanisms of Mitophagy in Liver Diseases. *Cells* 2020;9(4):837. doi:10.3390/cells9040837, PMID:32244304.
- [76] Sullivan EM, Pennington ER, Green WD, Beck MA, Brown DA, Shaikh SR. Mechanisms by Which Dietary Fatty Acids Regulate Mitochondrial Structure-Function in Health and Disease. *Adv Nutr* 2018;9(3):247–262. doi:10.1093/advances/nmy007, PMID:29767698.
- [77] Zhang C, Zhao Y, Yu M, Qin J, Ye B, Wang Q. Mitochondrial Dysfunction and Chronic Liver Disease. *Curr Issues Mol Biol* 2022;44(7):3156–3165. doi:10.3390/cimb44070218, PMID:35877442.
- [78] Middleton P, Vergis N. Mitochondrial dysfunction and liver disease: role, relevance, and potential for therapeutic modulation. *Therap Adv Gastroen-*
- terol* 2021;14:17562848211031394. doi:10.1177/17562848211031394, PMID:34377148.
- [79] Prola A, Pilot-Storck F. Cardiolipin Alterations during Obesity: Exploring Therapeutic Opportunities. *Biology (Basel)* 2022;11(11):1638. doi:10.3390/biology11111638, PMID:36358339.
- [80] Sullivan EM, Fix A, Crouch MJ, Sparagna GC, Zeczycki TN, Brown DA, *et al.* Murine diet-induced obesity remodels cardiac and liver mitochondrial phospholipid acyl chains with differential effects on respiratory enzyme activity. *J Nutr Biochem* 2017;45:94–103. doi:10.1016/j.jnutbio.2017.04.004, PMID:28437736.
- [81] Moreno-Lastres D, Fontanesi F, García-Consuegra I, Martín MA, Arenas J, Barrientos A, *et al.* Mitochondrial complex I plays an essential role in human respirasome assembly. *Cell Metab* 2012;15(3):324–335. doi:10.1016/j.cmet.2012.01.015, PMID:22342700.
- [82] Lim SC, Hroudová J, Van Bergen NJ, Lopez Sanchez MI, Trousseau IA, McKenzie M. Loss of mitochondrial DNA-encoded protein ND1 results in disruption of complex I biogenesis during early stages of assembly. *FASEB J* 2016;30(6):2236–2248. doi:10.1096/fj.201500137R, PMID:26929434.
- [83] Jeon TJ, Lee SG, Yoo SH, Kim M, Song D, Ryu J, *et al.* A Dynamic Substrate Pool Revealed by cryo-EM of a Lipid-Preserved Respiratory Supercomplex. *Antioxid Redox Signal* 2022. doi:10.1089/ars.2021.0114, PMID:34913730.
- [84] Gu J, Yao M, Yao D, Wang L, Yang X, Yao D. Nonalcoholic Lipid Accumulation and Hepatocyte Malignant Transformation. *J Clin Transl Hepatol* 2016;4(2):123–130. doi:10.14218/JCTH.2016.00010, PMID:27350942.
- [85] Sun Q, Zhong W, Zhang W, Zhou Z. Defect of mitochondrial respiratory chain is a mechanism of ROS overproduction in a rat model of alcoholic liver disease: role of zinc deficiency. *Am J Physiol Gastrointest Liver Physiol* 2016;310(3):G205–G214. doi:10.1152/ajpgi.00270.2015, PMID:26585415.
- [86] Sakurai T, Chen Z, Yamahata A, Hayasaka T, Satoh H, Sekiguchi H, *et al.* A mouse model of short-term, diet-induced fatty liver with abnormal cardiolipin remodeling via downregulated Tafazzin gene expression. *J Sci Food Agric* 2021;101(12):4995–5001. doi:10.1002/jsfa.11144, PMID:33543498.
- [87] Dukić M, Radonjić T, Jovanović I, Zdravković M, Todorović Z, Krašnik N, *et al.* Alcohol, Inflammation, and Microbiota in Alcoholic Liver Disease. *Int J Mol Sci* 2023;24(4):3735. doi:10.3390/ijms24043735, PMID:36835145.
- [88] García-Ruiz I, Solís-Muñoz P, Fernández-Moreira D, Grau M, Colina F, Muñoz-Yague T, *et al.* High-fat diet decreases activity of the oxidative phosphorylation complexes and causes nonalcoholic steatohepatitis in mice. *Dis Model Mech* 2014;7(11):1287–1296. doi:10.1242/dmm.016766, PMID:25261569.
- [89] Shang L, Ren H, Wang S, Liu H, Hu A, Gou P, *et al.* SS-31 Protects Liver from Ischemia-Reperfusion Injury via Modulating Macrophage Polarization. *Oxid Med Cell Longev* 2021;2021:6662156. doi:10.1155/2021/6662156, PMID:33869618.
- [90] Yoon JR, Lee EJ, Kim HD, Lee JH, Kang HC. Polyunsaturated fatty acid-enriched diet therapy for a child with epilepsy. *Brain Dev* 2014;36(2):163–166. doi:10.1016/j.braindev.2013.01.017, PMID:23465587.
- [91] de Mello AH, Schraiber RB, Goldim MPS, Garcez ML, Gomes ML, de Bem Silveira G, *et al.* Omega-3 Fatty Acids Attenuate Brain Alterations in High-Fat Diet-Induced Obesity Model. *Mol Neurobiol* 2019;56(1):513–524. doi:10.1007/s12035-018-1097-6, PMID:29728888.
- [92] Real LM, Caruz A, Rivero-Juarez A, Soriano V, Neukam K, Rivero A, *et al.* A polymorphism linked to RRAS, SCAF1, IRF3 and BCL2L12 genes is associated with cirrhosis in hepatitis C virus carriers. *Liver Int* 2014;34(4):558–566. doi:10.1111/liv.12330, PMID:24131527.
- [93] Daly S, Bulovaite E, Handa A, Morris K, Muresan L, Adams C, *et al.* 3D Super-Resolution Imaging of PSD95 Reveals an Abundance of Diffuse Protein Supercomplexes in the Mouse Brain. *ACS Chem Neurosci* 2025;16(1):40–51. doi:10.1021/acschemneuro.4c00684, PMID:39702971.
- [94] Moreno-Justicia R, Gonzalez-Franquesa A, Stocks B, Deshmukh AS. Protocol to characterize mitochondrial supercomplexes from mouse tissues by combining BN-PAGE and MS-based proteomics. *STAR Protoc* 2022;3(1):101135. doi:10.1016/j.xpro.2022.101135, PMID:35128478.
- [95] Barros MH, McStay GP. Modular biogenesis of mitochondrial respiratory complexes. *Mitochondrion* 2020;50:94–114. doi:10.1016/j.mito.2019.10.008, PMID:31669617.
- [96] Yang F, Cao H, Su R, Guo J, Li C, Pan J, *et al.* Liver mitochondrial dysfunction and electron transport chain defect induced by high dietary copper in broilers. *Poult Sci* 2017;96(9):3298–3304. doi:10.3382/ps/pex137, PMID:28575470.
- [97] Lin S, Zhang H, Wang C, Su XL, Song Y, Wu P, *et al.* Metabolomics Reveal Nanoplastic-Induced Mitochondrial Damage in Human Liver and Lung Cells. *Environ Sci Technol* 2022;56(17):12483–12493. doi:10.1021/acs.est.2c03980, PMID:36005547.
- [98] Wang HW, Liu J, Wei SS, Zhao WP, Zhu SQ, Zhou BH. Mitochondrial respiratory chain damage and mitochondrial fusion disorder are involved in liver dysfunction of fluoride-induced mice. *Chemosphere* 2020;241:125099. doi:10.1016/j.chemosphere.2019.125099, PMID:31629238.
- [99] Abate M, Festa A, Falco M, Lombardi A, Luce A, Grimaldi A, *et al.* Mitochondria as playmakers of apoptosis, autophagy and senescence. *Semin Cell Dev Biol* 2020;98:139–153. doi:10.1016/j.semcdb.2019.05.022, PMID:31154010.
- [100] Green DR, Galluzzi L, Kroemer G. Mitochondria and the autophagy-inflammation-cell death axis in organismal aging. *Science* 2011;333(6046):1109–1112. doi:10.1126/science.1201940, PMID:21868666.
- [101] Kujoth GC, Hiona A, Pugh TD, Someya S, Panzer K, Wohlgenuth SE, *et al.* Mitochondrial DNA mutations, oxidative stress, and apoptosis in mammalian aging. *Science* 2005;309(5733):481–484. doi:10.1126/science.1112125, PMID:16020738.

- [102] Pirola CJ, Garaycoechea M, Flichman D, Castañó GO, Sookoian S. Liver mitochondrial DNA damage and genetic variability of Cytochrome b - a key component of the respirasome - drive the severity of fatty liver disease. *J Intern Med* 2021;289(1):84-96. doi:10.1111/joim.13147, PMID:32634278.
- [103] Valente WJ, Ericson NG, Long AS, White PA, Marchetti F, Bielas JH. Mitochondrial DNA exhibits resistance to induced point and deletion mutations. *Nucleic Acids Res* 2016;44(18):8513-8524. doi:10.1093/nar/gkw716, PMID:27550180.
- [104] Chiang Morales MD, Chang CY, Le VL, Huang IT, Tsai IL, Shih HJ, et al. High-Fructose/High-Fat Diet Downregulates the Hepatic Mitochondrial Oxidative Phosphorylation Pathway in Mice Compared with High-Fat Diet Alone. *Cells* 2022;11(21):3425. doi:10.3390/cells11213425, PMID:36359820.
- [105] Liu C, Sekine S, Ito K. Assessment of mitochondrial dysfunction-related, drug-induced hepatotoxicity in primary rat hepatocytes. *Toxicol Appl Pharmacol* 2016;302:23-30. doi:10.1016/j.taap.2016.04.010, PMID:27095095.
- [106] Musatov A, Robinson NC. Susceptibility of mitochondrial electron-transport complexes to oxidative damage. Focus on cytochrome c oxidase. *Free Radic Res* 2012;46(11):1313-1326. doi:10.3109/10715762.2012.717273, PMID:22856385.
- [107] Adisa RA, Sulaimon LA, Okeke EG, Ariyo OC, Abdulkareem FB. Mitoquinol mesylate (MITOQ) attenuates diethyl nitrosamine-induced hepatocellular carcinoma through modulation of mitochondrial antioxidant defense systems. *Toxicol Res* 2022;38(3):275-291. doi:10.1007/s43188-021-00105-1, PMID:35874502.
- [108] Duduk NI, Kravchuk RI, Zimatkin SM. [Morpho-functional changes in the liver and the possibility of their correction in the offspring of rats with cholestasis]. *Morfologiya* 2015;147(1):48-53. PMID:25958728.
- [109] Ezgu F, Senaca S, Gunduz M, Tumer L, Hasanoglu A, Tiras U, et al. Severe renal tubulopathy in a newborn due to BCS1L gene mutation: effects of different treatment modalities on the clinical course. *Gene* 2013;528(2):364-366. doi:10.1016/j.gene.2013.07.007, PMID:23892085.
- [110] Xu X, Feng J, Wang X, Zeng X, Luo Y, He X, et al. Mitochondrial GRIM19 Loss Induces Liver Fibrosis through NLRP3/IL33 Activation via Reactive Oxygen Species/NF- κ B Signaling. *J Clin Transl Hepatol* 2024;12(6):539-550. doi:10.14218/JCTH.2023.00562, PMID:38974954.
- [111] Briet F, Aqel BA, Jeejeebhoy KN, Scapio JS. A low pretransplant peripheral blood mononuclear cell complex I activity predicts metabolic disturbances and inability to regain fat free mass in cirrhotic patients undergoing liver transplantation. *Nutr Res* 2009;29(1):26-34. doi:10.1016/j.nutres.2008.10.004, PMID:19185774.
- [112] Zhou M, Yu Y, Luo Y, Luo X, Zhang Y, Zhou X, et al. Mitochondrial Transplantation: A Unique Treatment Strategy. *J Cardiovasc Pharmacol* 2022;79(6):759-768. doi:10.1097/FJC.0000000000001247, PMID:35639247.
- [113] Li X, Ruan T, Wang S, Sun X, Liu C, Peng Y, et al. Mitochondria at the Crossroads of Cholestatic Liver Injury: Targeting Novel Therapeutic Avenues. *J Clin Transl Hepatol* 2024;12(9):792-801. doi:10.14218/JCTH.2024.00087, PMID:39280065.
- [114] Li B, Li B, Qiao X, Meng W, Xie Y, Gong J, et al. Targeting mitochondrial transfer as a promising therapeutic strategy. *Trends Mol Med* 2025. doi:10.1016/j.molmed.2025.04.002, PMID:40335384.
- [115] Zhao Z, Hou Y, Zhou W, Keerthiga R, Fu A. Mitochondrial transplantation therapy inhibit carbon tetrachloride-induced liver injury through scavenging free radicals and protecting hepatocytes. *Bioeng Transl Med* 2021;6(2):e10209. doi:10.1002/btm2.10209, PMID:34027095.
- [116] Chen Y, Yang F, Chu Y, Yun Z, Yan Y, Jin J. Mitochondrial transplantation: opportunities and challenges in the treatment of obesity, diabetes, and nonalcoholic fatty liver disease. *J Transl Med* 2022;20(1):483. doi:10.1186/s12967-022-03693-0, PMID:36273156.
- [117] Shi X, Bai H, Zhao M, Li X, Sun X, Jiang H, et al. Treatment of acetaminophen-induced liver injury with exogenous mitochondria in mice. *Transl Res* 2018;196:31-41. doi:10.1016/j.trsl.2018.02.003, PMID:29548626.
- [118] Li J, Lv M, Yuan Z, Ge J, Geng T, Gong D, et al. PGC-1 α Promotes mitochondrial biosynthesis and energy metabolism of goose fatty liver. *Poult Sci* 2025;104(1):104617. doi:10.1016/j.psj.2024.104617, PMID:39644719.
- [119] Liu X, Khouri-Farah N, Wu CH, Wu GY. Targeted delivery of mitochondria to the liver in rats. *J Gastroenterol Hepatol* 2020;35(12):2241-2247. doi:10.1111/jgh.15091, PMID:32386240.
- [120] Mokhtari B, Alihemmati A, Bafadam S, Boraghi S, Badalzadeh R, Mahmoodpoor A. Mitotherapy attenuates sepsis-induced brain injury in rats subjected to cecal ligation and puncture: Role of SIRT-1/PGC-1 α network. *Iran J Basic Med Sci* 2025;28(8):1065-1074. doi:10.22038/ijbms.2025.84848.18363, PMID:40584438.
- [121] Azevedo RDS, Falcão KVG, Almeida SMV, Araújo MC, Silva-Filho RC, Souza Maia MB, et al. The tissue-specific nature of physiological zebrafish mitochondrial bioenergetics. *Mitochondrion* 2024;77:101901. doi:10.1016/j.mito.2024.101901, PMID:38777222.
- [122] Hu WJ, Wei H, Cai LL, Xu YH, Du R, Zhou Q, et al. Magnetic targeting enhances the neuroprotective function of human mesenchymal stem cell-derived iron oxide exosomes by delivering miR-1228-5p. *J Nanobiotechnology* 2024;22(1):665. doi:10.1186/s12951-024-02941-3, PMID:39468528.
- [123] Yang C, Liao X, Wang Y, Zhong W, Xian F, Gao M, et al. Biomimetic Nanoparticle Integrating CeO $_2$ Nanoparticles and Anti-Inflammatory Peptides for Osteoarthritis Therapy. *Small* 2025;21(32):e2502677. doi:10.1002/smll.202502677, PMID:40391633.
- [124] Hayashida K, Takegawa R, Endo Y, Yin T, Choudhary RC, Aoki T, et al. Exogenous mitochondrial transplantation improves survival and neurological outcomes after resuscitation from cardiac arrest. *BMC Med* 2023;21(1):56. doi:10.1186/s12916-023-02759-0, PMID:36922820.
- [125] Khalfi P, Suspène R, Raymond KA, Caval V, Caignard G, Berry N, et al. Antagonism of ALAS1 by the Measles Virus V protein contributes to degradation of the mitochondrial network and promotes interferon response. *PLoS Pathog* 2023;19(2):e1011170. doi:10.1371/journal.ppat.1011170, PMID:36802406.
- [126] Cano-Sanchez M, Ben-Hassen K, Louis OP, Dantin F, Gueye P, Roques F, et al. Bothrops lanceolatus snake venom impairs mitochondrial respiration and induces DNA release in human heart preparation. *PLoS Negl Trop Dis* 2022;16(6):e0010523. doi:10.1371/journal.pntd.0010523, PMID:35727836.
- [127] Cherry AD. Mitochondrial transplantation in cardiac surgical patients: optimism, caveats, and outstanding questions. *Curr Opin Anaesthesiol* 2023;36(1):5-10. doi:10.1097/ACO.0000000000001202, PMID:36550601.
- [128] Sun X, Gao R, Li W, Zhao Y, Yang H, Chen H, et al. Alda-1 treatment promotes the therapeutic effect of mitochondrial transplantation for myocardial ischemia-reperfusion injury. *Bioact Mater* 2021;6(7):2058-2069. doi:10.1016/j.bioactmat.2020.12.024, PMID:33511307.
- [129] Jabbari H, Roushandeh AM, Rostami MK, Razavi-Toosi MT, Shokrgozar MA, Jahani-Najafabadi A, et al. Mitochondrial transplantation ameliorates ischemia/reperfusion-induced kidney injury in rat. *Biochim Biophys Acta Mol Basis Dis* 2020;1866(8):165809. doi:10.1016/j.bbdis.2013.165809, PMID:32353613.
- [130] Zou X, Yan C, Shi Y, Cao K, Xu J, Wang X, et al. Mitochondrial dysfunction in obesity-associated nonalcoholic fatty liver disease: the protective effects of pomegranate with its active component punicalagin. *Antioxid Redox Signal* 2014;21(11):1557-1570. doi:10.1089/ars.2013.5538, PMID:24393106.
- [131] Zhang J, Chen F. Integrated transcriptome and metabolome study reveal the therapeutic effects of nicotinamide riboside and nicotinamide mononucleotide on nonalcoholic fatty liver disease. *Biomed Pharmacother* 2024;175:116701. doi:10.1016/j.biopha.2024.116701, PMID:38729053.
- [132] Ardekani A, Tabrizi R, Maleki E, Bagheri Lankarani K, Heydari ST, Moradinazar M, et al. Effects of coenzyme Q10 supplementation on lipid profiles and liver enzymes of nonalcoholic fatty liver disease (NAFLD) patients: A systematic review and meta-analysis of randomized controlled trials. *Food Sci Nutr* 2023;11(6):2580-2588. doi:10.1002/fsn3.3315, PMID:37324909.
- [133] Pang J, Yin L, Jiang W, Wang H, Cheng Q, Jiang Z, et al. Sirt1-mediated deacetylation of PGC-1 α alleviated hepatic steatosis in type 2 diabetes mellitus via improving mitochondrial fatty acid oxidation. *Cell Signal* 2024;124:111478. doi:10.1016/j.cellsig.2024.111478, PMID:39428026.
- [134] Kanazawa A, Tanaka A, Iwata S, Satoh S, Hatano E, Shinohara H, et al. The beneficial effect of phosphocreatine accumulation in the creatine kinase transgenic mouse liver in endotoxin-induced hepatic cell death. *J Surg Res* 1998;80(2):229-235. doi:10.1006/jsre.1998.5482, PMID:9878318.
- [135] Deng Y, Dong Y, Zhang S, Feng Y. Targeting mitochondrial homeostasis in the treatment of non-alcoholic fatty liver disease: a review. *Front Pharmacol* 2024;15:1463187. doi:10.3389/fphar.2024.1463187, PMID:39290869.
- [136] Sunny NE, Bril F, Cusi K. Mitochondrial Adaptation in Nonalcoholic Fatty Liver Disease: Novel Mechanisms and Treatment Strategies. *Trends Endocrinol Metab* 2017;28(4):250-260. doi:10.1016/j.tem.2016.11.006, PMID:27986466.
- [137] Mposhi A, Cortés-Mancera F, Heegsma J, de Meijer VE, van de Sluis B, Sydor S, et al. Mitochondrial DNA methylation in metabolic associated fatty liver disease. *Front Nutr* 2023;10:964337. doi:10.3389/fnut.2023.964337, PMID:37305089.
- [138] Priego-Parra BA, Gallego-Durán R, Román-Calleja BM, Velarde-Ruiz Velasco JA, Romero-Gómez M, Gracia-Sancho J. Advancing precision medicine in metabolic dysfunction-associated steatotic liver disease. *Trends Endocrinol Metab* 2025. doi:10.1016/j.tem.2025.03.006, PMID:40221323.
- [139] Longo M, Paolini E, Meroni M, Dongiovanni P. Remodeling of Mitochondrial Plasticity: The Key Switch from NAFLD/NASH to HCC. *Int J Mol Sci* 2021;22(8):4173. doi:10.3390/ijms22084173, PMID:33920670.
- [140] Chen Z, Tian R, She Z, Cai J, Li H. Role of oxidative stress in the pathogenesis of nonalcoholic fatty liver disease. *Free Radic Biol Med* 2020;152:116-141. doi:10.1016/j.freeradbiomed.2020.02.025, PMID:32156524.
- [141] Tao S, Cui D, Cheng H, Liu X, Jiang Z, Chen H, et al. High expression of TBRG4 in relation to unfavorable outcome and cell ferroptosis in hepatocellular carcinoma. *BMC Cancer* 2024;24(1):194. doi:10.1186/s12885-024-11943-1, PMID:38347489.
- [142] Wang H, Xu J, Li D, Zhang S, Guo Z. Identification of sequence polymorphisms in the mitochondrial cytochrome c oxidase genes as risk factors for hepatocellular carcinoma. *J Clin Lab Anal* 2018;32(3):e22299. doi:10.1002/jcla.22299, PMID:28703354.
- [143] Bi Y, Liu S, Qin X, Abudureyimu M, Wang L, Zou R, et al. FUNDC1 interacts with GPx4 to govern hepatic ferroptosis and fibrotic injury through a mitophagy-dependent manner. *J Adv Res* 2024;55:45-60. doi:10.1016/j.jare.2023.02.012, PMID:36828120.
- [144] Lu X, Xuan W, Li J, Yao H, Huang C, Li J. AMPK protects against alcohol-induced liver injury through UQCRC2 to up-regulate mitophagy. *Autophagy* 2021;17(11):3622-3643. doi:10.1080/15548627.2021.1886829, PMID:33719895.
- [145] Oaks Z, Winans T, Caza T, Fernandez D, Liu Y, Landas SK, et al. Mitochondrial Dysfunction in the Liver and Antiphospholipid Antibody Production Precede Disease Onset and Respond to Rapamycin in Lupus-Prone Mice. *Arthritis Rheumatol* 2016;68(11):2728-2739. doi:10.1002/art.39791, PMID:27332042.
- [146] Sun B, Ding P, Song Y, Zhou J, Chen X, Peng C, et al. FDX1 down-regulation activates mitophagy and the PI3K/AKT signaling pathway to promote hepatocellular carcinoma progression by inducing ROS produc-

- tion. *Redox Biol* 2024;75:103302. doi:10.1016/j.redox.2024.103302, PMID:39128228.
- [147] Gopal RK, Vantaku VR, Panda A, Reimer B, Rath S, To TL, *et al.* Effectors Enabling Adaptation to Mitochondrial Complex I Loss in Hürthle Cell Carcinoma. *Cancer Discov* 2023;13(8):1904–1921. doi:10.1158/2159-8290.CD-22-0976, PMID:37262067.
- [148] Munro B, Horvath R, Müller JS. Nucleoside supplementation modulates mitochondrial DNA copy number in the dguok $\pm$ zebrafish. *Hum Mol Genet* 2019;28(5):796–803. doi:10.1093/hmg/ddy389, PMID:30428046.
- [149] Fernandes MSS, Fidelis DEDS, Aidar FJ, Badicu G, Greco G, Cataldi S, *et al.* Coenzyme Q10 Supplementation in Athletes: A Systematic Review. *Nutrients* 2023;15(18):3990. doi:10.3390/nu15183990, PMID:37764774.
- [150] Gu YF, Chen YP, Jin R, Wang C, Wen C, Zhou YM. Age-related changes in liver metabolism and antioxidant capacity of laying hens. *Poult Sci* 2021;100(12):101478. doi:10.1016/j.psj.2021.101478, PMID:34695635.
- [151] Murali M, Shivanandappa T. Endosulfan causes oxidative stress in the liver and brain that involves inhibition of NADH dehydrogenase and altered antioxidant enzyme status in rat. *Ecotoxicol Environ Saf* 2022;239:113593. doi:10.1016/j.ecoenv.2022.113593, PMID:35567928.
- [152] Testai L, Martelli A, Flori L, Cicero AFG, Colletti A. Coenzyme Q(10): Clinical Applications beyond Cardiovascular Diseases. *Nutrients* 2021;13(5):1697. doi:10.3390/nu13051697, PMID:34067632.
- [153] Xu B, Zhang P, Tang X, Wang S, Shen J, Zheng Y, *et al.* Metabolic Rewiring of Kynurenine Pathway during Hepatic Ischemia-Reperfusion Injury Exacerbates Liver Damage by Impairing NAD Homeostasis. *Adv Sci (Weinh)* 2022;9(35):e2204697. doi:10.1002/advs.202204697, PMID:36310151.
- [154] Machado CM, de-Souza-Ferreira E, Silva GFS, Pimentel FSA, De-Souza EA, Silva-Rodrigues T, *et al.* Galactose-1-phosphate inhibits cytochrome c oxidase and causes mitochondrial dysfunction in classic galactosemia. *Biochim Biophys Acta Mol Basis Dis* 2024;1870(7):167340. doi:10.1016/j.bbdis.2024.167340, PMID:38986816.
- [155] Slaughter D, Takenaga N, Lu P, Assang C, Walsh DJ, Arison BH, *et al.* Metabolism of rofecoxib in vitro using human liver subcellular fractions. *Drug Metab Dispos* 2003;31(11):1398–1408. doi:10.1124/dmd.31.11.1398, PMID:14570773.
- [156] Luo P, Yan H, Du J, Chen X, Shao J, Zhang Y, *et al.* PLK1 (polo like kinase 1)-dependent autophagy facilitates gefitinib-induced hepatotoxicity by degrading COX6A1 (cytochrome c oxidase subunit 6A1). *Autophagy* 2021;17(10):3221–3237. doi:10.1080/15548627.2020.1851492, PMID:33315519.
- [157] Luo T, Yang S, Zhao T, Zhu H, Chen C, Shi X, *et al.* Hepatocyte DDX3X protects against drug-induced acute liver injury via controlling stress granule formation and oxidative stress. *Cell Death Dis* 2023;14(7):400. doi:10.1038/s41419-023-05913-x, PMID:37407573.
- [158] Mansouri A, Gattolliat CH, Asselah T. Mitochondrial Dysfunction and Signaling in Chronic Liver Diseases. *Gastroenterology* 2018;155(3):629–647. doi:10.1053/j.gastro.2018.06.083, PMID:30012333.
- [159] Yu L, Lin Z, Cheng X, Chu J, Li X, Chen C, *et al.* Thorium inhibits human respiratory chain complex IV (cytochrome c oxidase). *J Hazard Mater* 2022;424(Pt B):127546. doi:10.1016/j.jhazmat.2021.127546, PMID:34879532.
- [160] Boreel DF, Span PN, Heskamp S, Adema GJ, Bussink J. Targeting Oxidative Phosphorylation to Increase the Efficacy of Radio- and Immune-Combination Therapy. *Clin Cancer Res* 2021;27(11):2970–2978. doi:10.1158/1078-0432.CCR-20-3913, PMID:33419779.
- [161] Jolly RA, Goldstein KM, Wei T, Gao H, Chen P, Huang S, *et al.* Pooling samples within microarray studies: a comparative analysis of rat liver transcription response to prototypical toxicants. *Physiol Genomics* 2005;22(3):346–355. doi:10.1152/physiolgenomics.00260.2004, PMID:15914576.
- [162] Sharifabadi HR, Zamiri MJ, Rowghani E, Bottje WG. Relationship between the activity of mitochondrial respiratory chain complexes and feed efficiency in fat-tailed Ghezel lambs. *J Anim Sci* 2012;90(6):1807–1815. doi:10.2527/jas.2011-4791, PMID:22147474.