



Review Article

Emerging Serum Biomarkers for Chronic Hepatitis B: Focus on Serum HBV RNA and HBcrAg

Yike Tian¹, Haibo Yu^{1*} and Juan Chen^{1,2*}

¹Department of Infectious Diseases, Key Laboratory of Molecular Biology for Infectious Diseases (Ministry of Education), Institute for Viral Hepatitis, the Second Affiliated Hospital, Chongqing Medical University, Chongqing, China; ²College of Biomedical Engineering, Chongqing Medical University, Chongqing, China

Received: February 11, 2025 | Revised: May 22, 2025 | Accepted: July 07, 2025 | Published online: July 22, 2025

Abstract

Chronic hepatitis B virus (HBV) infection remains a major cause of liver diseases, including cirrhosis and hepatocellular carcinoma. Reliable biomarkers for assessing viral replication, liver damage, and predicting clinical outcomes are essential for effective patient management. This review focuses on two promising biomarkers: serum HBV RNA and hepatitis B core-related antigen, both of which show strong correlations with viral replication and disease progression. Serum HBV RNA levels reflect the quantity and transcriptional activity of intrahepatic covalently closed circular DNA, providing insights into viral replication. They also correlate with other markers of replicative activity and have predictive value for key clinical outcomes, including hepatitis B e antigen and hepatitis B surface antigen seroconversion, relapse after therapy cessation, and liver fibrosis. Similarly, hepatitis B core-related antigen is closely associated with covalently closed circular DNA levels, correlates with markers of viral replication, and shows promise in predicting liver fibrosis, cirrhosis, and the risk of hepatocellular carcinoma. This review highlights the potential of both biomarkers for monitoring disease progression and guiding therapeutic decisions, particularly in the context of personalized treatment strategies and risk assessment for liver-related complications.

Citation of this article: Tian Y, Yu H, Chen J. Emerging Serum Biomarkers for Chronic Hepatitis B: Focus on Serum HBV RNA and HBcrAg. J Clin Transl Hepatol 2025;13(9):766–775. doi: 10.14218/JCTH.2025.00064.

Introduction

Hepatitis B virus (HBV) remains a significant global health challenge, with over 250 million people worldwide living with chronic HBV infection despite the widespread promotion of

vaccination for more than 40 years.¹ If left untreated or inadequately managed, chronic HBV infection can progress to severe complications, including cirrhosis, liver failure, and hepatocellular carcinoma (HCC).² HBV is a hepatotropic virus with a 3.2-kb partially double-stranded relaxed circular DNA (rcDNA) genome.³ During the HBV life cycle, rcDNA is released into the nucleus, where it is converted into covalently closed circular DNA (cccDNA) via host DNA repair mechanisms.⁴ The cccDNA serves as a transcriptional template, generating a 3.5-kb pregenomic RNA (pgRNA) and multiple subgenomic RNAs (including preCore, preS1, preS2/S, and X mRNAs).⁴ The pgRNA is subsequently reverse-transcribed into rcDNA within nucleocapsids composed of hepatitis B core antigen (HBcAg). These nucleocapsids are then enveloped by hepatitis B surface antigen (HBsAg) to form mature virions, which are secreted from the cell.^{5,6} Notably, during pgRNA reverse transcription, 5–20% of transcripts undergo failed primer translocation, resulting in the formation of double-stranded linear DNA.⁷ These double-stranded linear DNA molecules can integrate into the host genome.^{8,9} Once integrated, they act as persistent transcriptional templates for HBsAg production,^{10,11} contributing to immune tolerance in the host.^{12–14}

Existing antiviral drugs cannot directly eliminate cccDNA or integrated HBV DNA within liver cells, making it very difficult to achieve a “complete cure” of hepatitis B.^{15–18} Given this limitation, one of the core goals of current antiviral treatment is to suppress the transcriptional activity of cccDNA and inhibit the functionality of integrated HBV DNA, ultimately achieving a “clinical cure” of hepatitis B.¹⁹ Therefore, accurately monitoring the activity levels of cccDNA and integrated HBV DNA is crucial for evaluating treatment efficacy.^{20–23} Since these viral forms are exclusively confined to hepatocytes, liver biopsy remains the gold standard for evaluation. However, its clinical utility is severely limited by invasiveness and the lack of standardized procedural guidelines.²⁴

Recent studies have demonstrated that serum HBV RNA and hepatitis B core-related antigen (HBcrAg) can reliably reflect the transcriptional activity of cccDNA, though their association with the transcriptional activity of integrated HBV DNA requires further investigation. Moreover, they have shown potential clinical value in predicting antiviral treatment response, assessing liver fibrosis progression, and estimating HCC risk, offering novel tools for the precision management of chronic HBV infection.^{25–28} Therefore, this review systematically summarizes the clinical applications and research ad-

Keywords: Chronic hepatitis B; Hepatitis B virus; Serum HBV RNA; Hepatitis B core-related antigen; Covalently closed circular DNA; Biomarkers.

***Correspondence to:** Haibo Yu and Juan Chen, Department of Infectious Diseases, Key Laboratory of Molecular Biology for Infectious Diseases (Ministry of Education), Institute for Viral Hepatitis, the Second Affiliated Hospital, Chongqing Medical University, Chongyi Building, 1 Yixueyuan Road, Yuzhong District, Chongqing 400016, China. ORCID: <https://orcid.org/0000-0002-7471-7923> (HY). Tel/Fax: +86-23-68486780, E-mail: 306842@hospital.cqmu.edu.cn (HY) and chenjuan2014@cqmu.edu.cn (JC)




Fig. 1. Production and clinical application of Serum HBV RNA and HBcrAg. Created with BioRender. cccDNA, covalently closed circular DNA; HBeAg, hepatitis B e antigen; HBsAg, hepatitis B surface antigen; HBV, hepatitis B virus; pgRNA, pregenomic RNA; rcDNA, relaxed circular DNA; p22cr, 22-kDa core-related antigen (precore protein).

vances of serum HBV RNA and HBcrAg in HBV infection and related diseases (Fig. 1).

Serum HBV RNA

In 1996, German researchers first discovered the presence of HBV RNA in the peripheral blood of chronic hepatitis B (CHB) patients.²⁹ These RNA molecules primarily consist of non- or partially reverse-transcribed pgRNA, with significant heterogeneity due to diverse post-transcriptional processing.³⁰ Current research confirms five predominant classes of serum HBV RNA: (1) Full-length pgRNA (3.5 kb)³¹: The major component, transcribed directly from cccDNA in the hepatocyte nucleus, containing the complete coding region with a polyadenylated tail.^{28,32–35} (2) 3'-Truncated pgRNA: Generated by incomplete template degradation mediated by the RNase H domain of the viral polymerase during reverse transcription; it lacks the 3' polyadenylated tail but retains the 5' ε packaging signal.^{15,36–38} Its proportion increases significantly with the duration of nucleos(t)ide analogue (NA) therapy.^{32,39} (3) pgRNA splicing variants: Arising from aberrant host cell splicing mechanisms, with at least 20 distinct secreted isoforms (sp1–sp20) identified^{15,36–38,40,41}; sp1 is the most abundant, constituting up to 30% of total pgRNA in HBV-infected HCC cell lines.^{16,42} (4) HBx transcripts: Generally present at low levels,^{15,28,43} including full-length or truncated HBx open reading frames as well as an ultra-long variant containing the polymerase (P) open reading frame.^{15,28,44} (5) HBV-human chimeric RNAs: Recently detected in serum samples from CHB patients harboring integrated HBV DNA. These include both 5'-HBV-human-3' and 5'-human-HBV-3' transcripts, although they represent only a minimal propor-

tion of total serum HBV RNA.^{28,45} Overall, the composition of serum HBV RNA is highly dynamic and complex. The specific RNA species present and their relative proportions are influenced by multiple factors, including infection stage, antiviral treatment status, and detection methodologies.³²

In addition, mounting evidence suggests that each distinct RNA species has its own clinical implications. For instance, splice variants can regulate HBV replication and promote the migration and invasion of liver cancer cells^{46–48}; 3'-truncated pgRNA, which accumulates during long-term NA therapy, is implicated in viral persistence^{32,39}; and the newly discovered HBV-human chimeric RNAs are closely associated with the development and progression of HCC.^{49,50} Nevertheless, current clinical research primarily focuses on detecting total serum pgRNA levels, which encompass full-length, 3'-truncated, and spliced variants.³⁰ This strategy is based on three key factors. Firstly, the total amount of pgRNA constitutes the majority of serum HBV RNA, providing a stable foundation for detection. Secondly, the relative proportions and contents of different components exhibit highly dynamic changes, and existing mainstream detection techniques⁵¹ make it difficult to analyze individual components accurately. Most importantly, as a direct transcription product of cccDNA, changes in serum pgRNA levels specifically reflect the transcriptional activity of cccDNA.³¹ Therefore, monitoring overall serum pgRNA levels is helpful for assessing viral replication status and the progression of HBV-related liver disease.^{52–54}

Serum HBV RNA quantification techniques

Quantitative detection of serum HBV RNA is critically important for understanding viral activity, assessing cccDNA transcriptional activity, and guiding clinical management. Cur-

rently, several molecular biology techniques are employed, primarily including reverse transcription (RT)-quantitative polymerase chain reaction (qPCR), RT-droplet digital polymerase chain reaction (PCR), fluorescence-based nucleic acid isothermal amplification testing (SAT), and rapid amplification of cDNA ends (RACE). These methods have distinct principles, advantages, and limitations, and differ in their capability to detect various HBV RNA isoforms.³⁰

Widely used serum HBV RNA quantification methods in clinical practice and research, such as RT-qPCR and SAT assays, cannot effectively differentiate between distinct HBV RNA isoforms.⁵¹ This limitation arises from their core technical principle: they target conserved, shared regions within the HBV genome (e.g., the X, preC/C, or S regions) rather than unique molecular markers specific to individual isoforms.^{31,55} RT-qPCR detects the sum of all RNA molecules containing the target sequence within the amplified region by reverse transcription and real-time amplification.⁵¹ While highly sensitive, this approach indiscriminately captures pgRNA, splice-generated SP1 variants, and truncated RNAs. SAT utilizes primers incorporating T7 promoters to achieve isothermal amplification, offering operational simplicity and eliminating the need for DNase treatment.⁵⁶ However, its amplification products similarly represent a mixed signal from all isoforms within the primer-binding region.

In contrast, RACE and droplet digital PCR possess inherent potential for isoform differentiation. RACE selectively amplifies polyadenylated transcripts by anchoring to the polyadenylated tail, enabling separation of full-length pgRNA generated by the canonical polyadenylation signal from truncated RNAs produced by the cryptic polyadenylation signal.^{57–60} However, this method cannot detect truncated RNAs lacking a polyadenylated tail. Droplet digital PCR leverages microfluidic partitioning for single-molecule isolation and Poisson distribution principles.³⁷ When combined with isoform-specific primer design, it enables absolute quantification of different splice variant proportions. Notably, conventional RT-qPCR also theoretically holds differentiation potential if designed with probes targeting isoform-specific markers; however, due to high sensitivity requirements, complex primer/probe design, and lack of standardization, it is rarely applied in routine clinical testing.^{51,61}

Given that distinct HBV RNA isoforms have unique biological properties and clinical significance, the inability of current mainstream methods to differentiate them obscures critical biological insights within composite detection signals.⁶² This represents a major obstacle to achieving precision medicine in HBV management. To overcome this bottleneck, future research must urgently focus on: first, developing novel detection platforms capable of highly sensitive, simultaneous differentiation of core isoforms; second, establishing international reference materials and unified detection standards encompassing these major isoforms to ensure result comparability and accelerate clinical translation.

Serum HBV RNA reflects cccDNA quantity and transcriptional activity

Unlike traditional HBV biomarkers such as HBV DNA and HBsAg, serum HBV RNA originates directly from the cccDNA template, making it a more precise indicator of intrahepatic cccDNA levels and transcriptional activity.⁶³ Wang *et al.* (2018) investigated this relationship in the natural history of CHB, finding a moderate correlation between serum HBV RNA and intrahepatic cccDNA ($r = 0.596$, $P < 0.001$).⁶⁴

Further studies have corroborated this correlation in untreated chronic HBV infections, with serum HBV RNA levels showing variable correlations with intrahepatic cccDNA ($r =$

$0.25–0.89$).^{58,64–66} These variations may arise from differences in patient characteristics or detection methodologies for circulating HBV RNA and intrahepatic cccDNA.⁶³

In patients undergoing antiviral therapy, serum HBV RNA also correlates with cccDNA levels. For example, in Peg-interferon-treated patients, serum HBV RNA demonstrated a stronger correlation with cccDNA than other HBV biomarkers after 48 weeks of treatment, irrespective of hepatitis B e antigen (HBeAg) seroconversion.⁶⁷ Similarly, in NAs-treated patients, serum HBV RNA, derived predominantly from cccDNA and minimally affected by antiviral drugs, has proven to be a reliable marker for monitoring cccDNA levels and activity.⁶⁸ Even in patients with suppressed HBV replication under NAs therapy, a significant correlation between serum HBV RNA and cccDNA transcriptional activity remains evident ($r = 0.78$, $P < 0.0001$).⁶⁹

In summary, serum HBV RNA, as a direct downstream product of cccDNA, serves as a robust biomarker reflecting intrahepatic cccDNA levels and activity. Its utility has been validated in both untreated CHB patients and those receiving antiviral therapy, highlighting its potential for infection monitoring and treatment response evaluation.

Correlation between serum HBV RNA and integrated DNA

In recent years, there has been growing research interest in the association between serum HBV RNA and integrated viral DNA. Studies reveal that integrated HBV DNA can transcribe two forms of replication-independent RNAs: (i) 5'-HBV-human-3' chimeric RNAs (integrant-derived RNAs, id-RNAs) initiated from viral promoters and polyadenylated using human polyadenylation signals, and (ii) 5'-human-HBV-3' RNAs initiated from upstream human promoters. Although this suggests the potential presence of integrant-derived envelope protein (cps) RNAs in serum, a systematic analysis of their complete molecular profiles remains lacking.²⁸ Furthermore, whether these RNAs accurately reflect the transcriptional activity and integration levels of integrated DNA remains unclear.

Given that the relevant molecular mechanisms have not been fully elucidated, an in-depth exploration of the characteristics of integrant-derived RNA in serum and its relationship with the transcriptional activity of viral integration sites will help elucidate the molecular mechanisms of HBV persistent infection and provide potential biomarkers for clinical monitoring.

Serum HBV RNA predicts HBeAg and HBsAg seroconversion

Serum HBV RNA levels and their dynamic changes are emerging as potential predictors of HBeAg and HBsAg seroconversion. During the immune-active phase, a rapid decline in serum HBV RNA strongly correlates with higher rates of spontaneous HBeAg seroconversion. Specifically, when HBV pgRNA levels at week 28 are below $5.63 \log_{10}$ copies/mL or when the reduction in HBV pgRNA from baseline exceeds $1.85 \log_{10}$ copies/mL, the likelihood of spontaneous HBeAg seroconversion within 48 weeks reaches approximately 87%. In contrast, patients with higher pgRNA levels or smaller reductions exhibit significantly lower conversion rates of 10–12%.⁷⁰

In treated patients, HBV RNA status after therapy also closely predicts HBeAg seroconversion. Studies have shown that patients remaining HBV RNA-positive after 48 weeks of NA therapy experience a prolonged time to seroconversion and a reduced likelihood of achieving it (hazard ratio (HR)

= 6.69, 95% CI: 1.88–23.84).⁷¹ Additionally, early dynamic changes in HBV RNA levels serve as reliable indicators of HBeAg seroconversion. For instance, HBV RNA levels at week 12 are significantly predictive of HBeAg seroconversion at 96 weeks of NAs therapy, with a threshold of 6.18 log₁₀ copies/mL (sensitivity 81%, specificity 80%, OR = 3.560, 95% CI: 1.39–9.110, *P* = 0.008). Fluctuations in HBV RNA at later stages, such as weeks 24 and 48, as well as three and six months post-treatment, further enhance predictive accuracy in HBeAg-positive patients.^{72–74}

Beyond HBeAg seroconversion, HBV RNA levels may also predict HBsAg seroconversion. HBV RNA-negative patients exhibit significantly lower quantitative HBsAg levels compared to HBV RNA-positive patients (2.2 vs. 3.1 log₁₀ IU/mL, *P* < 0.001).⁷⁵ Furthermore, patients with HBV RNA levels below 1,000 copies/mL at treatment cessation demonstrate a significantly higher cumulative HBsAg clearance rate over six years (30.9% vs. 1.6%, *P* = 0.007).⁷⁶

In conclusion, serum HBV RNA levels and their changes are useful predictors of both HBeAg and HBsAg seroconversion. These findings highlight the potential of serum HBV RNA as a valuable tool for guiding individualized treatment strategies and monitoring therapeutic outcomes in CHB.

Serum HBV RNA predicts relapse after therapy cessation

Current cessation criteria for HBV antiviral therapy continue to face clinical challenges, with over 40% of patients discontinuing treatment requiring reinitiation due to virological relapse.⁷⁷ The underlying mechanism lies in the persistent viral reservoir maintained by cccDNA, which serves as the fundamental source of post-treatment recurrence. Serum HBV RNA, as a direct transcriptional product of cccDNA, dynamically reflects the transcriptional activity of this viral reservoir in real time.⁵⁴ Leveraging this biological characteristic, multiple clinical investigations have demonstrated the potential utility of serum HBV RNA quantification in predicting post-therapeutic viral rebound and optimizing personalized treatment cessation strategies.

Multiple studies have consistently demonstrated a significant association between serum HBV RNA levels at the end of treatment (EOT) and the risk of virological relapse.^{78,79} A study involving 74 patients who completed one year of NAs therapy revealed a marked disparity in relapse rates between the EOT HBV RNA-negative and -positive groups (25.4% vs. 71%, *P* = 0.011).⁸⁰ This correlation was further quantified in a cohort of 114 entecavir-treated patients, where an HBV RNA cutoff ≥ 44.6 U/mL demonstrated over 90% predictive power for post-treatment relapse.⁸¹

Furthermore, emerging evidence supports the enhanced prognostic capacity of combined virological biomarker profiling for post-treatment recurrence prediction. Fan *et al.* conducted a four-year longitudinal follow-up study of 130 treatment-naïve HBeAg-positive patients, demonstrating that dual negativity of HBV DNA and RNA at EOT conferred a four-fold lower clinical relapse rate compared to double-positive cases (8.0% vs. 31.4%, *P* = 0.018).⁸² Innovatively, the Seto research group developed a novel composite criterion integrating HBV RNA negativity with HBsAg < 10 IU/mL, which reliably identifies candidates with minimal rebound risk post-cessation.⁸¹ Current studies suggest that integrating multidimensional viral markers may enable precise risk stratification in clinical practice. However, attention should be paid to the biological heterogeneity of different markers (e.g., HBV RNA reflecting cccDNA activity vs. HBsAg characterizing host immune response) when setting thresholds. Future validation through multicenter cohorts and dynamic marker combina-

tion models is warranted.

In summary, serum HBV RNA holds significant translational medical value in optimizing antiviral treatment endpoints for CHB and predicting recurrence risk after drug withdrawal. However, for routine clinical application, large-scale, multicenter prospective cohort studies are needed to address key issues: (1) determining optimal predictive thresholds at different treatment stages (e.g., NAs therapy, interferon therapy); (2) establishing standardized protocols for dynamic monitoring; and (3) validating combined biomarker prediction models across diverse populations. These efforts will provide a solid evidence base for developing individualized drug withdrawal strategies guided by biomarkers.

Serum HBV RNA assesses liver fibrosis

The early diagnosis and monitoring of liver fibrosis are critical for effective management of CHB. Although liver biopsy remains the gold standard for fibrosis evaluation, its invasiveness and risk of sampling errors limit routine clinical use.⁸³ Recent studies highlight serum HBV RNA as a non-invasive biomarker that correlates strongly with both the progression and regression of liver fibrosis, offering an alternative diagnostic tool.^{84–86}

Wang *et al.* (2017) found significant correlations between serum HBV RNA levels and histopathological scores for necroinflammation and fibrosis (*r* = 0.665, *P* < 0.001 for grading; *r* = 0.722, *P* < 0.001 for staging).⁸⁷ Using a cutoff value of 2.45 log₁₀ copies/mL, serum HBV RNA effectively differentiates samples with inflammation activity scores and fibrosis scores of <2 versus ≥2, achieving AUROCs of 0.88 and 0.85, respectively, surpassing the diagnostic accuracy of HBsAg.⁸⁷ Furthermore, Huang *et al.* (2020) demonstrated that serum HBV RNA levels are independent predictors of liver fibrosis in both HBeAg-positive (OR = 0.514, *P* < 0.001) and HBeAg-negative patients (OR = 3.574, *P* < 0.001), outperforming traditional indices such as APRI and FIB-4.⁸³

In addition to assessing fibrosis progression, serum HBV RNA levels are also effective in predicting fibrosis regression. Lower HBV RNA levels are observed in patients with regression compared to those without.^{84,85} Notably, a reduction in HBV RNA levels exceeding 0.63 log₁₀ copies/mL within the first six months of treatment predicts fibrosis regression at 60 months, with a sensitivity of 53.8% and specificity of 92.3%.⁸⁴ This likely reflects decreased cccDNA transcriptional activity in hepatocytes, a key factor in fibrosis resolution.^{65,88}

In summary, serum HBV RNA serves as a pivotal biomarker for evaluating liver fibrosis in CHB patients. Its ability to assess both fibrosis progression and regression provides valuable insights for optimizing treatment strategies and improving patient outcomes.

HBV RNA predicts HCC risk and prognosis

The strong relationship between HBV RNA and cccDNA has positioned serum HBV RNA as a promising biomarker for predicting HCC risk and prognosis in CHB patients. A 2021 case-control study from Hong Kong first demonstrated significantly higher serum HBV RNA positivity and levels in HCC patients compared to non-HCC patients during prior treatment (undetectable pgRNA: 9.6% vs. 36.5%, *P* < 0.001).⁸⁹ These findings were corroborated by a large prospective cohort study that established elevated serum HBV RNA levels as a significant risk factor for HCC in CHB patients.⁹⁰

In CHB patients undergoing long-term antiviral therapy with undetectable serum HBV DNA, elevated serum pgRNA levels were associated with poorer overall survival and higher

recurrence rates following hepatectomy. Conversely, pgRNA-negative patients exhibited significantly improved overall survival ($P < 0.001$).⁹¹ Cox multivariate analysis further identified high serum HBV RNA expression as an independent predictor of HCC recurrence, with HRs of 2.1 ($P = 0.003$) in Cohort A and 1.6 ($P = 0.033$) in Cohort B. The underlying mechanism may involve multiple carcinogenic pathways: (1) elevated HBV pgRNA levels promote expression of oncoproteins⁹¹; (2) HBV RNA can act as a microRNA sponge, sequestering and inhibiting tumor-suppressing host microRNAs, thereby promoting HCC growth and invasion.⁹²

Interestingly, patients with poorly differentiated HCC or lymphovascular invasion were found to have lower serum HBV RNA levels, particularly one year after diagnosis (1.71 [IQR 1.71–2.37] vs. 2.14 [IQR 1.71–3.59] $\log_{10}$ IU/mL, $P = 0.076$).^{89,93} This may be due to the high metabolic state in poorly differentiated HCC being unfavorable to HBV survival.⁹⁴

Overall, comprehensive analysis indicates that elevated serum HBV RNA levels are positively correlated with HCC risk, postoperative recurrence, and poor prognosis in CHB patients. However, patients with highly aggressive or poorly differentiated advanced HCC exhibit markedly reduced serum HBV RNA levels. This paradox suggests that the predictive value of HBV RNA may be disease stage-specific: during early HCC stages, active viral replication may directly drive tumorigenesis, whereas in advanced disease, dynamic tumor microenvironment remodeling might suppress viral replication, resulting in an inverse correlation between HBV RNA levels and tumor malignancy. Future studies should systematically elucidate the dynamic fluctuation patterns of HBV RNA according to HCC clinical staging and molecular subtyping. Additionally, developing multidimensional predictive models integrating clinicopathological features, imaging findings, and molecular biomarkers is essential to enhance precision in HCC risk prediction and prognosis evaluation.

HBcrAg

HBcrAg is a composite biomarker comprising three viral proteins—HBcAg, HBeAg, and the 22 kDa truncated precursor core protein—which share an identical 149-amino acid sequence but differ in their processing pathways.⁹⁵ HBcAg is produced by translation of the 3.5 kb pgRNA transcribed from cccDNA; HBeAg is generated via translation of precore mRNA into the precore protein, followed by N-terminal specific proteolytic processing to remove the signal peptide; and the 22 kDa truncated precursor core protein represents a distinct processing form of the precore protein undergoing both N-terminal and C-terminal modifications.^{96–100}

Because these proteins are almost entirely dependent on cccDNA-driven transcription and translation, HBcrAg quantitatively reflects the transcriptional activity of this viral reservoir, making it a potential clinical marker for evaluating HBV persistence and therapeutic efficacy.

HBcrAg correlates with intrahepatic cccDNA

HBcrAg has been demonstrated as a reliable surrogate marker for intrahepatic cccDNA due to its strong and consistent correlation with cccDNA levels.^{101–105} Studies show serum HBcrAg is significantly associated with intrahepatic cccDNA, independent of HBeAg status. In HBeAg-positive patients, multivariate regression analysis revealed that serum HBcrAg correlated more strongly with intrahepatic cccDNA than HBsAg ($\beta = 0.563$ vs. 0.328, both $P < 0.001$). In contrast, among HBeAg-negative patients, serum HBcrAg was the only biomarker significantly correlated with intrahepatic cccDNA

levels ($\beta = 0.774$, $P < 0.001$).¹⁰²

Further evidence indicates that, in patients undergoing NA therapy, reductions in serum HBcrAg closely parallel decreases in intrahepatic cccDNA. This correlation is stronger than that between serum HBsAg and intrahepatic cccDNA, both before and during treatment.^{101,103–105} While prolonged NA therapy often results in undetectable serum HBV DNA in most patients, serum HBcrAg remains detectable in approximately 78% of cases.¹⁰⁶ This persistence is attributed to NAs' ability to inhibit HBV DNA replication with minimal effect on cccDNA transcription and HBcrAg synthesis.¹⁰⁷ Even after five years of entecavir therapy, a moderate correlation between serum HBcrAg and intrahepatic cccDNA persists ($r = 0.419$, $P = 0.005$).^{106,108}

Overall, HBcrAg demonstrates superior and more consistent correlation with intrahepatic cccDNA compared to HBsAg and HBV DNA across diverse clinical contexts. These findings establish HBcrAg as a reliable surrogate marker for cccDNA in clinical practice. With the advent of novel therapies targeting cccDNA, HBcrAg holds significant potential as a non-invasive indicator for directly assessing cccDNA activity, thereby guiding precision therapeutic interventions.

HBcrAg correlates with other HBV markers reflecting replicative activity

Suzuki *et al.* (2019) first reported a significant correlation between HBcrAg and other virological markers in a study involving 57 CHB patients, observing a strong positive correlation between serum HBcrAg and HBV DNA levels ($r = 0.713$, $P < 0.001$).¹⁰⁹ Subsequent studies confirmed that this correlation remains consistent regardless of HBeAg status ($r = 0.59–0.85$, $P < 0.001$)^{104,110} or treatment history ($r = 0.786–0.820$, $P < 0.001$).^{105,111}

While the correlation between HBcrAg and HBV DNA is well established, evidence suggests a comparatively weaker relationship between HBcrAg and HBsAg. A cohort study of 2,666 patients in Taiwan found a strong correlation between HBcrAg and HBV DNA ($r = 0.83$, $P < 0.001$), whereas the correlation with HBsAg was moderate ($r = 0.59$, $P < 0.001$).¹¹² This discrepancy may be explained by the distinct origins of these markers: both HBcrAg and HBV DNA derive from cccDNA, whereas HBsAg can also originate from integrated viral genomes.¹¹³ Additionally, a moderate correlation between HBcrAg and HBeAg levels has been reported ($r = 0.491$, $P < 0.001$).¹⁰²

In summary, HBcrAg exhibits strong correlations with HBV DNA and moderate correlations with HBsAg and HBeAg, effectively reflecting HBV replicative activity. These findings highlight its potential utility in monitoring viral dynamics in CHB patients.

HBcrAg predicts HBeAg and HBsAg seroconversion

Emerging evidence suggests HBcrAg may serve as a supplementary biomarker for monitoring HBeAg and HBsAg seroconversion in CHB management. Studies show that patients with lower HBcrAg levels or significant reductions in HBcrAg are more likely to achieve HBeAg seroconversion, either spontaneously or during NA therapy.^{114,115} For example, at the third month of treatment, an HBcrAg level of 6.20 $\log_{10}$ U/mL moderately predicts HBeAg seroconversion (AU-ROC = 0.663).¹¹⁶ In patients undergoing combined NAs and interferon- α therapy, baseline HBcrAg levels $> 4.5 \log$ U/mL predict non-response and failure to achieve HBeAg seroconversion at 24 months ($P < 0.003$).¹¹⁷

Additionally, HBcrAg may serve as a valuable indicator in predicting HBsAg clearance. In cases of spontaneous HBsAg

clearance, 79% of patients had undetectable HBcrAg levels.^{118,119} Further research demonstrated that baseline HBcrAg levels are closely associated with virological response rates and HBsAg seroconversion in treated patients. Specifically, a baseline HBcrAg level of 2.550 log₁₀ U/mL moderately predicts HBsAg seroconversion (AUROC = 0.552).¹²⁰ At the EOT, HBcrAg levels below 2 log₁₀ U/mL significantly associate with higher virological response rates and HBsAg conversion ($P < 0.001$).¹²¹ These findings suggest that lower HBcrAg levels during treatment may indicate a greater likelihood of successful HBsAg conversion.

In conclusion, existing studies suggest HBcrAg levels provide supplementary insights for anticipating HBeAg and HBsAg seroconversion. Further investigations should focus on validating these findings through standardized multicenter studies, exploring integration with emerging biomarkers, and establishing long-term clinical correlations to refine its prognostic utility.

HBcrAg evaluates the risk of HBV recurrence after treatment

HBcrAg, as a marker of cccDNA transcriptional activity, has been extensively validated for its role in predicting HBV relapse after discontinuation of antiviral therapy. Numerous studies have demonstrated a strong correlation between baseline HBcrAg levels and the risk of virological relapse. A prospective study in Hong Kong revealed that the rate of HBV reactivation in patients with baseline HBcrAg positivity was significantly higher than in those with HBcrAg negativity (71.8% vs. 31%, $P = 0.002$). Multivariate analysis further identified baseline HBcrAg positivity as an independent risk factor for HBV reactivation ($P = 0.004$; HR = 2.94; 95% CI: 1.43–6.07).¹²²

In addition to baseline levels, HBcrAg at the EOT has emerged as a critical predictor of relapse risk. Studies consistently show that patients who experience reactivation after antiviral therapy have significantly higher HBcrAg levels at EOT compared to those who remain relapse-free (4.9 log U/mL vs. 3.2 log U/mL, $P = 0.009$).¹²³ Multiple analyses have confirmed that HBcrAg levels at EOT are an independent risk factor for HBV recurrence, with optimal cutoff values ranging from 3.7 to 4.0 log U/mL ($P = 0.002$ –0.024).^{124–126}

Shinkai *et al.* (2006) further highlighted the predictive value of HBcrAg at treatment cessation, showing that an HBcrAg level below 3.4 log U/mL at EOT was the only independent factor effectively predicting non-recurrence after therapy ($P = 0.042$).¹²⁷ Moreover, the Japanese Society of Hepatology has incorporated HBcrAg into clinical guidelines for relapse risk assessment, recommending a cutoff of <3.0 log U/mL to define low relapse risk.¹²⁸

In summary, both baseline and EOT HBcrAg levels are valuable for predicting HBV reactivation and recurrence. Regular monitoring of these levels enables clinicians to stratify patients by relapse risk, optimize treatment discontinuation strategies, and improve long-term clinical management.

HBcrAg predicts liver fibrosis and cirrhosis risk

Severe necroinflammatory activity is widely recognized as the initial stage in liver fibrosis progression. Studies have shown that HBcrAg independently predicts both significant necroinflammation ($P = 0.000$; OR = 2.290; 95% CI: 1.524–3.441) and significant liver fibrosis ($P = 0.000$; OR = 2.456; 95% CI: 1.631–3.699). The predictive accuracy for necroinflammation and fibrosis, measured by the area under the curve, is 0.807 (95% CI: 0.707–0.885) and 0.804 (95% CI: 0.703–0.883), respectively.¹²⁹ These associations

remain consistent across different HBeAg statuses (both $P < 0.001$).^{105,129,130}

Notably, HBcrAg demonstrates high diagnostic accuracy for liver fibrosis with optimal cutoff values specific to HBeAg status. For HBeAg-positive patients, a cutoff of $\leq 2.45 \times 10^4$ kU/mL effectively excludes severe lesions, while for HBeAg-negative patients, a cutoff of ≥ 4.02 kU/mL is more suitable for confirming significant liver damage (area under the curve > 0.70 for both). Based on the Youden index, HBcrAg thus serves distinct roles in fibrosis diagnosis depending on HBeAg status.¹²⁹

As liver fibrosis progresses, cirrhosis represents its end stage, often accompanied by severe liver damage and functional impairment.¹³¹ A Japanese study first established the association between HBcrAg levels and cirrhosis progression, identifying HBcrAg ≥ 3.7 log U/mL as an independent risk factor for cirrhosis (HR = 3.28; 95% CI: 1.60–6.75).¹³² In a long-term follow-up study of 1,673 CHB patients, Tseng *et al.* confirmed a significant correlation between elevated HBcrAg levels and cirrhosis risk ($P < 0.001$). Risk stratification analysis revealed that compared to patients with HBcrAg < 10 kU/mL, those with HBcrAg levels of 10–99 kU/mL and ≥ 100 kU/mL had HRs of 3.32 (95% CI: 1.99–5.52) and 7.35 (95% CI: 4.28–12.64), respectively.¹³³

In summary, HBcrAg is a promising non-invasive biomarker for assessing the risk of liver fibrosis and cirrhosis. Its strong predictive capabilities, especially when stratified by HBeAg status, highlight its clinical value. However, further research is needed to validate its applicability across diverse patient populations and clinical settings.

HBcrAg predicts HCC risk

HCC accounts for 75–85% of primary liver cancers and is associated with poor prognosis.¹³⁴ The limited sensitivity of traditional imaging and liver function tests for early diagnosis underscores the urgent need for novel serological markers.^{135,136} Recent studies have identified HBcrAg as a promising biomarker for predicting HCC risk, offering potential for early diagnosis and individualized risk assessment.

Kumada *et al.* (2013) first demonstrated a significant association between serum HBcrAg levels and HCC occurrence, reporting that elevated HBcrAg was independently associated with HCC development (HR = 2.77; 95% CI: 1.07–7.17; $P = 0.036$).¹³⁷ Subsequent studies have corroborated this finding, demonstrating that high serum HBcrAg levels are closely linked to HCC risk regardless of treatment status. For example, Tada *et al.* conducted a retrospective cohort study in untreated patients, revealing that baseline serum HBcrAg levels exceeding 2.9 log U/mL were associated with a fivefold increased risk of HCC compared to lower levels.¹³⁸

In patients receiving NA therapy, serum HBcrAg levels also strongly correlate with HCC risk. Research indicates that patients with serum HBcrAg levels exceeding 3.89 log U/mL after treatment exhibit a threefold higher risk of developing HCC.¹³⁹ Additionally, Hosaka *et al.* (2019) reported that patients with continuously rising HBcrAg levels during treatment face significantly greater HCC risk compared to those with stable or declining levels.¹⁴⁰

Overall, serum HBcrAg has been established as an independent predictive biomarker for HCC, with elevated levels significantly correlating with HCC risk in both untreated and antiviral-treated patients. Dynamic monitoring of HBcrAg facilitates evaluation of disease progression and prognosis, supporting early risk stratification. Future research should focus on integrating HBcrAg with other biomarkers to develop predictive models, while elucidating its mechanistic links to HBV-related carcinogenesis.

Conclusions

Serum HBcrAg and HBV RNA have emerged as promising biomarkers for the monitoring and prognostication of CHB. Their applications span a broad range of clinical scenarios, including predicting HBeAg and HBsAg seroconversion, assessing the risk of HBV recurrence following treatment cessation, evaluating liver fibrosis and cirrhosis, and forecasting the development of HCC. These biomarkers hold significant potential to advance personalized treatment strategies and improve clinical outcomes in CHB management. Despite their promise, further validation studies are necessary to establish standardized cutoff values and refine their clinical utility across diverse patient populations.

Funding

This work was supported by the National Key R&D Program of China (2022YFA1303600), the National Natural Science Foundation of China (Nos. 82273423, 82302507, and U23A20472), the Science Foundation of Chongqing (CST-B2023NSCQ-LZX0053), and the Joint Project of the Pinnacle Disciplinary Group of the Second Affiliated Hospital of Chongqing Medical University.

Conflict of interest

The authors have no conflict of interests related to this publication.

Author contributions

Conceptualization: YT; Drafting of the manuscript: YT; Critical review and editing of the manuscript: HY and JC. All authors have approved the final version and publication of the manuscript.

References

- [1] WHO. Hepatitis B. 9 April 2024. Available from: <https://www.who.int/news-room/fact-sheets/detail/hepatitis-b>.
- [2] Jeng WJ, Papatheodoridis GV, Lok ASF. Hepatitis B. *Lancet* 2023;401(10381):1039–1052. doi:10.1016/S0140-6736(22)01468-4, PMID:36774930.
- [3] Iannaccone M, Guidotti LG. Immunobiology and pathogenesis of hepatitis B virus infection. *Nat Rev Immunol* 2022;22(1):19–32. doi:10.1038/s41577-021-00549-4, PMID:34002067.
- [4] Gómez-Moreno A, Ploss A. Mechanisms of Hepatitis B Virus cccDNA and Minichromosome Formation and HBV Gene Transcription. *Viruses* 2024;16(4):609. doi:10.3390/v16040609, PMID:38675950.
- [5] Mendenhall MA, Hong X, Hu J. Hepatitis B Virus Capsid: The Core in Productive Entry and Covalently Closed Circular DNA Formation. *Viruses* 2023;15(3):642. doi:10.3390/v15030642, PMID:36992351.
- [6] Hong X, Schneider WM, Rice CM. Hepatitis B Virus Nucleocapsid Assembly. *J Mol Biol* 2025. doi:10.1016/j.jmb.2025.169182, PMID:40316009.
- [7] Caballero A, Tabernero D, Buti M, Rodríguez-Frias F. Hepatitis B virus: The challenge of an ancient virus with multiple faces and a remarkable replication strategy. *Antiviral Res* 2018;158:34–44. doi:10.1016/j.antiviral.2018.07.019, PMID:30059722.
- [8] Staprans S, Loeb DD, Ganem D. Mutations affecting hepatitis B virus plus-strand DNA synthesis dissociate primer cleavage from translocation and reveal the origin of linear viral DNA. *J Virol* 1991;65(3):1255–1262. doi:10.1128/JVI.65.3.1255-1262.1991, PMID:1704925.
- [9] Zhao XL, Yang JR, Lin SZ, Ma H, Guo F, Yang RF, *et al*. Serum viral duplex-linear DNA proportion increases with the progression of liver disease in patients infected with HBV. *Gut* 2016;65(3):502–511. doi:10.1136/gutjnl-2014-308989, PMID:26045139.
- [10] van Buuren N, Ramirez R, Soulette C, Suri V, Han D, May L, *et al*. Targeted long-read sequencing reveals clonally expanded HBV-associated chromosomal translocations in patients with chronic hepatitis B. *JHEP Rep* 2022;4(4):100449. doi:10.1016/j.jhepr.2022.100449, PMID:35295767.
- [11] Meier MA, Calabrese D, Suslov A, Terracciano LM, Heim MH, Wieland S. Ubiquitous expression of HBsAg from integrated HBV DNA in patients with low viral load. *J Hepatol* 2021;75(4):840–847. doi:10.1016/j.jhep.2021.04.051, PMID:34004216.
- [12] Le Bert N, Gill US, Hong M, Kunasegaran K, Tan DZM, Ahmad R, *et al*. Effects of Hepatitis B Surface Antigen on Virus-Specific and Global T

- Cells in Patients With Chronic Hepatitis B Virus infection. *Gastroenterology* 2020;159(2):652–664. doi:10.1053/j.gastro.2020.04.019, PMID:32302614.
- [13] Salimzadeh L, Le Bert N, Dutertre CA, Gill US, Newell EW, Frey C, *et al*. PD-1 blockade partially recovers dysfunctional virus-specific B cells in chronic hepatitis B infection. *J Clin Invest* 2018;128(10):4573–4587. doi:10.1172/JCI121957, PMID:30084841.
- [14] Fanning GC, Zoulim F, Hou J, Bertolotti A. Therapeutic strategies for hepatitis B virus infection: towards a cure. *Nat Rev Drug Discov* 2019;18(11):827–844. doi:10.1038/s41573-019-0037-0, PMID:31455905.
- [15] Vachon A, Seo GE, Patel NH, Coffin CS, Marinier E, Eyraas E, *et al*. Hepatitis B virus serum RNA transcript isoform composition and proportion in chronic hepatitis B patients by nanopore long-read sequencing. *Front Microbiol* 2023;14:1233178. doi:10.3389/fmicb.2023.1233178, PMID:37645229.
- [16] Lim CS, Sozzi V, Littlejohn M, Yuen LKW, Warner N, Betz-Stablein B, *et al*. Quantitative analysis of the splice variants expressed by the major hepatitis B virus genotypes. *Microb Genom* 2021;7(1):mgen000492. doi:10.1099/mgen.0.000492, PMID:33439114.
- [17] He W, Zheng Z, Zhao Q, Zhang R, Zheng H. Targeting HBV cccDNA Levels: Key to Achieving Complete Cure of Chronic Hepatitis B. *Pathogens* 2024;13(12):1100. doi:10.3390/pathogens13121100, PMID:39770359.
- [18] Kim SW, Yoon JS, Lee M, Cho Y. Toward a complete cure for chronic hepatitis B: Novel therapeutic targets for hepatitis B virus. *Clin Mol Hepatol* 2022;28(1):17–30. doi:10.3350/cmh.2021.0093, PMID:34281294.
- [19] Wong GLH, Gane E, Lok ASF. How to achieve functional cure of HBV: Stopping NUCs, adding interferon or new drug development? *J Hepatol* 2022;76(6):1249–1262. doi:10.1016/j.jhep.2021.11.024, PMID:35589248.
- [20] Nassal M. HBV cccDNA: viral persistence reservoir and key obstacle for a cure of chronic hepatitis B. *Gut* 2015;64(12):1972–1984. doi:10.1136/gutjnl-2015-309809, PMID:26048673.
- [21] Tu T, Budzinska MA, Shackel NA, Urban S. HBV DNA Integration: Molecular Mechanisms and Clinical Implications. *Viruses* 2017;9(4):E75. doi:10.3390/v9040075, PMID:28394272.
- [22] Xia Y, Guo H. Hepatitis B virus cccDNA: Formation, regulation and therapeutic potential. *Antiviral Res* 2020;180:104824. doi:10.1016/j.antiviral.2020.104824, PMID:32450266.
- [23] Allweiss L, Testoni B, Yu M, Lucifora J, Ko C, Qu B, *et al*. Quantification of the hepatitis B virus cccDNA: evidence-based guidelines for monitoring the key obstacle of HBV cure. *Gut* 2023;72(5):972–983. doi:10.1136/gutjnl-2022-328380, PMID:36707234.
- [24] Chowdhury AB, Mehta KJ. Liver biopsy for assessment of chronic liver diseases: a synopsis. *Clin Exp Med* 2023;23(2):273–285. doi:10.1007/s10238-022-00799-z, PMID:35192111.
- [25] Adraneda C, Tan YC, Yeo EJ, Kew GS, Khakpoor A, Lim SG. A critique and systematic review of the clinical utility of hepatitis B core-related antigen. *J Hepatol* 2023;78(4):731–741. doi:10.1016/j.jhep.2022.12.017, PMID:36586590.
- [26] Inoue T, Watanabe T, Tanaka Y. Hepatitis B core-related antigen: A novel and promising surrogate biomarker to guide anti-hepatitis B virus therapy. *Clin Mol Hepatol* 2023;29(4):851–868. doi:10.3350/cmh.2022.0434, PMID:36891607.
- [27] Lok J, Dusheiko G, Carey I, Agarwal K. Review article: novel biomarkers in hepatitis B infection. *Aliment Pharmacol Ther* 2022;56(5):760–776. doi:10.1111/apt.17105, PMID:35770458.
- [28] Zaiets I, Gunewardena S, Menne S, Weinman SA, Gudima SO. Sera of Individuals Chronically Infected with Hepatitis B Virus (HBV) Contain Diverse RNA Types Produced by HBV Replication or Derived from Integrated HBV DNA. *J Virol* 2023;97(3):e0195022. doi:10.1128/jvi.01950-22, PMID:36877036.
- [29] Köck J, Theilmann L, Galle P, Schlicht HJ. Hepatitis B virus nucleic acids associated with human peripheral blood mononuclear cells do not originate from replicating virus. *Hepatology* 1996;23(3):405–413. doi:10.1002/hep.510230303, PMID:8617418.
- [30] Venkatakrishnan B, Zlotnick A. The Structural Biology of Hepatitis B Virus: Form and Function. *Annu Rev Virol* 2016;3(1):429–451. doi:10.1146/annurev-virology-110615-042238, PMID:27482896.
- [31] Wang J, Shen T, Huang X, Kumar GR, Chen X, Zeng Z, *et al*. Serum hepatitis B virus RNA is encapsidated pregenome RNA that may be associated with persistence of viral infection and rebound. *J Hepatol* 2016;65(4):700–710. doi:10.1016/j.jhep.2016.05.029, PMID:27245431.
- [32] Su Q, Wang SF, Chang TE, Breikreutz R, Hennig H, Takegoshi K, *et al*. Circulating hepatitis B virus nucleic acids in chronic infection: representation of differently polyadenylated viral transcripts during progression to nonreplicative stages. *Clin Cancer Res* 2001;7(7):2005–2015. PMID:11448918.
- [33] Hacker HJ, Zhang W, Tokus M, Bock T, Schröder CH. Patterns of circulating hepatitis B virus serum nucleic acids during lamivudine therapy. *Ann N Y Acad Sci* 2004;1022:271–281. doi:10.1196/annals.1318.042, PMID:15251972.
- [34] Breikreutz R, Zhang W, Lee M, Hoffmann A, Tokus M, Su Q, *et al*. Hepatitis B virus nucleic acids circulating in the blood: distinct patterns in HBs carriers with hepatocellular carcinoma. *Ann N Y Acad Sci* 2001;945:195–206. doi:10.1111/j.1749-6632.2001.tb03886.x, PMID:11708479.
- [35] Prakash K, Rydell GE, Larsson SB, Andersson M, Norkrans G, Norder H, *et al*. High serum levels of pregenomic RNA reflect frequently failing reverse transcription in hepatitis B virus particles. *Virol J* 2018;15(1):86. doi:10.1186/s12985-018-0994-7, PMID:29764511.
- [36] Bai L, Zhang X, Kozłowski M, Li W, Wu M, Liu J, *et al*. Extracellular Hepatitis B Virus RNAs Are Heterogeneous in Length and Circulate as Capsid-Antibody Complexes in Addition to Virions in Chronic Hepatitis B Patients. *J Virol* 2018;92(24):e00798–18. doi:10.1128/JVI.00798-18, PMID:30282709.
- [37] Shen S, Xie Z, Cai D, Yu X, Zhang H, Kim ES, *et al*. Biogenesis and mo-

- lecular characteristics of serum hepatitis B virus RNA. *PLoS Pathog* 2020;16(10):e1008945. doi:10.1371/journal.ppat.1008945, PMID:33079954.
- [38] Shen S, Liu W, Zeng G, Liang H, Yu X, Zhang H, *et al*. Conditional replication and secretion of hepatitis B virus genome uncover the truncated 3' terminus of encapsidated viral pregenomic RNA. *J Virol* 2023;97(10):e0076023. doi:10.1128/jvi.00760-23, PMID:37754759.
- [39] Kairat A, Beerheide W, Zhou G, Tang ZY, Edler L, Schröder CH. Truncated hepatitis B virus RNA in human hepatocellular carcinoma: its representation in patients with advancing age. *Intervirol* 1999;42(4):228–237. doi:10.1159/000024982, PMID:10567841.
- [40] Lam AM, Ren S, Espiritu C, Kelly M, Lau V, Zheng L, *et al*. Hepatitis B Virus Capsid Assembly Modulators, but Not Nucleoside Analogs, Inhibit the Production of Extracellular Pregenomic RNA and Spliced RNA Variants. *Antimicrob Agents Chemother* 2017;61(8):e00680–17. doi:10.1128/AAC.00680-17, PMID:28559265.
- [41] Wang J, Sheng Q, Ding Y, Chen R, Sun X, Chen X, *et al*. HBV RNA virion-like particles produced under nucleos(t)ide analogues treatment are mainly replication-deficient. *J Hepatol* 2018;68(4):847–849. doi:10.1016/j.jhep.2017.10.030, PMID:29113914.
- [42] Wu HL, Chen PJ, Tu SJ, Lin MH, Lai MY, Chen DS. Characterization and genetic analysis of alternatively spliced transcripts of hepatitis B virus in infected human liver tissues and transfected HepG2 cells. *J Virol* 1991;65(4):1680–1686. doi:10.1128/JVI.65.4.1680-1686.1991, PMID:1705988.
- [43] Stadelmayer B, Diederichs A, Chapus F, Rivoire M, Neveu G, Alam A, *et al*. Full-length 5'RACE identifies all major HBV transcripts in HBV-infected hepatocytes and patient serum. *J Hepatol* 2020;73(1):40–51. doi:10.1016/j.jhep.2020.01.028, PMID:32087349.
- [44] Niu C, Livingston CM, Li L, Beran RK, Daffis S, Ramakrishnan D, *et al*. The Smc5/6 Complex Restricts HBV when Localized to ND10 without Inducing an Innate Immune Response and Is Counteracted by the HBV X Protein Shortly after Infection. *PLoS One* 2017;12(1):e0169648. doi:10.1371/journal.pone.0169648, PMID:28095508.
- [45] Chang S, Hedskog C, Parhy B, Martin R, Mo H, Maiorova E, *et al*. Sequence characterization of extracellular HBV RNA in patient plasma. *J Viral Hepat* 2023;30(1):29–38. doi:10.1111/jvh.13760, PMID:36208116.
- [46] Soussan P, Garreau F, Zylberberg H, Ferray C, Brechot C, Kremsdorf D. In vivo expression of a new hepatitis B virus protein encoded by a spliced RNA. *J Clin Invest* 2000;105(1):55–60. doi:10.1172/JCI8098, PMID:10619861.
- [47] Soussan P, Pol J, Garreau F, Schneider V, Le Pendeven C, Nalpas B, *et al*. Expression of defective hepatitis B virus particles derived from singly spliced RNA is related to liver disease. *J Infect Dis* 2008;198(2):218–225. doi:10.1086/589623, PMID:18532883.
- [48] Chen WN, Chen JY, Jiao BY, Lin WS, Wu YL, Liu LL, *et al*. Interaction of the hepatitis B spliced protein with cathepsin B promotes hepatoma cell migration and invasion. *J Virol* 2012;86(24):13533–13541. doi:10.1128/JVI.02095-12, PMID:23035214.
- [49] Lau CC, Sun T, Ching AK, He M, Li JW, Wong AM, *et al*. Viral-human chimeric transcript predisposes host to liver cancer development and progression. *Cancer Cell* 2014;25(3):335–349. doi:10.1016/j.ccr.2014.01.030, PMID:24582836.
- [50] Liang HW, Wang N, Wang Y, Wang F, Fu Z, Yan X, *et al*. Hepatitis B virus-human chimeric transcript HBx-LINE1 promotes hepatic injury via sequestering cellular microRNA-122. *J Hepatol* 2016;64(2):278–291. doi:10.1016/j.jhep.2015.09.013, PMID:26409216.
- [51] Gao M, Feng C, Ying R, Nie Y, Deng X, Zhu Y, *et al*. A novel one-step quantitative reverse transcription PCR assay for selective amplification of hepatitis B virus pregenomic RNA from a mixture of HBV DNA and RNA in serum. *Arch Virol* 2019;164(11):2683–2690. doi:10.1007/s00705-019-04372-0, PMID:31428915.
- [52] Lu F, Wang J, Chen X, Xu D, Xia N. Potential use of serum HBV RNA in antiviral therapy for chronic hepatitis B in the era of nucleos(t)ide analogs. *Front Med* 2017;11(4):502–508. doi:10.1007/s11684-017-0590-z, PMID:29170915.
- [53] Wu Y, Wen J, Xiao W, Zhang B. Pregenomic RNA: How to assist the management of chronic hepatitis B? *Rev Med Virol* 2019;29(4):e2051. doi:10.1002/rmv.2051, PMID:31074177.
- [54] Liu S, Zhou B, Valdes JD, Sun J, Guo H. Serum Hepatitis B Virus RNA: A New Potential Biomarker for Chronic Hepatitis B Virus Infection. *Hepatology* 2019;69(4):1816–1827. doi:10.1002/hep.30325, PMID:30362148.
- [55] Rokuhara A, Matsumoto A, Tanaka E, Umemura T, Yoshizawa K, Kimura T, *et al*. Hepatitis B virus RNA is measurable in serum and can be a new marker for monitoring lamivudine therapy. *J Gastroenterol* 2006;41(8):785–790. doi:10.1007/s00535-006-1856-4, PMID:16988768.
- [56] Hu X, Zhao L, Ou M, Chen Y, Wei H, Xia Y, *et al*. Evaluation of reverse transcription-polymerase chain reaction and simultaneous amplification and testing for quantitative detection of serum hepatitis B virus RNA. *Heliyon* 2023;9(8):e18557. doi:10.1016/j.heliyon.2023.e18557, PMID:37560627.
- [57] Serum hepatitis B virus RNA levels as an early predictor of hepatitis B envelope antigen seroconversion during treatment with polymerase inhibitors. *Hepatology* 2016;63(1):349. doi:10.1002/hep.28349, PMID:26689731.
- [58] Gao Y, Li Y, Meng Q, Zhang Z, Zhao P, Shang Q, *et al*. Serum Hepatitis B Virus DNA, RNA, and HBsAg: Which Correlated Better with Intrahepatic Covalently Closed Circular DNA before and after Nucleos(t)ide Analogue Treatment? *J Clin Microbiol* 2017;55(10):2972–2982. doi:10.1128/JCM.00760-17, PMID:28747369.
- [59] van Bömmel F, van Bömmel A, Krauel A, Wat C, Pavlovic V, Yang L, *et al*. Serum HBV RNA as a Predictor of Peginterferon Alfa-2a Response in Patients With HBeAg-Positive Chronic Hepatitis B. *J Infect Dis* 2018;218(7):1066–1074. doi:10.1093/infdis/jiy270, PMID:29741634.
- [60] van Campenhout MJH, van Bömmel F, Pfefferkorn M, Fischer J, Deichsel D, Boonstra A, *et al*. Host and viral factors associated with serum hepatitis B virus RNA levels among patients in need for treatment. *Hepatology* 2018;68(3):839–847. doi:10.1002/hep.29872, PMID:29514389.
- [61] Nguyen UD, Le Do Q, Vu QAN, Trieu NT, Dao TT, Van Le N, *et al*. Selective detection of HBV pre-genomic RNA in chronic hepatitis B patients using a novel RT-PCR assay. *Clin Exp Med* 2023;23(8):5281–5289. doi:10.1007/s10238-023-01162-6, PMID:37572154.
- [62] Yan R, Cai D, Ouyang L, Colonno R, Huang Q, Kitrinis KM. Development of a sensitive, multi-assay platform to monitor low levels of HBV DNA and pgRNA in patients with chronic hepatitis B virus infection. *J Virol Methods* 2023;311:114640. doi:10.1016/j.jviromet.2022.114640, PMID:36332714.
- [63] Deng R, Liu S, Shen S, Guo H, Sun J. Circulating HBV RNA: From biology to clinical applications. *Hepatology* 2022;76(5):1520–1530. doi:10.1002/hep.32479, PMID:35342969.
- [64] Wang J, Yu Y, Li G, Shen C, Li J, Chen S, *et al*. Natural history of serum HBV-RNA in chronic HBV infection. *J Viral Hepat* 2018;25(9):1038–1047. doi:10.1111/jvh.12908, PMID:29633430.
- [65] Giersch K, Allweiss L, Volz T, Dandri M, Lütgehetmann M. Serum HBV pgRNA as a clinical marker for cccDNA activity. *J Hepatol* 2017;66(2):460–462. doi:10.1016/j.jhep.2016.09.028, PMID:27826059.
- [66] Huang H, Wang J, Li W, Chen R, Chen X, Zhang F, *et al*. Serum HBV DNA plus RNA shows superiority in reflecting the activity of intrahepatic cccDNA in treatment-naïve HBV-infected individuals. *J Clin Virol* 2018;99–100:71–78. doi:10.1016/j.jcv.2017.12.016, PMID:29353073.
- [67] Wang X, Chi X, Wu R, Xu H, Gao X, Yu L, *et al*. Serum HBV RNA correlated with intrahepatic cccDNA more strongly than other HBV markers during peg-interferon treatment. *Virol J* 2021;18(1):4. doi:10.1186/s12985-020-01471-2, PMID:33407619.
- [68] Liu Y, Jiang M, Xue J, Yan H, Liang X. Serum HBV RNA quantification: useful for monitoring natural history of chronic hepatitis B infection. *BMC Gastroenterol* 2019;19(1):53. doi:10.1186/s12876-019-0966-4, PMID:30991954.
- [69] Testoni B, Scholtès C, Plissonnier ML, Patrel A, Berby F, Facchetti F, *et al*. Quantification of circulating HBV RNA expressed from intrahepatic cccDNA in untreated and NUC treated patients with chronic hepatitis B. *Gut* 2024;73(4):659–667. doi:10.1136/gutjnl-2023-330644, PMID:37879886.
- [70] Song G, Yang R, Jin Q, Liu J, Rao H, Feng B, *et al*. HBV pregenome RNA as a predictor of spontaneous HBeAg seroconversion in HBeAg-positive chronic hepatitis B patients. *BMC Gastroenterol* 2023;23(1):381. doi:10.1186/s12876-023-03023-8, PMID:37946120.
- [71] Luo H, Tan N, Kang Q, Pan J, Chen H, Xi H, *et al*. Hepatitis B virus pregenomic RNA status can reveal the long-term prognoses of chronic hepatitis B patients treated with nucleos(t)ide analogues. *J Viral Hepat* 2020;27(3):323–328. doi:10.1111/jvh.13227, PMID:31667945.
- [72] Ji X, Xia M, Zhou B, Liu S, Liao G, Cai S, *et al*. Serum Hepatitis B Virus RNA Levels Predict HBeAg Seroconversion and Virological Response in Chronic Hepatitis B Patients with High Viral Load Treated with Nucleos(t)ide Analog. *Infect Drug Resist* 2020;13:1881–1888. doi:10.2147/IDR.S252994, PMID:32606837.
- [73] van Bömmel F, Bartens A, Mysickova A, Hofmann J, Krüger DH, Berg T, *et al*. Serum hepatitis B virus RNA levels as an early predictor of hepatitis B envelope antigen seroconversion during treatment with polymerase inhibitors. *Hepatology* 2015;61(1):66–76. doi:10.1002/hep.27381, PMID:25132147.
- [74] Ye F, Zhao W, Yang X, Zhang X, An X, Zhu R, *et al*. The decline of HBV RNA associated with HBeAg seroconversion and double-negative HBV DNA and RNA in chronic hepatitis B patients who received entecavir therapy: a 10-year retrospective cohort study. *Ann Transl Med* 2022;10(16):897. doi:10.21037/atm-22-3265, PMID:36110993.
- [75] Fan R, Peng J, Xie Q, Tan D, Xu M, Niu J, *et al*. Combining Hepatitis B Virus RNA and Hepatitis B Core-Related Antigen: Guidance for Safely Stopping Nucleos(t)ide Analogues in Hepatitis B e Antigen-Positive Patients With Chronic Hepatitis B. *J Infect Dis* 2020;222(4):611–618. doi:10.1093/infdis/jiaa136, PMID:32211776.
- [76] Xia M, Chi H, Wu Y, Hansen BE, Li Z, Liu S, *et al*. Serum hepatitis B virus RNA level is associated with biochemical relapse in patients with chronic hepatitis B infection who discontinue nucleos(t)ide analogue treatment. *Aliment Pharmacol Ther* 2021;54(5):709–714. doi:10.1111/apt.16538, PMID:34275138.
- [77] Kao JH, Jeng WJ, Ning Q, Su TH, Tseng TC, Ueno Y, *et al*. APASL guidance on stopping nucleos(t)ide analogues in chronic hepatitis B patients. *Hepatol Int* 2021;15(4):833–851. doi:10.1007/s12072-021-10223-5, PMID:34297329.
- [78] Brakenhoff SM, de Knegt RJ, van Campenhout MJH, van der Eijk AA, Brouwer WP, van Bömmel F, *et al*. End-of-treatment HBsAg, HBcrAg and HBV RNA predict the risk of off-treatment ALT flares in chronic hepatitis B patients. *J Microbiol Immunol Infect* 2023;56(1):31–39. doi:10.1016/j.jmii.2022.06.002, PMID:35941076.
- [79] Carey I, Gersch J, Wang B, Moigboi C, Kuhns M, Cloherty G, *et al*. Pregenomic HBV RNA and Hepatitis B Core-Related Antigen Predict Outcomes in Hepatitis B e Antigen-Negative Chronic Hepatitis B Patients Suppressed on Nucleos(T)ide Analogue Therapy. *Hepatology* 2020;72(1):42–57. doi:10.1002/hep.31026, PMID:31701544.
- [80] Laras A, Papatheodoridi M, Panopoulou E, Papatheodoridis GV, Hadziyannis SJ, Hadziyannis E. Serum hepatitis B virus RNA detectability, composition and clinical significance in patients with ab initio hepatitis B e antigen negative chronic hepatitis B. *Virol J* 2022;19(1):22. doi:10.1186/s12985-022-01749-7, PMID:35093105.
- [81] Seto WK, Liu KS, Mak LY, Cloherty G, Wong DK, Gersch J, *et al*. Role of serum HBV RNA and hepatitis B surface antigen levels in identifying Asian

- patients with chronic hepatitis B suitable for entecavir cessation. *Gut* 2021;70(4):775–783. doi:10.1136/gutjnl-2020-321116, PMID:32759300.
- [82] Fan R, Zhou B, Xu M, Tan D, Niu J, Wang H, *et al*. Association Between Negative Results From Tests for HBV DNA and RNA and Durability of Response After Discontinuation of Nucleos(t)ide Analogue Therapy. *Clin Gastroenterol Hepatol* 2020;18(3):719–727.e7. doi:10.1016/j.cgh.2019.07.046, PMID:31362119.
- [83] Huang C, Li Q, Xu W, Wang Q, Hu Q, Zhang X, *et al*. Serum HBV RNA levels predict significant liver fibrosis in patients with chronic HBV infection. *Discov Med* 2020;29(157):119–128. PMID:33002408.
- [84] Bian D, Zhao J, Liao H, Wang Y, Ren Y, Jiang Y, *et al*. Serum HBV RNA is associated with liver fibrosis regression in HBeAg-positive chronic hepatitis B patients treated with nucleos(t)ide analogues. *J Viral Hepat* 2023;30(4):303–309. doi:10.1111/jvh.13790, PMID:36533536.
- [85] Sun Y, Wu X, Zhou J, Meng T, Wang B, Chen S, *et al*. Persistent Low Level of Hepatitis B Virus Promotes Fibrosis Progression During Therapy. *Clin Gastroenterol Hepatol* 2020;18(11):2582–2591.e6. doi:10.1016/j.cgh.2020.03.001, PMID:32147592.
- [86] Xu Q, Ding H, Bai T, Huang R, Wang J, Zhang J, *et al*. Serum HBV RNA levels among untreated adults with chronic hepatitis B in distinct immune phases and liver histopathology statuses. *J Mol Histol* 2023;54(6):739–749. doi:10.1007/s10735-023-10162-5, PMID:37843699.
- [87] Wang J, Yu Y, Li G, Shen C, Meng Z, Zheng J, *et al*. Relationship between serum HBV-RNA levels and intrahepatic viral as well as histologic activity markers in entecavir-treated patients. *J Hepatol* 2017;68(1):16–24. doi:10.1016/j.jhep.2017.08.021, PMID:28870671.
- [88] Wang J, Du M, Huang H, Chen R, Niu J, Jiang J, *et al*. Reply to: “Serum HBV pgRNA as a clinical marker for cccDNA activity”: Consistent loss of serum HBV RNA might predict the “para-functional cure” of chronic hepatitis B. *J Hepatol* 2017;66(2):462–463. doi:10.1016/j.jhep.2016.10.034, PMID:27826054.
- [89] Mak LY, Huang Q, Wong DK, Stamm L, Cheung KS, Ko KL, *et al*. Residual HBV DNA and pgRNA viraemia is associated with hepatocellular carcinoma in chronic hepatitis B patients on antiviral therapy. *J Gastroenterol* 2021;56(5):479–488. doi:10.1007/s00535-021-01780-5, PMID:33772643.
- [90] Hilger C, Velhagen I, Zentgraf H, Schröder CH. Diversity of hepatitis B virus X gene-related transcripts in hepatocellular carcinoma: a novel polyadenylation site on viral DNA. *J Virol* 1991;65(8):4284–4291. doi:10.1128/JVI.65.8.4284-4291.1991, PMID:1649331.
- [91] Ding WB, Wang MC, Yu J, Huang G, Sun DP, Liu L, *et al*. HBV/Pregenomic RNA Increases the Stemness and Promotes the Development of HBV-Related HCC Through Reciprocal Regulation With Insulin-Like Growth Factor 2 mRNA-Binding Protein 3. *Hepatology* 2021;74(3):1480–1495. doi:10.1002/hep.31850, PMID:33825218.
- [92] Wang S, Qiu L, Yan X, Jin W, Wang Y, Chen L, *et al*. Loss of microRNA 122 expression in patients with hepatitis B enhances hepatitis B virus replication through cyclin G(1)-modulated P53 activity. *Hepatology* 2012;55(3):730–741. doi:10.1002/hep.24809, PMID:22105316.
- [93] Halgand B, Desterke C, Rivière R, Fallot G, Sebagh M, Calderaro J, *et al*. Hepatitis B Virus Pregenomic RNA in Hepatocellular Carcinoma: A Nosological and Prognostic Determinant. *Hepatology* 2018;67(1):86–96. doi:10.1002/hep.29463, PMID:28802063.
- [94] Ding W-B, Wang M-C, Zhang J-N, Sun D-P, Dong J-P, Zhou W-P, *et al*. Novel insights of HBV RNA in hepatitis B virus pathogenesis and clinical application. *Hepato Res* 2019;5(0):10. doi:10.20517/2394-5079.2018.115.
- [95] Hong X, Luckenbaugh L, Mendenhall M, Walsh R, Cabuang L, Soppe S, *et al*. Characterization of Hepatitis B Precore/Core-Related Antigens. *J Virol* 2021;95(3):e01695–20. doi:10.1128/JVI.01695-20, PMID:33148795.
- [96] Wu JW, Kao JH, Tseng TC. Three heads are better than two: Hepatitis B core-related antigen as a new predictor of hepatitis B virus-related hepatocellular carcinoma. *Clin Mol Hepatol* 2021;27(4):524–534. doi:10.3350/cmh.2021.0012, PMID:33618507.
- [97] Mak LY, Wong DK, Cheung KS, Seto WK, Lai CL, Yuen MF. Review article: hepatitis B core-related antigen (HBcrAg): an emerging marker for chronic hepatitis B virus infection. *Aliment Pharmacol Ther* 2018;47(1):43–54. doi:10.1111/apt.14376, PMID:29035003.
- [98] Vachon A, Osiowy C. Novel Biomarkers of Hepatitis B Virus and Their Use in Chronic Hepatitis B Patient Management. *Viruses* 2021;13(6):951. doi:10.3390/v13060951, PMID:34064049.
- [99] Baudi I, Inoue T, Tanaka Y. Novel Biomarkers of Hepatitis B and Hepatocellular Carcinoma: Clinical Significance of HBcrAg and M2BPGI. *Int J Mol Sci* 2020;21(3):949. doi:10.3390/ijms21030949, PMID:32023902.
- [100] Tsukuda S, Watashi K. Hepatitis B virus biology and life cycle. *Antiviral Res* 2020;182:104925. doi:10.1016/j.antiviral.2020.104925, PMID:32866519.
- [101] Chen EQ, Feng S, Wang ML, Liang LB, Zhou LY, Du LY, *et al*. Serum hepatitis B core-related antigen is a satisfactory surrogate marker of intrahepatic covalently closed circular DNA in chronic hepatitis B. *Sci Rep* 2017;7(1):173. doi:10.1038/s41598-017-00111-0, PMID:28282964.
- [102] Chen EQ, Wang ML, Tao YC, Wu DB, Liao J, He M, *et al*. Serum HBcrAg is better than HBV RNA and HBsAg in reflecting intrahepatic covalently closed circular DNA. *J Viral Hepat* 2019;26(5):586–595. doi:10.1111/jvh.13061, PMID:30632235.
- [103] Chen S, Jia J, Gao Y, Li H, Fang M, Feng H, *et al*. Clinical evaluation of hepatitis B core-related antigen in chronic hepatitis B and hepatocellular carcinoma patients. *Clin Chim Acta* 2018;486:237–244. doi:10.1016/j.cca.2018.07.027, PMID:30025756.
- [104] Wong DK, Seto WK, Cheung KS, Chong CK, Huang FY, Fung J, *et al*. Hepatitis B virus core-related antigen as a surrogate marker for covalently closed circular DNA. *Liver Int* 2017;37(7):995–1001. doi:10.1111/liv.13346, PMID:27992681.
- [105] Wong DK, Tanaka Y, Lai CL, Mizokami M, Fung J, Yuen MF. Hepatitis B virus core-related antigens as markers for monitoring chronic hepatitis B infection. *J Clin Microbiol* 2007;45(12):3942–3947. doi:10.1128/JCM.00366-07, PMID:17942661.
- [106] Suzuki F, Miyakoshi H, Kobayashi M, Kumada H. Correlation between serum hepatitis B virus core-related antigen and intrahepatic covalently closed circular DNA in chronic hepatitis B patients. *J Med Virol* 2009;81(1):27–33. doi:10.1002/jmv.21339, PMID:19031469.
- [107] Broquetas T, Carrión JA. Current Perspectives on Nucleos(t)ide Analogue Therapy for the Long-Term Treatment of Hepatitis B Virus. *Hepat Med* 2022;14:87–100. doi:10.2147/HMER.S291976, PMID:35936810.
- [108] Testoni B, Lebossé F, Scholtes C, Berby F, Miaglia C, Subic M, *et al*. Serum hepatitis B core-related antigen (HBcrAg) correlates with covalently closed circular DNA transcriptional activity in chronic hepatitis B patients. *J Hepatol* 2019;70(4):615–625. doi:10.1016/j.jhep.2018.11.030, PMID:30529504.
- [109] Inoue T, Tanaka Y. The Role of Hepatitis B Core-Related Antigen. *Genes (Basel)* 2019;10(5):357. doi:10.3390/genes10050357, PMID:31075974.
- [110] Rokuhara A, Tanaka E, Matsumoto A, Kimura T, Yamaura T, Orii K, *et al*. Clinical evaluation of a new enzyme immunoassay for hepatitis B virus core-related antigen; a marker distinct from viral DNA for monitoring lamivudine treatment. *J Viral Hepat* 2003;10(4):324–330. doi:10.1046/j.1365-2893.2003.00437.x, PMID:12823601.
- [111] Rokuhara A, Sun X, Tanaka E, Kimura T, Matsumoto A, Yao D, *et al*. Hepatitis B virus core and core-related antigen quantitation in Chinese patients with chronic genotype B and C hepatitis B virus infection. *J Gastroenterol Hepatol* 2005;20(11):1726–1730. doi:10.1111/j.1440-1746.2005.04087.x, PMID:16246193.
- [112] Tseng TC, Liu CJ, Hsu CY, Hong CM, Su TH, Yang WT, *et al*. High Level of Hepatitis B Core-Related Antigen Associated With Increased Risk of Hepatocellular Carcinoma in Patients With Chronic HBV Infection of Intermediate Viral Load. *Gastroenterology* 2019;157(6):1518–1529.e3. doi:10.1053/j.gastro.2019.08.028, PMID:31470004.
- [113] Wooddell CI, Yuen MF, Chan HL, Gish RG, Locarnini SA, Chavez D, *et al*. RNAi-based treatment of chronically infected patients and chimpanzees reveals that integrated hepatitis B virus DNA is a source of HBsAg. *Sci Transl Med* 2017;9(409):eaan0241. doi:10.1126/scitranslmed.aan0241, PMID:28954926.
- [114] Song G, Yang R, Rao H, Feng B, Ma H, Jin Q, *et al*. Serum HBV core-related antigen is a good predictor for spontaneous HBeAg seroconversion in chronic hepatitis B patients. *J Med Virol* 2017;89(3):463–468. doi:10.1002/jmv.24657, PMID:27505145.
- [115] Wang ML, Deng R, Chen EQ, Tao CM, Liao J, Zhou TY, *et al*. Performance of serum HBcrAg in chronic hepatitis B patients with 8-year nucleos(t)ide analogs therapy. *Clin Res Hepatol Gastroenterol* 2019;43(3):301–309. doi:10.1016/j.clinre.2018.10.020, PMID:30497844.
- [116] Wang B, Carey I, Bruce M, Montague S, Dusheiko G, Agarwal K. HBsAg and HBcrAg as predictors of HBeAg seroconversion in HBeAg-positive patients treated with nucleos(t)ide analogues. *J Viral Hepat* 2018;25(8):886–893. doi:10.1111/jvh.12889, PMID:29532589.
- [117] Matsumoto A, Yatsushashi H, Nagaoka S, Suzuki Y, Hosaka T, Tsuge M, *et al*. Factors associated with the effect of interferon- α sequential therapy in order to discontinue nucleoside/nucleotide analog treatment in patients with chronic hepatitis B. *Hepatol Res* 2015;45(12):1195–1202. doi:10.1111/hepr.12488, PMID:25594111.
- [118] Seto WK, Tanaka Y, Wong DK, Lai CL, Shinkai N, Yuen JC, *et al*. Evidence of serologic activity in chronic hepatitis B after surface antigen (HBsAg) seroclearance documented by conventional HBsAg assay. *Hepatol Int* 2012;7(1):98–105. doi:10.1007/s12072-012-9354-7, PMID:24014110.
- [119] Seto WK, Wong DK, Fung J, Huang FY, Liu KS, Lai CL, *et al*. Linearized hepatitis B surface antigen and hepatitis B core-related antigen in the natural history of chronic hepatitis B. *Clin Microbiol Infect* 2014;20(11):1173–1180. doi:10.1111/1469-0691.12739, PMID:24975365.
- [120] Martinot-Peignoux M, Lalalus M, Maylin S, Boyer N, Castelnau C, Giuli N, *et al*. Baseline HBsAg and HBcrAg titres allow peginterferon-based ‘precision medicine’ in HBeAg-negative chronic hepatitis B patients. *J Viral Hepat* 2016;23(11):905–911. doi:10.1111/jvh.12565, PMID:27375231.
- [121] Sonneveld MJ, Park JY, Kaewdech A, Seto WK, Tanaka Y, Carey I, *et al*. Prediction of Sustained Response After Nucleos(t)ide Analogue Cessation Using HBsAg and HBcrAg Levels: A Multicenter Study (CREATE). *Clin Gastroenterol Hepatol* 2022;20(4):e784–e793. doi:10.1016/j.cgh.2020.12.005, PMID:33309804.
- [122] Seto WK, Wong DK, Chan TS, Hwang YY, Fung J, Liu KS, *et al*. Association of Hepatitis B Core-Related Antigen With Hepatitis B Virus Reactivation in Occult Viral Carriers Undergoing High-Risk Immunosuppressive Therapy. *Am J Gastroenterol* 2016;111(12):1788–1795. doi:10.1038/ajg.2016.436, PMID:27644733.
- [123] Matsumoto A, Tanaka E, Minami M, Okanoue T, Yatsushashi H, Nagaoka S, *et al*. Low serum level of hepatitis B core-related antigen indicates unlikely reactivation of hepatitis after cessation of lamivudine therapy. *Hepatol Res* 2007;37(8):661–666. doi:10.1111/j.1872-034X.2007.00094.x, PMID:17584261.
- [124] Hsu YC, Nguyen MH, Mo LR, Wu MS, Yang TH, Chen CC, *et al*. Combining hepatitis B core-related and surface antigens at end of nucleos(t)ide analogue treatment to predict off-therapy relapse risk. *Aliment Pharmacol Ther* 2019;49(1):107–115. doi:10.1111/apt.15058, PMID:30450681.
- [125] Jung KS, Park JY, Chon YE, Kim HS, Kang W, Kim BK, *et al*. Clinical outcomes and predictors for relapse after cessation of oral antiviral treatment in chronic hepatitis B patients. *J Gastroenterol* 2016;51(8):830–839. doi:10.1007/s00535-015-1153-1, PMID:26687058.
- [126] Matsumoto A, Tanaka E, Suzuki Y, Kobayashi M, Tanaka Y, Shinkai N, *et al*. Combination of hepatitis B viral antigens and DNA for prediction of re-

- lapse after discontinuation of nucleos(t)ide analogs in patients with chronic hepatitis B. *Hepatol Res* 2012;42(2):139–149. doi:10.1111/j.1872-034X.2011.00910.x, PMID:22103237.
- [127] Shinkai N, Tanaka Y, Orito E, Ito K, Ohno T, Hirashima N, *et al*. Measurement of hepatitis B virus core-related antigen as predicting factor for relapse after cessation of lamivudine therapy for chronic hepatitis B virus infection. *Hepatol Res* 2006;36(4):272–276. doi:10.1016/j.hepres.2006.08.005, PMID:16971173.
- [128] Drafting Committee for Hepatitis Management Guidelines and the Japan Society of Hepatology. JSH Guidelines for the Management of Hepatitis B Virus Infection. *Hepatol Res* 2014;44(Suppl S1):1–58. doi:10.1111/hepr.12269, PMID:24397839.
- [129] Zhang ZQ, Lu W, Wang YB, Weng QC, Zhang ZY, Yang ZQ, *et al*. Measurement of the hepatitis B core-related antigen is valuable for predicting the pathological status of liver tissues in chronic hepatitis B patients. *J Virol Methods* 2016;235:92–98. doi:10.1016/j.jviromet.2016.05.016, PMID:27230224.
- [130] Chang XJ, Sun C, Chen Y, Li XD, Yu ZJ, Dong Z, *et al*. On-treatment monitoring of liver fibrosis with serum hepatitis B core-related antigen in chronic hepatitis B. *World J Gastroenterol* 2019;25(32):4764–4778. doi:10.3748/wjg.v25.i32.4764, PMID:31528100.
- [131] Tsochatzis EA, Bosch J, Burroughs AK. Liver cirrhosis. *Lancet* 2014;383(9930):1749–1761. doi:10.1016/S0140-6736(14)60121-5, PMID:24480518.
- [132] Tada T, Kumada T, Toyoda H, Kobayashi N, Akita T, Tanaka J. Hepatitis B virus core-related antigen levels predict progression to liver cirrhosis in hepatitis B carriers. *J Gastroenterol Hepatol* 2018;33(4):918–925. doi:10.1111/jgh.13989, PMID:28914957.
- [133] Tseng TC, Liu CJ, Yang WT, Hsu CY, Hong CM, Su TH, *et al*. Serum hepatitis B core-related antigen level stratifies risk of disease progression in chronic hepatitis B patients with intermediate viral load. *Aliment Pharmacol Ther* 2021;53(8):908–918. doi:10.1111/apt.16266, PMID:33465271.
- [134] Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018;68(6):394–424. doi:10.3322/caac.21492, PMID:30207593.
- [135] Brown ZJ, Tsilimigras DI, Ruff SM, Mohseni A, Kamel IR, Cloyd JM, *et al*. Management of Hepatocellular Carcinoma: A Review. *JAMA Surg* 2023;158(4):410–420. doi:10.1001/jamasurg.2022.7989, PMID:36790767.
- [136] Craig AJ, von Felden J, Garcia-Lezana T, Sarcognato S, Villanueva A. Tumour evolution in hepatocellular carcinoma. *Nat Rev Gastroenterol Hepatol* 2020;17(3):139–152. doi:10.1038/s41575-019-0229-4, PMID:31792430.
- [137] Kumada T, Toyoda H, Tada T, Kiriya S, Tanikawa M, Hisanaga Y, *et al*. Effect of nucleos(t)ide analogue therapy on hepatocarcinogenesis in chronic hepatitis B patients: a propensity score analysis. *J Hepatol* 2013;58(3):427–433. doi:10.1016/j.jhep.2012.10.025, PMID:23123221.
- [138] Tada T, Kumada T, Toyoda H, Kiriya S, Tanikawa M, Hisanaga Y, *et al*. HBcrAg predicts hepatocellular carcinoma development: An analysis using time-dependent receiver operating characteristics. *J Hepatol* 2016;65(1):48–56. doi:10.1016/j.jhep.2016.03.013, PMID:27034253.
- [139] Cheung KS, Seto WK, Wong DK, Lai CL, Yuen MF. Relationship between HBsAg, HBcrAg and hepatocellular carcinoma in patients with undetectable HBV DNA under nucleos(t)ide therapy. *J Viral Hepat* 2017;24(8):654–661. doi:10.1111/jvh.12688, PMID:28185363.
- [140] Hosaka T, Suzuki F, Kobayashi M, Fujiyama S, Kawamura Y, Sezaki H, *et al*. Impact of hepatitis B core-related antigen on the incidence of hepatocellular carcinoma in patients treated with nucleos(t)ide analogues. *Aliment Pharmacol Ther* 2019;49(4):457–471. doi:10.1111/apt.15108, PMID:30663078.