



## Review Article

# Potential Applications of Cannabis Plant Extracts and Phytochemicals as Natural Antimicrobials



Hebah M.S. AL Ubeed<sup>1\*</sup>, Asgar Farahnaky<sup>1</sup>, Emma L. Beckett<sup>2,3</sup>, Momena Khandaker<sup>4</sup> and Christopher J. Pillidge<sup>1</sup>

<sup>1</sup>School of Science, STEM College, RMIT University, VIC, Australia; <sup>2</sup>School of Environmental and Life Sciences, University of Newcastle, NSW, Australia; <sup>3</sup>Hunter Medical Research Institute, New Lambton Heights, NSW, Australia; <sup>4</sup>School of Environmental and Rural Sciences, University of New England, NSW, Australia

Received: July 27, 2022 | Revised: October 09, 2022 | Accepted: October 24, 2022 | Published online: December 05, 2022

## Abstract

*Cannabis* has a long history of use in treating human diseases, but scientific research on its properties has only recently gained momentum. The increasing prevalence of antibiotic resistance in human and animal pathogens has sparked renewed interest in exploring alternative antimicrobial therapies from *Cannabis* and other plant sources. There is also potential for *Cannabis* extracts or purified cannabinoids to be applied in novel medical contexts. Industrial hemp extracts may find applications in food manufacturing, veterinary purposes, and microbial control in cleaners and sanitizers. This review highlights the latest discoveries regarding *Cannabis* plant extracts and phytochemicals as potent antimicrobial agents against various microorganisms, including Gram-positive and Gram-negative bacteria. More importantly, the challenges of using cannabinoids as effective and affordable natural antimicrobial agents are reviewed. While antimicrobial and other applications of *Cannabis* extracts and phytochemicals

**Keywords:** *Cannabis*; Antibacterial; Antifungal; Antiviral; Natural phytochemicals; Medicinal Cannabis extract.

**Abbreviations:**  $\Delta^8$ -THC, Delta-8-Tetrahydrocannabinol; ABTS, 2,2'-Azinobis [3-ethylbenzothiazoline-6-sulfonic acid]-diammonium salt; ACE-2, Angiotensin-converting enzyme-2; AgNP<sub>s</sub>, Silver nanoparticles; *B. cereus*, *Bacillus cereus*; *B. longum*, *Bifidobacterium longum*; *B. subtilis*, *Bacillus subtilis*; CBC, Cannabichromene; CBCA, Cannabichromene acid; CBD, Cannabidiol; CBDA, Cannabidiol acid; CBDV, Cannabidivarin; CBDVA, Cannabidivarin acid; CBG, Cannabigerol; CBGA, Cannabigerolic acid; CBL, Cannabicyclol; CBN, Cannabinol; CBNA, Cannabinol acid; CLSI, Clinical Laboratory and standards Institute; COVID-19, Human coronavirus; SARS-CoV2, severe acute respiratory syndrome coronavirus 2; DPPH, 2,2-diphenyl-1-picrylhydrazyl; *E. coli*, *Escherichia coli*; *E. faecalis*, *Enterococcus faecalis*; EOs, Essential Oils; GC-MS, Gas Chromatography Mass Spectrometry; GI, Gastrointestinal; GOT, Geranyl pyrophosphate, Olivetolate geranyltransferase; GPP, Geranyl pyrophosphate; HBV, Hepatitis B virus; HCV, Hepatitis C virus; *K. pneumoniae*, *Klebsiella pneumoniae*; KSHV, Kaposi's sarcoma-associated herpesvirus; *L. donovani*, *Leishmania donovani*; *M. smegmatis*, *Mycobacterium smegmatis*; MBC, minimum bactericidal concentration; MIC, minimum inhibitory concentration; MRSA, Methicillin-resistant *Staphylococcus aureus*; NCs, Nanocarriers; OLA, olivetol acid; *P. aeruginosa*, *Pseudomonas aeruginosa*; *P. falciparum*, *Plasmodium falciparum*; *P. guajava*, *Psidium guajava*; *S. aureus*, *Staphylococcus aureus*; *S. milleri*, *Streptococcus milleri*; *S. mutans*, *Streptococcus mutans*; *S. pyogenes*, *Streptococcus pyogenes*; SFE-CO<sub>2</sub>, Supercritical fluid extraction with carbon dioxide; THC, Tetrahydrocannabinol; THCA, Tetrahydrocannabinolic acid; THCAA, Delta9-tetrahydrocannabinolic acid-A; THCAS, Tetrahydrocannabinolic acid synthase; THCV, Tetrahydrocannabivarin; THCVA, Tetrahydrocannabivarin acid; TMRSS2, Transmembrane serine protease 2; ZnNP<sub>s</sub>, Zinc Nanoparticles;  $\Delta^9$ -THC, Delta-9-trans-Tetrahydrocannabinol.

\*Correspondence to: Hebah M.S. AL Ubeed, School of Science, STEM College, RMIT University, Bundoora, Melbourne, VIC 3083, Australia. ORCID: <https://orcid.org/0000-0003-4344-9513>. Tel: +61 3 9925 3347, Fax: RITM0806372, E-mail: hebah.al.ubeed@rmit.edu.au

**How to cite this article:** AL Ubeed HMS, Farahnaky A, Beckett EL, Khandaker M, Pillidge CJ. Potential Applications of Cannabis Plant Extracts and Phytochemicals as Natural Antimicrobials. *J Explor Res Pharmacol* 2023;8(2):121–130. doi: 10.14218/JERP.2022.00062.

appear promising, concerns about possible toxic side effects exist. Therefore, future research should focus on addressing the safety of these compounds, evaluating their *in vivo* activity, and understanding structural changes that influence their pharmacokinetic properties. Standardized tests will be crucial for facilitating valid inter-laboratory comparisons. The review article also discusses future research directions aimed at developing novel broad-spectrum antibiotics based on *Cannabis*.

## Introduction

*Cannabis sativa* L. or marijuana, is a flowering plant that has been used for millennia for food, drugs (both legal and illegal), textile (hemp), and religious purposes.<sup>1</sup> Selective breeding has resulted in numerous *C. sativa* strains (cultivars) with different properties. For example, hemp strains are fibrous and low in cannabinoids, while medicinal strains are highly flowering and contain both phytonutrients and phytochemicals.<sup>2–4</sup> Bioactive compounds can be extracted from oils or as aqueous phases from seeds, flowers, leaves, and stems using traditional techniques such as cold-pressing and solvent extraction or by contemporary procedures like ultrasound.<sup>5</sup> Supercritical fluid extraction with carbon dioxide (SFE-CO<sub>2</sub>) is another technology used in industry to extract phytochemical compounds.<sup>6,7</sup> This is no different from other plant products, which have been historically used as rich sources of natural products for human health.<sup>8</sup>

Potential therapeutic applications and bioactive mechanisms

of crude *Cannabis* extracts and purified compounds derived from *C. sativa* have been investigated in various pharmacological scenarios, including their use as anti-convulsive, analgesic, anti-anxiety, and anti-emetic therapeutic drugs.<sup>9</sup> While much research has centered around the psychoactive properties of cannabinoids,<sup>10</sup> the antimicrobial properties of compounds extracted from *C. sativa* are now becoming of particular interest due to the emergence of antimicrobial resistance as a vital threat to human health globally.<sup>11</sup> Mitigating the human and economic impacts of this problem, and more broadly, the emergence of new microbial pathogens,<sup>12</sup> requires identifying and elucidating new antimicrobial therapies.

One possibility to combat such infections, apart from antibiotics, is to use phage-based therapies, including lysis therapy and engineered phage enzymes.<sup>13,14</sup> Another alternative is to use plant-derived compounds or extracts with antimicrobial properties (bacterial, fungal, and viral). Among these, *Cannabis* extracts and (more specifically) cannabinoids show colossal promise, especially towards multi-drug resistant microorganisms like MRSA (methicillin-resistant *Staphylococcus aureus*), which can be very difficult to treat.<sup>15</sup> While there is much anecdotal evidence on the efficacy of *Cannabis* compounds or extracts, controlled laboratory investigations of these novel antimicrobial agents, their mode of action, efficacy, and safety, together with the development of safe and approved disease treatment therapies, is certainly needed. *In vitro* antifungal, antibacterial, antimalarial, antileishmanial, and cytotoxic properties of *C. sativa* extracts and compounds have been investigated since the early 2000s.<sup>13,16–19</sup> However, significant challenges exist in assessing the efficacy of these agents (both impure phytochemical extracts and purified compounds), determining their safety, and implementing research findings. These challenges include high variability in cannabinoid content in different *C. sativa* strains; also for different parts of plants and in the extraction methods;<sup>5</sup> a lack of uniform methods for antimicrobial screening assays; diversity in microbial targets (strain variability); failure to consider the stability of extracts or the effects of additional compounds (impurities), both in cell culture experiments and animal trials. Furthermore, challenges regarding the negative perception of *Cannabis*, its safety and potential side effects, variable delivery methods, local and systemic effects, and synergistic effects need to be addressed.<sup>20,21</sup>

This review will describe the potential uses of agents found in medicinal *Cannabis* extracts as antimicrobials, the current knowledge on their mechanism of action, the challenges around phytochemicals' stability and bioavailability, optimized extraction protocols, and isolation of active antibiotic compounds. Also, the prospect of using new formulations containing 'old' antibiotics combined with therapeutic plant compounds to provide a synergistic killing effect is noted. Such combined treatments could more effectively kill or inhibit antibiotic-resistant bacteria causing local and systemic infections.

### New classes of potent antimicrobial agents

It has been known for decades that *Cannabis* plant extracts can be effective antimicrobial agents.<sup>22,23</sup> More than 525 phytochemicals have been extracted and isolated in *C. sativa*.<sup>24–26</sup> The most important classes of *Cannabis* phytochemicals are the C21 terpenes, phenolics, and cannabinoids. The respective antimicrobial properties of these compounds will be discussed in this report.

There are many benefits for isolated cannabinoids, including antifungal, antibacterial, antimalarial, antileishmanial, and cytotoxic

properties.<sup>26</sup> Active cannabinoids like CBG,<sup>27,28</sup> CBN, CBC,<sup>29</sup> and psychoactive cannabinoids delta-9-trans-tetrahydrocannabinol ( $\Delta^9$ -THC)<sup>17</sup> (summarized in Table 1 and Fig. 1) and their precursors have a high level of antimicrobial activity.<sup>16,26,30,31</sup> Synthesis of cannabinoids is complex; essentially, they are derived from cannabigerolic acid (CBGA), which is the precursor of olivetol acid (OLA) and geranyl pyrophosphate (GPP), and is induced by the prenyltransferase geranyl pyrophosphate: olivetolategeranyltransferase (GOT).<sup>32</sup> There are some examples of co-enzymes involvement like tetrahydrocannabinolic acid synthase (THCAS), cannabidiolic acid synthase (CBDAS), or cannabichromeneacid synthase (CBCAS). With the assistance of the co-enzyme, CBGA is finally converted to tetrahydrocannabinolic acid (THCA), cannabidiolic acid (CBDA), and cannabichromenic acid (CBCA). Then, oxidation of THCA occurs to produce cannabinolic acid (CBNA) in the buds.<sup>33,34</sup> Cannabinoids  $\Delta^9$ -THC, CBD, CBG, CBC, and CBN displayed anti-staphylococcal activity with even greater antibiotic effect than traditional antibiotics like norfloxacin, erythromycin, tetracycline, and oxacillin.<sup>3</sup> CBG exhibits antibacterial activities against *Streptococcus mutans* (*S. mutans*).<sup>35,36</sup>

Both compounds  $\Delta^9$ -THC and CBD showed significant antibacterial activity towards *S. aureus*, *Streptococcus pyogenes* (*S. pyogenes*), *Streptococcus milleri* (*S. milleri*), *Enterococcus faecalis* (*E. faecalis*), *E. coli*, *Salmonella typhi*, and *Proteus vulgaris*.<sup>37</sup> Farha, *et al.*<sup>37</sup> further described the antibacterial activity of specific cannabinoids CBC, CBCA, CBD, CBDV, CBDVA, CBG, CBGA, CBN, CBL, THC,  $\Delta^8$ -THC, exo-tetrahydrocannabinol (exo-THC),  $\Delta^9$ -tetrahydrocannabinolic acid-A (THCAA), THCV, ( $\pm$ ) 11-nor-9-carboxy- $\Delta^9$  – THC, and ( $\pm$ ) 11-hydroxy- $\Delta^9$ -THC against MRSA to inhibit its ability to form biofilms and stationary phase cells' resistance to antibiotics. CBD also showed potent activity against various MRSA strains.<sup>38</sup> A detailed paper on the antimicrobial activity of CBD, including mechanistic mode-of-action studies, was later provided by Blaskovich *et al.*<sup>17</sup> They showed that cannabidiol has a superior effect against biofilms. This cannabinoid could potentially treat Gram-negative bacteria infections, including *Neisseria gonorrhoeae* and Gram-positive bacterial infections. CBD was also shown to be a bigger inhibitor of membrane vesicle emission from *E. coli* (strain VCS257) compared to *S. aureus*. Whenever CBD was used in combination with selected antibiotics, it caused higher inhibition action against Gram-negative bacteria.<sup>39</sup> Similar to terpenes, cannabinoids show synergistic effects with antibiotics. Grassi *et al.*<sup>40</sup> demonstrated that when cannabinoids were combined with polymyxin B, there was an effective inhibition against Gram-negative bacteria.


Apart from cannabinoids, geranyl pyrophosphate is the precursor in synthesizing the terpenoids, leading to the creation of monoterpenoids in secretory cell plastids.<sup>41</sup> For instance, volatile oil fractions incorporate monoterpenoids (C10)<sup>42</sup> or sesquiterpenoids and triterpenoids in the cytoplasm.<sup>43,44</sup> Sesquiterpenes (C15) are major compounds in hemp extracts.<sup>42</sup> After harvest, *Cannabis* buds must be dried to remove the carboxylic acid functional group to extract high-purity CBD, CBC, and CBG. Then, oxidation of THC yields delta-8-tetrahydrocannabinol ( $\Delta^8$ -THC, the main psychoactive compound) and CBN.<sup>26,45,46</sup>

Many factors affect the quality and consistency of these antibacterial extracts, including environmental and climatic conditions for the growth of leaves, flowers, and seeds. Also, Muscarà *et al.*<sup>47</sup> studied two standardized hexane extracts for their phytonutrients and antibiotic activities. The first extract consisted of cannabidivarinic acid (CBDVA) and tetrahydrocannabivarinic acid (THCVA). The second extract contained cannabidivarin (CBDV)

**Table 1. Cannabinoids from *Cannabis sativa* L. with antimicrobial activity**

Active compound	Organism	Mode of action	References
CBD	<i>S. aureus</i> ; <i>B. subtilis</i>	CBD is more active than CBDA due to exchanged positions of the one hydroxyl group and lipophilic side chain.	3,89,100
	Hepatitis C virus	CBD interacts with the CB <sub>2</sub> receptor and stimulates apoptosis in thymocytes and splenocytes, then inhibiting the proliferation of T-cells and macrophages as such mechanism cause to indirectly slows the pathogenic process of the HBV virus.	
	SARS-CoV-2	CBD additive or synergic effect with terpene control viral replication or clonal stable conformations with the binding of the transmembrane protease serine 2 (SARS-CoV-2) and angiotensin-converting enzyme-2 (ACE2).	
CBG	<i>Mycobacterium donovani</i> ; <i>S. mutans</i> ; <i>S. sanguis</i> ; <i>S. sobrinus</i> ; <i>S. salivarius</i> ; MRSA	CBG alters membrane structures of treated bacteria and disorders of the cytoplasm activity.	70
CBC	<i>S. aureus</i> ; <i>B. Subtilis</i> ; <i>C. albicans</i> ; <i>Mycobacterium smegmatis</i> ; <i>Saccharomyces cerevisiae</i> ; <i>Trichophyton mentagrophytes</i>	The activity of CBC due to bearing the lipophilic side chain and one hydroxyl group in exchanged positions displayed similarly potent activity.	3,74
CBN	Anti-MRSA; antileishmanial; <i>Plasmodium falciparum</i>	Decreased or increased esterification or methylation of the carboxylic acid moieties were detrimental to the activity and additional hydroxy function in the lipophilic side chain.	3,76
$\Delta^9$ -THC	<i>S. aureus</i> , <i>Streptococcus pyogenes</i> , <i>Streptococcus. milleri</i> , <i>Enterococcus faecalis</i> , <i>E. coli</i> , <i>Salmonella typhi</i> , <i>Proteus vulgaris</i> , MRSA	Decreased or increased esterification or methylation of the carboxylic acid moieties were detrimental to the activity.	3,16

ACE-2, Angiotensin-converting enzyme-2; CBD, Cannabidiol; CBDA, Cannabidiol acid; CBG, Cannabigerol; CBN, Cannabinol; MRSA, Methicillin-resistant *Staphylococcus aureus*; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2;  $\Delta^9$ -THC, Delta-9-trans-Tetrahydrocannabinol.



**Fig. 1. Antimicrobial activity of cannabinoids like cannabidiol (CBD), cannabigerol (CBG), cannabichromene (CBC), cannabinol (CBN), and delta-8-tetrahydrocannabinol ( $\Delta^8$ -THC).**

and tetrahydrocannabivarin (THCV) from a new Chinese *C. sativa* variety and other non-psychoactive strains. Both extracts showed extraordinary antioxidant activity and antimicrobial and antifungal properties against clinical strains of MRSA and other microorganisms.<sup>28</sup> Various methods have been applied to extract and isolate these and other antimicrobial agents, such as solvent extraction, physical methods, and supercritical fluid extraction, giving different results.<sup>5,6</sup>

Other *Cannabis* compounds with antimicrobial activity include terpenoids. These form a large percentage of essential oils.<sup>48-51</sup> Some 120 terpenoids have been found in marijuana and hemp,<sup>52</sup> but relatively few terpenes have been isolated, purified, and tested.<sup>53</sup> Hemp seed oil was also effective in controlling spoilage of food and phytopathogenic microorganisms.<sup>16,54</sup> An *in vitro* or cell culture study by Nostro *et al.*<sup>55</sup> considered a range of plant extracts, potentially inhibiting biofilms in food. They found that extracts were effective against *S. aureus*, including MRSA strains, *Escherichia coli* (*E. coli*), and *Pseudomonas aeruginosa* (*P. aeruginosa*). Another study focused on linalool and  $\alpha$ -Phellandrene, which were investigated due to their very low toxicity and ease of use and were registered as a potential new therapeutic.<sup>47,56</sup> Another type of active compound extracted from *Cannabis* seeds are the flavonoids, also effective against bacteria and yeast.<sup>57</sup>

Another active compound class that can be extracted from *Cannabis* and used as an antibiotic alone or in combination with other compounds is *Cannabis* alkaloids. Natural alkaloids accumulate to differing extents in particular parts of the plant, such as barks, roots, and leaves.<sup>58,59</sup> These alkaloids can be chemically modified to produce synthetic and semi-synthetic smart drugs. However, natural compounds are generally safer and cheaper than alternative synthetic compounds.<sup>60</sup> Currently, natural alkaloids are used as a therapeutic compound for human disease and as pesticides in agriculture.<sup>61</sup> Natural alkaloids and phenolic compounds have potential antimicrobial activity due to their ability to alter the structure of the bacterial wall.<sup>62</sup>

In summary, the most valuable compounds in *Cannabis*, besides cannabinoids, include its alkaloids, terpenoids, steroids, phenols, glycosides, and tannins.<sup>63</sup>

### A novel mode of action

The most active compounds in *C. sativa* with antimicrobial potential are THC, CBD, CBG, and CBC.<sup>56,64-66</sup> These compounds may also effectively treat psychological and physiological disorders.<sup>67,68</sup> Medicinal *Cannabis* phytochemicals applied to specific medical devices were active against biofilms composed of Gram-positive and Gram-negative bacteria, and their mode of action was to alter and penetrate the bacteria cells<sup>69</sup> are summarized in Table 1. Various methods like scanning electron microscopy, transmission electron microscopy, Nile Red membrane staining, and laurdan membrane fluidity assays have shown that CBG inhibits bacteria via alterations to the bacterial cell structure by inducing membrane hyperpolarization and decreasing the membrane fluidity.<sup>70</sup> CBG exhibited moderate antimicrobial activity towards *Mycobacterium*<sup>25</sup> and *Leishmania donovani* (*L. donovani*).<sup>3</sup> Not only CBG but also its derivative iso-CBG-C1 showed potential activity against *S. aureus* and *Bacillus subtilis* (*B. subtilis*), and *Mycobacterium smegmatis* (*M. smegmatis*).<sup>27</sup>

Other studies found that CBD has a higher effect against Gram-positive bacteria than their acid form due to differences in lipophilicity.<sup>27,71,72</sup> Studies have shown that CBD and CBG have a higher antimicrobial effect than their precursors (acidic cannabinoids or

raw material before drying) due to exchanged positions of a hydroxyl group and lipophilic side chain.<sup>3</sup>

While CBD and CBG are effective against Gram-positive bacteria, Krejci *et al.*<sup>73</sup> reported that both cannabinoids were inactive against Gram-negative bacteria. Moreover, compared to other natural cannabinoids and antibiotics like streptomycin, CBC has the highest activity against *S. aureus* and *B. subtilis*,<sup>74</sup> and similar effects were confirmed with iso-CBC due to connecting with the lipophilic side chain and a hydroxyl group in interchanged positions.<sup>75</sup> Furthermore, a cannabinol derivative CBN<sup>76,77</sup> showed moderate anti-MRSA, antileishmanial, and anti-*Plasmodium falciparum* (*P. falciparum*) activity. However, both  $\Delta^9$ -THCVA and CBDVA showed lower anti-staphylococcal activity based on deleting a side chain compared to  $\Delta^9$ -THCV and CBDV.<sup>78,79</sup> The primary action of CBG against MRSA is the disordering of the principal activity of the cytoplasm.<sup>37</sup> Recent structure-function studies have helped to determine the most effective cannabinoids.<sup>80</sup> Cannabinoids showed significant differences in antimicrobial activity due to a decrease or increase in methylation or esterification. The most active antimicrobial cannabinoids are  $\Delta^9$ -THC, CBN, and CBG.<sup>16,70</sup>

No significant antifungal activity has been observed yet for cannabinoids.<sup>19</sup> However, the antiviral potential of CBD has received some attention.<sup>81-86</sup> CBD was effective against hepatitis C virus (HCV),<sup>87-92</sup> which causes liver inflammation.<sup>93</sup> In contrast, CBD was ineffective against the hepatitis B virus (HBV). CBD was more efficient than sofosbuvir and had a less cytotoxic effect.<sup>94</sup> Other research indirectly showed that CBD could effectively work and inhibit Kaposi's sarcoma-associated herpesvirus (KSHV).<sup>95</sup>

One key research target is control of the human coronavirus (COVID-19).<sup>96-99</sup> In a recent article, CBD in combination with terpenes showed antiviral effects against Human Coronavirus E229.<sup>100</sup> Also, in an earlier investigation, terpenes were found to show an antiviral effect on severe acute respiratory syndrome coronavirus 1 (SARS-CoV-1).<sup>101</sup> However, while such compounds show promise, most of these antivirals have not been trialed on humans or undergone controlled clinical trials.<sup>102,103</sup> According to Chatow *et al.*,<sup>100</sup> the combined possibility of plant terpenes with CBD against a human coronavirus strain could be seen using a tissue cell culture model.

Although such studies are a good start, they cannot determine side effects or potential cytotoxicity as *in vivo* animal model studies. Recent studies showed phytochemicals have stable conformations with the binding enzyme of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), which has a vital role in viral replication, cloning, suppressing viral entry and activation, and down-regulating ACE2 receptor and TMPRSS2 enzyme.<sup>104-106</sup> Further investigation on developing new vaccines or antivirals will be needed to address strain variability, i.e., target all strains of COVID-19. This needs more intensive research in the future, together with studies giving mechanistic insights into their action.

### Challenges in applications of cannabinoids as an antimicrobial agent

As noted earlier, the spread of antibiotic resistance genes is common among many pathogens, and with little or no pending discovery or development of new classes of antibiotics, the need to find other agents for combating infections is vital.<sup>37,107,108</sup> Recently, the World Health Organization announced an archive of 12 microorganisms that have developed antibacterial resistance and are now considered a severe threat to health care. Eight of these pathogens



are Gram-negative and are spectacularly hard to cure, while the other four Gram-positive microorganisms are the most significant threat.<sup>109</sup> The antimicrobial activity of *Cannabis* derives from essential oils (Eos) and purified cannabinoids, the latter receiving increased attention in recent years as possible therapeutic agents.<sup>16,19,56,110</sup> This section will first consider the antimicrobial effects of impure extracts and Eos, then discuss studies on purified cannabinoids.

The disc diffusion assay and the two-fold serial broth dilution assay, which generate a MIC value in a 96-well microtiter plate, are the most commonly used methods to measure antimicrobial effects.<sup>111</sup> The methodology for the MIC assay is standard and specified by the Clinical Laboratory and Standards Institute (CLSI).<sup>112</sup> However, a problem highlighted by Sadgrove and Jones<sup>113</sup> is that MIC values are normally quoted in µg/ml, which does not take into account molarity. The authors also noted that antimicrobial outcomes might be naively extrapolated beyond any reasonable systemic concentration level to show an effect. Even with method standardization, caution must be exercised when comparing inter-laboratory results to ensure that results cross-comparisons are valid. In one study, even though there was the bactericidal effect of both cannabinoids like THC and CBD against staphylococci and streptococci tested in broth media, the impact was much higher in horse blood agar or with 4% horse serum.<sup>109</sup>

Even though essential oils (Eos) can be extracted from low-THC *Cannabis* varieties cultivated without legal restrictions, relatively few studies have considered these Eos or extracts in terms of their antimicrobial activities. The use of *Cannabis* in this way strengthens the idea of growing hemp as a multi-use crop.<sup>54</sup> A fundamental challenge in medical *Cannabis* and hemp research requires production methods or post-harvesting processes that maintain the uniform quality of the product. Also, drying, extraction, isolation, and purification techniques must be optimized to obtain pure chemical composition and pharma grade for different products.<sup>6,114–116</sup>

One problem with assessing multiple studies is that *Cannabis* extracts vary enormously in composition and contain multiple different compounds, making result comparisons extremely difficult. Some key variables that make inter-laboratory comparisons difficult include the absence of standardized antimicrobial testing methodologies; microbiological media components in disk diffusion tests reacting with or modifying extracts; plant extracts containing unstable compounds making accurate quantitative analysis impossible; synergistic or antagonistic effects between multiple compounds; different cultivars or parts of plants used in different studies; and extraction protocols of varying efficiency leading to different outcomes. In one study, *C. sativa* and *Psidium guajava* extracts contained various active compounds like alkaloids, saponins, flavonoids, steroids, cardiac glycosides, terpenes, resins, tannins, and phenols. However, steroids, resins, and cardiac glycosides were absent in another medicinal plant, *Thuja orientalis*. Furthermore, hemp essential oils demonstrated high antimicrobial activity against Gram-positive bacteria.<sup>20,54</sup>

Some studies have focused on extracts of seeds of *Cannabis* plants.<sup>117</sup> Hemp seed extracts show good antimicrobial activity due to their composition of polyphenols, essentially caffeine, tyramine and cannabisin.<sup>118</sup> In another study,<sup>117</sup> *C. sativa* L. seed extracts inhibited biofilm formation by *S. aureus* ATCC 35556. In contrast, adding hemp seed extract to media stimulated the growth of *Bifidobacterium* and *Lactobacillus* probiotic strains.<sup>117</sup> The increased growth of *Bifidobacterium Longum* (*B. longum*) in the presence of hemp extract was due to protection from oxidative

stress.<sup>119</sup>

Most studies on the effects of extracts use classical disc diffusion methods. However, recently, the study reported by Frassinetti *et al.*<sup>117</sup> was to determine the lowest concentrations with activity via MIC and a biofilm production and inhibition assay. A further study by Iseppi *et al.*<sup>120</sup> aimed to standardize the antibacterial method and provided better designs for experiments. Firstly, the separation of essential oils (EO) compounds was done by gas chromatography mass spectrometry (GC-MS). The EOs were tested via an agar well-diffusion assay, and MIC testing was done against multiple Gram-positive bacteria. The EOs and purified compounds were highly effective against *Enterococcus*, an opportunistic pathogen. These authors also found that two EOs and two purified compounds had lower MICs than amoxicillin or ampicillin against *Bacillus cereus* (*B. cereus*).

Another active compound extracted from the seed was flavonoids; these were shown to be active against *Candida albicans*, *S. aureus*, and *P. aeruginosa*.<sup>57</sup> EOs of fiber-type hemp have also been applied to prevent food spoilage, incorporating antimicrobial extracts into food packaging. Further studies are needed on pure compounds.<sup>28</sup>

An exciting use of *Cannabis* extracts has been forming silver or zinc nanoparticles (Ag NPs and Zn NPs) with antimicrobial activity. Chouhan and Guleria,<sup>18</sup> Chauhan *et al.*<sup>121</sup> used *C. sativa* extracts to form stable nanoparticle emulsions with Ag-doped, Zn, and ZnO nanoparticles; AgNPs revealed an excellent antioxidant capacity and significant antibiotic activity against several human infection diseases by disc diffusion test, including *E. coli*, *Klebsiella pneumoniae* (*K. pneumoniae*), MRSA, *P. aeruginosa*, *S. typhi*, and *S. aureus*, and as an antifungal against *Fusarium* spp. *Rosellinia necatrix*. *C. sativa* aqueous leaf extract (CSE) derived AgNPs also had antifungal and α-amylase inhibitory activity. There was minor activity against *Bacillus subtilis*, *S. aureus*, and *K. pneumoniae* tested by a well-diffusion assay with increasing concentrations of AgNPs. Other researchers<sup>122</sup> suggested a synergistic or symbiotic effect between terpenes, flavonoids, and cannabinoids in industry hemp strains to improve the effect of silver nanoparticles (AgNPs). Other studies showed the effective use of hemp extracts to produce nanoparticles effective against biofilms.<sup>123</sup> The main challenge with using nanoparticles based on *C. sativa* extracts is to develop suitable technologies for obtaining nanoparticles with specific properties used in pharmaceutical products. For recent reviews see.<sup>124,125</sup>

Synergistic effects between *Cannabis sativa* and plants such as *Allium sativum* (Garlic) have been shown.<sup>126</sup> Further research is required to study the impact of terpenes, in particular, as antimicrobial agents.<sup>127</sup> There may also be a synergistic effect when these compounds are applied with antibiotics simultaneously, as demonstrated for other plant extracts by Blesson *et al.*<sup>128</sup> In comparative testing, the presence of other compounds must be considered.

Several reviews have summarized the antimicrobial properties of the major cannabinoids against essential pathogenic microorganisms or viruses, including Gram-negative pathogens, MRSA, and SARS-CoV-2.<sup>3,19,22,129,130</sup> From this extensive compilation of data, there is a need for standardization of methodology and approaches so that valid comparisons and conclusions between laboratories can be made. This includes how *in vitro* results can be extrapolated to determine *in vivo* efficacy, determined (usually) by using animal models.<sup>37</sup> However, further *in vivo* or animal model investigation is required to determine how the body reacts and responds to cannabinoids, including side effects.

Another challenge for using cannabinoids in terms of antimi-

crobal activity is to control their stability and bioavailability once introduced into the body. Cannabinoids have significant pharmacological activities but may show poor water solubility and become labile during processing and storage.<sup>131</sup> A commonly used method for the delivery of cannabinoids and improvements in bioavailability is using nanocarriers (NCs) to protect core materials from degradation during passage through the gastrointestinal (GI) tract.<sup>132</sup> Transmucosal oral routes of delivery offer distinct advantages.<sup>5</sup> Determining dosage forms for any medicinal product must consider product instability, such as at room temperature. In this regard, CBD is highly unstable and sensitive to oxidation.<sup>133</sup>

### Future directions

For the potential of *Cannabis* plant extracts and phytochemicals to be applied as natural antimicrobials, further work is required to standardize and harmonize research. As is genuine for all-natural extracts, variability in natural bioactive contents between plants, parts of plants, and extracts needs to be addressed with assessment and clear transparent reporting of properties. Additional research to identify active compounds (in isolation and synergistic applications) will potentially circumvent these issues by refining test substances. However, challenges persist regarding stability, delivery, safety, non-uniform testing procedures, and negative perceptions of these products. A multi-disciplinary approach is required to facilitate the progress from promising potential antimicrobials to pharmacological success, with roles for biological sciences, pharmacology, chemistry, and social and other sciences.

### Conclusions

The evolution of antibiotic-resistant bacteria and the emergence of new viruses such as COVID-19 have stimulated the search for non-traditional antimicrobial or antiviral treatments. In the case of bacteria, this includes alternatives to using antibiotics. While the use of *Cannabis* as a medicinal agent has been known for thousands of years, there is now great interest in discovering and analyzing potent antimicrobial agents derived from this plant, including its cannabinoids, terpenes, phenolics, and alkaloids. While this is welcome, inter-laboratory comparisons of results showing antimicrobial or antiviral activities are complex, as many variables affect results (such as plant-to-plant variations, differences in extraction protocols, synergistic/antagonistic effects of other compounds, and differences in antimicrobial assay protocols). There are also challenges for commercial applications, such as finding the best method for extraction and isolation of the active compound, not to mention understanding the stability and bioavailability of the cannabinoids and their pharmacokinetics once introduced into the body. Also, more study is necessary to determine the synergic effect of natural therapeutic plants in combination with *Cannabis*, to design functional nanoparticle delivery devices, and to test the safety and efficacy of an experimental new antibiotic in clinical trials.

### Acknowledgments

None.

### Funding

HMSAU and AF were funded by MGC Pharmaceuticals Limited, Australia.

### Conflict of interest

Funding received from a commercial company (MGC Pharmaceuticals Limited) is a potential conflict.

### Author contributions

Conceptualization and design (HMSAU); writing original draft preparation, Abstract, New classes of potent antimicrobial agents, A novel mode of action, Conclusion (HMSAU); Introduction and future research directions (ELB); A novel mode of action (MK); Abstract, Introduction, Challenges on applications of Cannabinoids as an antimicrobial agent (CP); Figure design (HMSAU); writing a review, and editing (AF, CP, ELB, HMSAU). All authors have read and agreed to the published version of the manuscript.

### References

- [1] Bonini SA, Premoli M, Tambaro S, Kumar A, Maccarinelli G, Memo M, *et al.* Cannabis sativa: A comprehensive ethnopharmacological review of a medicinal plant with a long history. *J Ethnopharmacol* 2018;227:300–315. doi:10.1016/j.jep.2018.09.004, PMID:30205181.
- [2] Fishedick JT, Hazekamp A, Erkelens T, Choi YH, Verpoorte R. Metabolic fingerprinting of Cannabis sativa L., cannabinoids and terpenoids for chemotaxonomic and drug standardization purposes. *Phytochemistry* 2010;71(17-18):2058–2073. doi:10.1016/j.phytochem.2010.10.001, PMID:21040939.
- [3] Klahn P. Cannabinoids-Promising Antimicrobial Drugs or Intoxicants with Benefits? *Antibiotics (Basel)* 2020;9(6):E297. doi:10.3390/antibiotics9060297, PMID:32498408.
- [4] Simiyu DC, Jang JH, Lee OR. Understanding *Cannabis sativa* L.: Current Status of Propagation, Use, Legalization, and Haploid-Inducer-Mediated Genetic Engineering. *Plants (Basel)* 2022;11(9):1236. doi:10.3390/plants11091236, PMID:35567237.
- [5] Fathordobady F, Singh A, Kitts DD, Pratap Singh A. Hemp (*Cannabis Sativa* L.) extract: anti-microbial properties, methods of extraction, and potential oral delivery. *Food Rev Int* 2019;35(7):664–684. doi:10.1080/87559129.2019.1600539.
- [6] Al Ubeed HMS, Bhuyan DJ, Alsherbiny MA, Basu A, Vuong QV. A Comprehensive Review on the Techniques for Extraction of Bioactive Compounds from Medicinal Cannabis. *Molecules* 2022;27(3):604. doi:10.3390/molecules27030604, PMID:35163863.
- [7] Rosenthal E, Zeman G. CO<sub>2</sub> extraction. Beyond buds: Next generation-marijuana extracts and cannabis infusions. *CA: Quick American Archives*; 2018:149–167.
- [8] Dias DA, Urban S, Roessner U. A historical overview of natural products in drug discovery. *Metabolites* 2012;2(2):303–336. doi:10.3390/metabo2020303, PMID:24957513.
- [9] Baker D, Pryce G, Giovannoni G, Thompson AJ. The therapeutic potential of cannabis. *Lancet Neurol* 2003;2(5):291–298. doi:10.1016/S1474-4422(03)00381-8, PMID:12849183.
- [10] Andre CM, Hausman JF, Guerriero G. Cannabis sativa: The Plant of the Thousand and One Molecules. *Front Plant Sci* 2016;7:19. doi:10.3389/fpls.2016.00019, PMID:26870049.
- [11] Ferri M, Ranucci E, Romagnoli P, Giaccone V. Antimicrobial resistance: A global emerging threat to public health systems. *Crit Rev Food Sci Nutr* 2017;57(13):2857–2876. doi:10.1080/10408398.2015.1077192, PMID:26464037.
- [12] Vouga M, Greub G. Emerging bacterial pathogens: the past and beyond. *Clin Microbiol Infect* 2016;22(1):12–21. doi:10.1016/j.cmi.2015.10.010, PMID:26493844.
- [13] Danis-Wlodarczyk KM, Wozniak DJ, Abedon ST. Treating Bacterial Infections with Bacteriophage-Based Enzybiotics: In Vitro, In Vivo and Clinical Application. *Antibiotics (Basel)* 2021;10(12):1497. doi:10.3390/antibiotics10121497, PMID:34943709.
- [14] Rios AC, Moutinho CG, Pinto FC, Del Fiol FS, Jozala A, Chaud MV, *et al.* Alternatives to overcoming bacterial resistances: State-of-the-art. *Microbiol Res* 2016;191:51–80. doi:10.1016/j.micres.2016.04.008, PMID:27524653.

- [15] David MZ, Dryden M, Gottlieb T, Tattevin P, Gould IM. Recently approved antibacterials for methicillin-resistant *Staphylococcus aureus* (MRSA) and other Gram-positive pathogens: the shock of the new. *Int J Antimicrob Agents* 2017;50(3):303–307. doi:10.1016/j.ijantimicag.2017.05.006, PMID:28666751.
- [16] Appendino G, Gibbons S, Giana A, Pagani A, Grassi G, Stavri M, *et al.* Antibacterial cannabinoids from *Cannabis sativa*: a structure-activity study. *J Nat Prod* 2008;71(8):1427–1430. doi:10.1021/np8002673, PMID:18681481.
- [17] Blaskovich MAT, Kavanagh AM, Elliott AG, Zhang B, Ramu S, Amado M, *et al.* The antimicrobial potential of cannabidiol. *Commun Biol* 2021;4(1):7. doi:10.1038/s42003-020-01530-y, PMID:33469147.
- [18] Chouhan S, Guleria S. Green synthesis of AgNPs using *Cannabis sativa* leaf extract: Characterization, antibacterial, anti-yeast and  $\alpha$ -amylase inhibitory activity. *Materials Science for Energy Technologies* 2020;3:536–544. doi:10.1016/j.mset.2020.05.004.
- [19] Karas JA, Wong LJM, Paulin OKA, Mazeh AC, Hussein MH, Li J, *et al.* The Antimicrobial Activity of Cannabinoids. *Antibiotics (Basel)* 2020;9(7):E406. doi:10.3390/antibiotics9070406, PMID:32668669.
- [20] Chakraborty S, Afaq N, Singh N, Majumdar S. Antimicrobial activity of *Cannabis sativa*, *Thuja orientalis* and *Psidium guajava* leaf extracts against methicillin-resistant *Staphylococcus aureus*. *J Integr Med* 2018;16(5):350–357. doi:10.1016/j.joim.2018.07.005, PMID:30120078.
- [21] Nashra A, Sujatha R, Sameer D, Bagoliwal A, Mishra V, Kumar A, *et al.* Comparative Evaluation of Antibacterial Efficacy of *Cannabis sativa*, *Allium sativum*, *Allium cepa*, *Thuja orientalis* and *Psidium guajava* against Drug Resistance Pathogens. *International Journal of Health Sciences and Research* 2018;8(8):89–97.
- [22] Mahmud MS, Hossain MS, Ahmed ATMF, Islam MZ, Sarker ME, Islam MR. Antimicrobial and Antiviral (SARS-CoV-2) Potential of Cannabinoids and *Cannabis sativa*: A Comprehensive Review. *Molecules* 2021;26(23):7216. doi:10.3390/molecules26237216, PMID:34885798.
- [23] Salami SA, Martinelli F, Giovino A, Bachari A, Arad N, Mantri N. It Is Our Turn to Get Cannabis High: Put Cannabinoids in Food and Health Baskets. *Molecules* 2020;25(18):E4036. doi:10.3390/molecules25184036, PMID:32899626.
- [24] Elsohly MA, Slade D. Chemical constituents of marijuana: the complex mixture of natural cannabinoids. *Life Sci* 2005;78(5):539–548. doi:10.1016/j.lfs.2005.09.011, PMID:16199061.
- [25] Radwan MM, Ross SA, Slade D, Ahmed SA, Zulficar F, Elsohly MA. Isolation and characterization of new Cannabis constituents from a high potency variety. *Planta Med* 2008;74(3):267–272. doi:10.1055/s-2008-1034311, PMID:18283614.
- [26] Radwan MM, Elsohly MA, Slade D, Ahmed SA, Wilson L, El-Alfy AT, *et al.* Non-cannabinoid constituents from a high potency Cannabis sativa variety. *Phytochemistry* 2008;69(14):2627–2633. doi:10.1016/j.phytochem.2008.07.010, PMID:18774146.
- [27] Elsohly HN, Turner CE, Clark AM, Elsohly MA. Synthesis and antimicrobial activities of certain cannabichromene and cannabigerol related compounds. *J Pharm Sci* 1982;71(12):1319–1323. doi:10.1002/jps.2600711204, PMID:7153877.
- [28] Khan BA, Warner P, Wang H. Antibacterial properties of hemp and other natural fibre plants: a review. *BioResources* 2014;9(2):3642–3659. doi:10.15376/biores.9.2.3642-3659.
- [29] Turner CE, Elsohly MA. Biological activity of cannabichromene, its homologs and isomers. *J Clin Pharmacol* 1981;21(S1):283S–291S. doi:10.1002/j.1552-4604.1981.tb02606.x, PMID:7298870.
- [30] Ahmed SA, Ross SA, Slade D, Radwan MM, Zulficar F, Matsumoto RR, *et al.* Cannabinoid ester constituents from high-potency Cannabis sativa. *J Nat Prod* 2008;71(4):536–542. doi:10.1021/np070454a, PMID:18303850.
- [31] Molnár J, Csiszár K, Nishioka I, Shoyama Y. The effects of cannabis-terpene compounds and tetrahydrocannabinolic acid on the plasmid transfer and maintenance in *Escherichia coli*. *Acta Microbiol Hung* 1986;33(3):221–231. PMID:3551476.
- [32] Fellermeier M, Zenk MH. Prenylation of olivetolate by a hemp transferase yields cannabigerolic acid, the precursor of tetrahydrocannabinol. *FEBS Lett* 1998;427(2):283–285. doi:10.1016/s0014-5793(98)00450-5, PMID:9607329.
- [33] Sirikantaramas S, Morimoto S, Shoyama Y, Ishikawa Y, Wada Y, Shoyama Y, *et al.* The gene controlling marijuana psychoactivity: molecular cloning and heterologous expression of Delta1-tetrahydrocannabinolic acid synthase from *Cannabis sativa* L. *J Biol Chem* 2004;279(38):39767–39774. doi:10.1074/jbc.M403693200, PMID:15190053.
- [34] Taura F, Morimoto S, Shoyama Y, Mechoulam R. First direct evidence for the mechanism of DELTA. 1-tetrahydrocannabinolic acid biosynthesis. *J Am Chem Soc* 1995;117(38):9766–9767. doi:10.1021/ja00143a024.
- [35] Feng J, Gu Y, Quan Y, Gao W, Dang Y, Cao M, *et al.* Construction of energy-conserving sucrose utilization pathways for improving poly- $\gamma$ -glutamic acid production in *Bacillus amyloliquefaciens*. *Microb Cell Fact* 2017;16(1):98. doi:10.1186/s12934-017-0712-y, PMID:28587617.
- [36] Feldman M, Sionov RV, Mechoulam R, Steinberg D. Anti-Biofilm Activity of Cannabidiol against *Candida albicans*. *Microorganisms* 2021;9(2):441. doi:10.3390/microorganisms9020441, PMID:33672633.
- [37] Farha MA, El-Halfawy OM, Gale RT, MacNair CR, Carfrae LA, Zhang X, *et al.* Uncovering the Hidden Antibiotic Potential of Cannabis. *ACS Infect Dis* 2020;6(3):338–346. doi:10.1021/acscinfed.9b00419, PMID:32017534.
- [38] Centers for Disease Control and Prevention (U.S.); National Center for Emerging Zoonotic and Infectious Diseases (U.S.). Division of Healthcare Quality Promotion. Antibiotic Resistance Coordination and Strategy Unit. Antibiotic resistance threats in the United States, 2019. <https://stacks.cdc.gov/view/cdc/82532>.
- [39] Kosgodage US, Mawelele P, Awamaria B, Kraev I, Warde P, Mastroianni G, *et al.* Cannabidiol Is a Novel Modulator of Bacterial Membrane Vesicles. *Front Cell Infect Microbiol* 2019;9:324. doi:10.3389/fcimb.2019.00324, PMID:31552202.
- [40] Grassi L, Maisetta G, Esin S, Batoni G. Combination Strategies to Enhance the Efficacy of Antimicrobial Peptides against Bacterial Biofilms. *Front Microbiol* 2017;8:2409. doi:10.3389/fmicb.2017.02409, PMID:29375486.
- [41] Loza-Tavera H. Monoterpenes in essential oils. Biosynthesis and properties. *Adv Exp Med Biol* 1999;464:49–62. doi:10.1007/978-1-4615-4729-7\_5, PMID:10335385.
- [42] Bertoli A, Tozzi S, Pistelli L, Angelini LG. Fibre hemp inflorescences: From crop-residues to essential oil production. *Industrial Crops Products* 2010;32(3):329–337. doi:10.1016/j.indcrop.2010.05.012.
- [43] Dewick PM. The biosynthesis of C5–C25 terpenoid compounds. *Nat Prod Rep* 2002;19(2):181–222. doi:10.1039/b002685i, PMID:12013278.
- [44] Russo EB. Taming THC: potential cannabis synergy and phytocannabinoid-terpenoid entourage effects. *Br J Pharmacol* 2011;163(7):1344–1364. doi:10.1111/j.1476-5381.2011.01238.x, PMID:21749363.
- [45] Gagne SJ, Stout JM, Liu E, Boubakir Z, Clark SM, Page JE. Identification of olivetolic acid cyclase from *Cannabis sativa* reveals a unique catalytic route to plant polyketides. *Proc Natl Acad Sci U S A* 2012;109(31):12811–12816. doi:10.1073/pnas.1200330109, PMID:22802619.
- [46] Thomas BF, Elsohly M. The analytical chemistry of cannabis: Quality assessment, assurance, and regulation of medicinal marijuana and cannabinoid preparations. Elsevier; 2015.
- [47] Muscarà C, Smeriglio A, Trombetta D, Mandalari G, La Camera E, Occhiuto C, *et al.* Antioxidant and antimicrobial activity of two standardized extracts from a new Chinese accession of non-psychoactive Cannabis sativa L. *Phytother Res* 2021;35(2):1099–1112. doi:10.1002/ptr.6891, PMID:33034400.
- [48] McPartland JM, Russo EB. Cannabis and cannabis extracts: greater than the sum of their parts? *Journal of Cannabis Therapeutics* 2001;1(3-4):103–132. doi:10.1300/J175v01n03\_08.
- [49] Nagy DU, Cianfaglione K, Maggi F, Sut S, Dall'Acqua S. Chemical Characterization of Leaves, Male and Female Flowers from Spontaneous Cannabis (*Cannabis sativa* L.) Growing in Hungary. *Chem Biodivers* 2019;16(3):e1800562. doi:10.1002/cbdv.201800562, PMID:30548994.
- [50] Novak J, Franz C. Composition of the essential oils and extracts of two populations of *Cannabis sativa* L. ssp. *spontanea* from Austria. *J Essent Oil Res* 2003;15(3):158–160. doi:10.1080/10412905.2003.9712100.
- [51] Ross SA, Elsohly MA. The volatile oil composition of fresh and air-dried buds of *Cannabis sativa*. *J Nat Prod* 1996;59(1):49–51. doi:10.1021/np960004a, PMID:8984153.



- [52] Turner CE, Elshohly MA, Boeren EG. Constituents of Cannabis sativa L. XVII. A review of the natural constituents. *J Nat Prod* 1980;43(2):169–234. doi:10.1021/np50008a001, PMID:6991645.
- [53] Chen F, Tholl D, Bohlmann J, Pichersky E. The family of terpene synthases in plants: a mid-size family of genes for specialized metabolism that is highly diversified throughout the kingdom. *Plant J* 2011;66(1):212–229. doi:10.1111/j.1365-3113.2011.04520.x, PMID:21443633.
- [54] Nissen L, Zatta A, Stefanini I, Grandi S, Sgorbati B, Biavati B, *et al.* Characterization and antimicrobial activity of essential oils of industrial hemp varieties (*Cannabis sativa* L.). *Fitoterapia* 2010;81(5):413–419. doi:10.1016/j.fitote.2009.11.010, PMID:19969046.
- [55] Nostro A, Guerrini A, Marino A, Tacchini M, Di Giulio M, Grandini A, *et al.* In vitro activity of plant extracts against biofilm-producing food-related bacteria. *Int J Food Microbiol* 2016;238:33–39. doi:10.1016/j.ijfoodmicro.2016.08.024, PMID:27591384.
- [56] Schofs L, Sparo MD, Sánchez Bruni SF. The antimicrobial effect behind Cannabis sativa. *Pharmacol Res Perspect* 2021;9(2):e00761. doi:10.1002/prp2.761, PMID:33822478.
- [57] Onsare JG, Arora DS. Antibiofilm potential of flavonoids extracted from Moringa oleifera seed coat against Staphylococcus aureus, Pseudomonas aeruginosa and Candida albicans. *J Appl Microbiol* 2015;118(2):313–325. doi:10.1111/jam.12701, PMID:25410525.
- [58] Das B, Mishra P. Antibacterial analysis of crude extracts from the leaves of Tagetes erecta and Cannabis sativa. *International Journal of Environmental Sciences* 2012;2(3):1605–1608. doi:10.6088/ijes.00202030045.
- [59] Sambasivam S, Karpagam G, Chandran R, Khan SA. Toxicity of leaf extract of yellow oleander Thevetia nerifolia on Tilapia. *J Environ Biol* 2003;24(2):201–204. PMID:12974464.
- [60] Kayser O, Kiderlen AF, Croft SL. Natural products as potential antiparasitic drugs. *Studies in Natural Products Chemistry* 2002;26:779–848. doi:10.1016/S1572-5995(02)80019-9.
- [61] Varma J, Dubey N. Prospectives of botanical and microbial products as pesticides of tomorrow. *Current Science* 1999;76(2):172–179.
- [62] Cowan MM. Plant products as antimicrobial agents. *Clin Microbiol Rev* 1999;12(4):564–582. doi:10.1128/CMR.12.4.564, PMID:10515903.
- [63] Audu B, Ofojekwu P, Ujah A, Ajima M. Phytochemical, proximate composition, amino acid profile and characterization of marijuana (*Cannabis sativa* L.). *The Journal of Phytopharmacology* 2014;3(1):35–43. doi:10.31254/phyto.2014.3106.
- [64] Pollastro F, Minassi A, Fresu LG. Cannabis Phenolics and their Bioactivities. *Curr Med Chem* 2018;25(10):1160–1185. doi:10.2174/0929867324666170810164636, PMID:28799497.
- [65] Zagožen M, Čerenak A, Krefl S. Cannabigerol and cannabichromene in Cannabis sativa L. *Acta Pharmaceutica* 2021;71(3):355–364. doi:10.2478/acph-2021-0021.
- [66] Sampson PB. Phytocannabinoid Pharmacology: Medicinal Properties of Cannabis sativa Constituents Aside from the “Big Two”. *J Nat Prod* 2021;84(1):142–160. doi:10.1021/acs.jnatprod.0c00965, PMID:33356248.
- [67] Gildea L, Ayariga JA, Ajayi OS, Xu J, Villafane R, Samuel-Foo M. Cannabis sativa CBD Extract Shows Promising Antibacterial Activity against Salmonella typhimurium and S. newington. *Molecules* 2022;27(9):2669. doi:10.3390/molecules27092669, PMID:35566019.
- [68] Gray RA, Whalley BJ. The proposed mechanisms of action of CBD in epilepsy. *Epileptic Disord* 2020;22(S1):10–15. doi:10.1684/epd.2020.1135, PMID:32053110.
- [69] Cannabinoids as antimicrobial agents. New York: LAVVAN; 2022. Available from: <https://www.lavvan.com/cannabinoids-as-antimicrobial-agents>. Accessed May 9, 2022.
- [70] Aqawi M, Sionov RV, Gallily R, Friedman M, Steinberg D. Anti-Bacterial Properties of Cannabigerol Toward Streptococcus mutans. *Front Microbiol* 2021;12:656471. doi:10.3389/fmicb.2021.656471, PMID:33967995.
- [71] Bucekova M, Jardekova L, Juricova V, Bugarova V, Di Marco G, Gismondi A, *et al.* Antibacterial Activity of Different Blossom Honey: New Findings. *Molecules* 2019;24(8):E1573. doi:10.3390/molecules24081573, PMID:31010070.
- [72] Martinenghi LD, Jønsson R, Lund T, Jenssen H. Isolation, Purification, and Antimicrobial Characterization of Cannabidiolic Acid and Cannabidiol from Cannabis sativa L. *Biomolecules* 2020;10(6):E900. doi:10.3390/biom10060900, PMID:32545687.
- [73] Krejci Z, Horok M, Santavy F. Hanf (cannabis sativa)-antibiotisches heilmittel. 3. Mitteilung: Isolierung und Konstitution zweier aus Cannabis sativa gewonnener sauren. *Die Pharmazie* 1959;14:349–355.
- [74] Izzo AA, Borrelli F, Capasso R, Di Marzo V, Mechoulam R. Non-psychoactive plant cannabinoids: new therapeutic opportunities from an ancient herb. *Trends Pharmacol Sci* 2009;30(10):515–527. doi:10.1016/j.tips.2009.07.006, PMID:19729208.
- [75] Dalzell HC, Uliss DB, Handrick GR, Razdan RK. Hashish. 26. Factors influencing double-bond stability in cannabinoids. *The Journal of Organic Chemistry* 1981;46(5):949–953. doi:10.1021/jo00318a021.
- [76] Clarke H, Roychoudhury P, Narouze SN. Other Phytocannabinoids. In: Narouze SN (ed). *Cannabinoids and Pain*. Cham: Springer; 2021:87–92. doi:10.1007/978-3-030-69186-8\_12.
- [77] Tahsin KN, Watson D, Rizkalla A, Heinrichs D, Charpentier P. Antimicrobial Studies of Cannabidiol as Biomaterials against superbug MRSA. *CMBES Proc* 2021;44.
- [78] Palomares Cañero B. Non psychoactive cannabinoids for the treatment of inflammatory diseases [Dissertation]. Universidad de Córdoba, UCO Press; 2020. Available from: <http://hdl.handle.net/10396/19632>. Accessed May 9, 2022.
- [79] Mnekni L, Ripoll L. Topical Use of Cannabis sativa L. *Biochemicals. Cosmetics* 2021;8(3):85. doi:10.3390/cosmetics8030085.
- [80] Khameneh B, Iranshahy M, Soheili V, Fazly Bazzaz BS. Review on plant antimicrobials: a mechanistic viewpoint. *Antimicrob Resist Infect Control* 2019;8:118. doi:10.1186/s13756-019-0559-6, PMID:31346459.
- [81] Sytar O, Brestic M, Hajhashemi S, Skalicky M, Kubeš J, Lamilla-Tamayo L, *et al.* COVID-19 Prophylaxis Efforts Based on Natural Antiviral Plant Extracts and Their Compounds. *Molecules* 2021;26(3):727. doi:10.3390/molecules26030727, PMID:33573318.
- [82] Pitakbut T, Nguyen GN, Kayser O. Activity of THC, CBD, and CBN on Human ACE2 and SARS-CoV1/2 Main Protease to Understand Antiviral Defense Mechanism. *Planta Med* 2022;88(12):1047–1059. doi:10.1055/a-1581-3707, PMID:34638139.
- [83] Onay A, Ertas A, Süzerer V, Yener İ, Yilmaz MA, Ayaz-Tilkat E, *et al.* Cannabinoids for SARS-CoV-2 and is there evidence of their therapeutic efficacy? *Turk J Biol* 2021;45(4):570–587. doi:10.3906/biy-2105-73, PMID:34803455.
- [84] Yadav V, Kaushik P. Phytochemicals against COVID-19 and a gap in clinical investigations: An outlook. *Indian Journal of Biochemistry Biophysics* 2021;58(5):403–407.
- [85] Zhao L, Yan Y, Dai Q, Wang Z, Yin J, Xu Y, *et al.* The CDK1 inhibitor, Ro-3306, is a potential antiviral candidate against influenza virus infection. *Antiviral Res* 2022;201:105296. doi:10.1016/j.antiviral.2022.105296, PMID:35367281.
- [86] Mabou Tagne A, Pacchetti B, Sodergren M, Cosentino M, Marino F. Cannabidiol for Viral Diseases: Hype or Hope? *Cannabis Cannabinoid Res* 2020;5(2):121–131. doi:10.1089/can.2019.0060, PMID:32656344.
- [87] Stieltjes N, Ounnoughene N, Sava E, Paugy P, Roussel-Robert V, Rosenberg AR, *et al.* Interest of transjugular liver biopsy in adult patients with haemophilia or other congenital bleeding disorders infected with hepatitis C virus. *Br J Haematol* 2004;125(6):769–776. doi:10.1111/j.1365-2141.2004.04968.x, PMID:15180867.
- [88] Leopold SJ, Grady BP, Lindenburg CE, Prins M, Beuers U, Weegink CJ. Common bile duct dilatation in drug users with chronic hepatitis C is associated with current methadone use. *J Addict Med* 2014;8(1):53–58. doi:10.1097/ADM.000000000000006, PMID:24394497.
- [89] Lowe HI, Toyang NJ, McLaughlin W. Potential of Cannabidiol for the Treatment of Viral Hepatitis. *Pharmacognosy Res* 2017;9(1):116–118. doi:10.4103/0974-8490.199780, PMID:28250664.
- [90] Bachmetov L, Gal-Tanamy M, Shapira A, Vorobeychik M, Giterman-Galam T, Sathiyamoorthy P, *et al.* Suppression of hepatitis C virus by the flavonoid quercetin is mediated by inhibition of NS3 protease activity. *J Viral Hepat* 2012;19(2):e81–e88. doi:10.1111/j.1365-2893.2011.01507.x, PMID:22239530.
- [91] Edmunds BL, Miller ER, Tsourtos G. The distribution and socio-economic burden of Hepatitis C virus in South Australia: a cross-sectional study 2010-2016. *BMC Public Health* 2019;19(1):527. doi:10.1186/s12889-019-6847-5, PMID:31068170.



- [92] de Paula Farah K, Carmo RA, de Figueiredo Antunes CM, Serufo JC, Nobre Júnior VA, Fonseca de Castro LP, *et al.* Hepatitis C, HCV genotypes and hepatic siderosis in patients with chronic renal failure on haemodialysis in Brazil. *Nephrol Dial Transplant* 2007;22(7):2027–2031. doi:10.1093/ndt/gfm028, PMID:17309883.
- [93] Hegde VL, Nagarkatti PS, Nagarkatti M. Role of myeloid-derived suppressor cells in amelioration of experimental autoimmune hepatitis following activation of TRPV1 receptors by cannabidiol. *PLoS One* 2011;6(4):e18281. doi:10.1371/journal.pone.0018281, PMID:21483776.
- [94] Lowe H, Steele B, Bryant J, Fouad E, Toyang N, Ngwa W. Antiviral Activity of Jamaican Medicinal Plants and Isolated Bioactive Compounds. *Molecules* 2021;26(3):607. doi:10.3390/molecules26030607, PMID:33503834.
- [95] Maor Y, Yu J, Kuzontkoski PM, Dezube BJ, Zhang X, Groopman JE. Cannabidiol inhibits growth and induces programmed cell death in kaposi sarcoma-associated herpesvirus-infected endothelium. *Genes Cancer* 2012;3(7-8):512–520. doi:10.1177/1947601912466556, PMID:23264851.
- [96] Camacho-Rivera M, Islam JY, Rodriguez DL, Vidot DC. Cannabis Use among Cancer Survivors amid the COVID-19 Pandemic: Results from the COVID-19 Cannabis Health Study. *Cancers (Basel)* 2021;13(14):3495. doi:10.3390/cancers13143495, PMID:34298708.
- [97] Beasley MB. Acute lung injury from cannabis to COVID. *Mod Pathol* 2022;35(Suppl 1):1–7. doi:10.1038/s41379-021-00915-6, PMID:34504310.
- [98] van Laar MW, Oomen PE, van Miltenburg CJA, Vercoulen E, Freeman TP, Hall WD. Cannabis and COVID-19: Reasons for Concern. *Front Psychiatry* 2020;11:601653. doi:10.3389/fpsy.2020.601653, PMID:33408655.
- [99] Chong WW, Acar ZI, West ML, Wong F. A Scoping Review on the Medical and Recreational Use of Cannabis During the COVID-19 Pandemic. *Cannabis Cannabinoid Res* 2022;7(5):591–602. doi:10.1089/can.2021.0054, PMID:34981958.
- [100] Chatow L, Nudel A, Neshet I, Hayo Hemo D, Rozenberg P, Voropaev H, *et al.* In Vitro Evaluation of the Activity of Terpenes and Cannabidiol against Human Coronavirus E229. *Life (Basel)* 2021;11(4):290. doi:10.3390/life11040290, PMID:33805385.
- [101] Loizzo MR, Saab AM, Tundis R, Menichini F, Bonesi M, Piccolo V, *et al.* In vitro inhibitory activities of plants used in Lebanon traditional medicine against angiotensin converting enzyme (ACE) and digestive enzymes related to diabetes. *J Ethnopharmacol* 2008;119(1):109–116. doi:10.1016/j.jep.2008.06.003, PMID:18601990.
- [102] Nguyen LC, Yang D, Nicolaescu V, Best TJ, Ohtsuki T, Chen SN, *et al.* Cannabidiol Inhibits SARS-CoV-2 Replication and Promotes the Host Innate Immune Response. *bioRxiv* 2021. doi:10.1101/2021.03.10.432967, PMID:33758843.
- [103] Perry E, Howes MJ. Medicinal plants and dementia therapy: herbal hopes for brain aging? *CNS Neurosci Ther* 2011;17(6):683–698. doi:10.1111/j.1755-5949.2010.00202.x, PMID:22070157.
- [104] Astuti I, Ysrafil. Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2): An overview of viral structure and host response. *Diabetes Metab Syndr* 2020;14(4):407–412. doi:10.1016/j.dsx.2020.04.020, PMID:32335367.
- [105] Davidson AM, Wysocki J, Battle D. Interaction of SARS-CoV-2 and Other Coronavirus With ACE (Angiotensin-Converting Enzyme)-2 as Their Main Receptor: Therapeutic Implications. *Hypertension* 2020;76(5):1339–1349. doi:10.1161/HYPERTENSIONAHA.120.15256, PMID:32851855.
- [106] Sallenave JM, Guillot L. Innate Immune Signaling and Proteolytic Pathways in the Resolution or Exacerbation of SARS-CoV-2 in Covid-19: Key Therapeutic Targets? *Front Immunol* 2020;11:1229. doi:10.3389/fimmu.2020.01229, PMID:32574272.
- [107] Anand U, Jacobo-Herrera N, Altemimi A, Lakhssassi N. A Comprehensive Review on Medicinal Plants as Antimicrobial Therapeutics: Potential Avenues of Biocompatible Drug Discovery. *Metabolites* 2019;9(11):E258. doi:10.3390/metabo9110258, PMID:31683833.
- [108] Cooper MA, Shlaes D. Fix the antibiotics pipeline. *Nature* 2011;472(7341):32. doi:10.1038/472032a, PMID:21475175.
- [109] WHO. The Burden of Health Care-Associated Infection Worldwide. A Summary; 2010: Geneva.
- [110] Datta S, Ramamurthy PC, Anand U, Singh S, Singh A, Dhanjal DS, *et al.* Wonder or evil?: Multifaceted health hazards and health benefits of *Cannabis sativa* and its phytochemicals. *Saudi J Biol Sci* 2021;28(12):7290–7313. doi:10.1016/j.sjbs.2021.08.036, PMID:34867033.
- [111] Eloff JN. A sensitive and quick microplate method to determine the minimal inhibitory concentration of plant extracts for bacteria. *Planta Med* 1998;64(8):711–713. doi:10.1055/s-2006-957563, PMID:9933989.
- [112] CLSI Performance Standards for Antimicrobial Disk Susceptibility Tests. 12th Edition ed. Wayne, PA: Clinical and Laboratory Standards Institute; 2017.
- [113] Sadgrove NJ, Jones GL. From Petri Dish to Patient: Bioavailability Estimation and Mechanism of Action for Antimicrobial and Immunomodulatory Natural Products. *Front Microbiol* 2019;10:2470. doi:10.3389/fmicb.2019.02470, PMID:31736910.
- [114] Al Ubeed HMS, Wills RBH, Chandrapala J. Post-Harvest Operations to Generate High-Quality Medicinal Cannabis Products: A Systemic Review. *Molecules* 2022;27(5):1719. doi:10.3390/molecules27051719, PMID:35268820.
- [115] Challa SKR, Misra N, Martynenko A. Drying of cannabis—state of the practices and future needs. *Drying Technology* 2021;39(14):2055–2064.
- [116] Chasiotis V, Tsakirakis A, Termentzi A, Machera K, Filios A. Drying and quality characteristics of Cannabis sativa L. Inflorescences under constant and time-varying convective drying temperature schemes. *Thermal Science and Engineering Progress* 2022;28:101076. doi:10.1016/j.tsep.2021.101076.
- [117] Frassinetti S, Gabriele M, Moccia E, Longo V, Di Gioia D. Antimicrobial and antibiofilm activity of Cannabis sativa L. seeds extract against *Staphylococcus aureus* and growth effects on probiotic *Lactobacillus* spp. *LWT* 2020;124:109149. doi:10.1016/j.lwt.2020.109149.
- [118] Patnaik T, Dey R, Gouda P. Antimicrobial Activity of Friedelan-3 [beta]-ol and trans-N-Caffeoyltyramine Isolated from the Root of *Vitis trifolia*. *Asian Journal of Chemistry* 2008;20(1):417–421.
- [119] Di Gioia D, Gaggia F, Baffoni L, Stenico VJB. Role of bifidobacteria in the production of bioactive compounds and detoxification of harmful compounds. *Beneficial microbes in fermented functional foods*. 2014:291–308.
- [120] Iseppi R, Brighenti V, Licata M, Lambertini A, Sabia C, Messi P, *et al.* Chemical Characterization and Evaluation of the Antibacterial Activity of Essential Oils from Fibre-Type *Cannabis sativa* L. (Hemp). *Molecules* 2019;24(12):E2302. doi:10.3390/molecules24122302, PMID:31234360.
- [121] Chauhan A, Verma R, Kumari S, Sharma A, Shandilya P, Li X, *et al.* Photocatalytic dye degradation and antimicrobial activities of Pure and Ag-doped ZnO using Cannabis sativa leaf extract. *Sci Rep* 2020;10(1):7881. doi:10.1038/s41598-020-64419-0, PMID:32398650.
- [122] Csakvari AC, Moisa C, Radu DG, Olariu LM, Lupitu AI, Panda AO, *et al.* Green Synthesis, Characterization, and Antibacterial Properties of Silver Nanoparticles Obtained by Using Diverse Varieties of *Cannabis sativa* Leaf Extracts. *Molecules* 2021;26(13):4041. doi:10.3390/molecules26134041, PMID:34279380.
- [123] Singh P, Pandit S, Garnæs J, Tunjic S, Mokkalapati VR, Sultan A, *et al.* Green synthesis of gold and silver nanoparticles from *Cannabis sativa* (industrial hemp) and their capacity for biofilm inhibition. *Int J Nanomedicine* 2018;13:3571–3591. doi:10.2147/IJN.S157958, PMID:29950836.
- [124] Ying S, Guan Z, Ying S, Ofoegbu PC, Clubb P, Rico C, *et al.* Green synthesis of nanoparticles: Current developments and limitations. *Environmental Technology & Innovation*. 2022:102336. doi:10.1016/j.eti.2022.102336.
- [125] Pal G, Rai P, Pandey A. Green synthesis of nanoparticles: A greener approach for a cleaner future. *Green synthesis, characterization and applications of nanoparticles*. Elsevier; 2019:1–26. doi:10.1016/B978-0-08-102579-6.00001-0.
- [126] Chakaraborty PS, Sapkota H, Prabhakar PK. Synergistic interaction of cannabis and garlic with commercial antibiotics. *International Journal of Pharmacognosy Phytochemical Research* 2015;7(1):193–196.
- [127] Nuutinen T. Medicinal properties of terpenes found in Cannabis sativa and Humulus lupulus. *Eur J Med Chem* 2018;157:198–228. doi:10.1016/j.ejmech.2018.07.076, PMID:30096653.

- [128] Blesson J, Saji C, Nivya R, Kumar R. Synergism of antibiotics and plant extracts in antibacterial activity against methicillin resistant *Staphylococcus aureus*. *Journal of Bioavailability & Bioequivalence* 2014;6(5):.
- [129] Janecki M, Graczyk M, Lewandowska AA, Pawlak Ł. Anti-Inflammatory and Antiviral Effects of Cannabinoids in Inhibiting and Preventing SARS-CoV-2 Infection. *Int J Mol Sci* 2022;23(8):4170. doi:10.3390/ijms23084170, PMID:35456990.
- [130] Sionov RV, Steinberg D. Anti-Microbial Activity of Phytocannabinoids and Endocannabinoids in the Light of Their Physiological and Pathophysiological Roles. *Biomedicines* 2022;10(3):631. doi:10.3390/biomedicines10030631, PMID:35327432.
- [131] Munjal M, ElSohly MA, Repka MA. Chemical stabilization of a Delta9-tetrahydrocannabinol prodrug in polymeric matrix systems produced by a hot-melt method: role of microenvironment pH. *AAPS PharmSciTech* 2006;7(3):71. doi:10.1208/pt070371, PMID:17025251.
- [132] Kumari A, Yadav SK, Pakade YB, Singh B, Yadav SC. Development of biodegradable nanoparticles for delivery of quercetin. *Colloids Surf B Biointerfaces* 2010;80(2):184–192. doi:10.1016/j.col-surf.2010.06.002, PMID:20598513.
- [133] Fraguas-Sánchez AI, Fernández-Carballido A, Martín-Sabroso C, Torres-Suárez AI. Stability characteristics of cannabidiol for the design of pharmacological, biochemical and pharmaceutical studies. *J Chromatogr B Analyt Technol Biomed Life Sci* 2020;1150:122188. doi:10.1016/j.jchromb.2020.122188, PMID:32506012.