



Review Article

Underwater Resection of Neuroendocrine Tumors of the Gastrointestinal Tract: A Systematic Review



Paula Arruda do Espirito Santo¹, Matheus Henrique Gonçalves de Souza^{2*} ,
Isabelle Kristal Grala Souza e Silva³, Fabio Shiguehissa Kawaguti⁴, Fauze Maluf-Filho^{3,5} and Luciano Lenz^{3,4}

¹Diagnostic Imaging and Specialized Diagnosis Unit, Hospital Universitario da Universidade Federal de São Carlos, São Carlos, Brazil; ²Department of Internal Medicine, Faculdade de Medicina da Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil; ³Department of Gastroenterology, University of São Paulo, Instituto do Cancer do Estado de São Paulo, São Paulo, Brazil; ⁴Fleury Medicina e Saúde, São Paulo, Brazil; ⁵National Council for Scientific and Technological Development – CNPq, Brasília, Brazil

Received: June 14, 2023 | Revised: August 22, 2023 | Accepted: September 08, 2023 | Published online: September 25, 2023

Abstract

Background and objectives: Over the past decade, the approach to gastrointestinal (GI) neuroendocrine tumors (NETs) has increasingly included endoscopic resection (ER) techniques. Underwater endoscopic mucosal resection (UEMR) has been introduced as a potential alternative to conventional mucosectomy. The objective of this systematic review is to investigate the feasibility and outcomes of UEMR in the treatment of GI NETs and to provide a reference for clinical management options.

Methods: A systematic search of PubMed, Cochrane Library, and EMBASE was performed to identify guidelines and primary literature published up to 8 August 2023.

Results: A total of 12 studies comprising 98 cases were included in this systematic review. No UEMR studies were identified for esophageal NETs. For gastric NETs, only one pilot study with two patients was available, reporting complete resection without complications. Eleven cases of duodenal NETs were reported, all successfully resected. Ileal NETs were documented in a single outlier case. The most substantial evidence exists for rectal NETs, with four comparative studies supporting UEMR as a viable alternative to conventional endoscopic resection techniques.

Conclusions: This systematic review synthesizes limited evidence, suggesting that UEMR may be a feasible option for well-selected rectal NETs, offering potential advantages in terms of efficiency and technical simplicity. For other gastrointestinal sites, evidence is currently insufficient to support its routine use, highlighting the need for further comparative studies.

Introduction

Neuroendocrine tumors (NETs) are rare and heterogeneous tumors that arise from the neuroendocrine cell system. NETs are more

commonly found in the gastrointestinal (GI) tract, with the most common sites being the stomach (23%), appendix (21%), small bowel (15%), rectum (14%) and pancreas (10%). Esophageal and colonic NETs are rare and account for a small percentage of GI NETs.¹ The incidence of GI NETs has gradually increased over the last few decades due to improved detection and an aging population.² The prognosis of these tumors is highly variable and depends on several factors including site, size, grading, depth of invasion, and local lymphadenopathy. Correct evaluation of the tumor and the clinical context has important prognostic implications. Careful assessment will also help determine the treatment modality. Over the past decade, the approach to GI NETs has progressively included endoscopic resection (ER) techniques. The indication for endoscopic treatment does not include NETs of the appendix, colon, or biliary tree.³ For pancreatic NETs, the appropriate endoscopic approaches are restricted to ablative techniques.¹ For other sites, ER briefly includes endoscopic mucosal resection (EMR), endoscopic

Keywords: Underwater endoscopic mucosal resection; Gastrointestinal neuroendocrine tumour; Systematic review; Endoscopy.

Abbreviations: EMR, endoscopic mucosal resection; ER, endoscopic resection; ESD, endoscopic submucosal dissection; GI, gastrointestinal; JBI, Joanna Briggs Institute; NET, neuroendocrine tumor; PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses; UEMR, underwater endoscopic mucosal resection.

***Correspondence to:** Matheus Henrique Gonçalves de Souza, Department of Internal Medicine, Faculdade de Medicina da Universidade Federal do Rio de Janeiro, Rio de Janeiro 21044-020, Brazil. ORCID: <https://orcid.org/0009-0005-3695-9701>. Tel: +55-21-39382731, Fax: +55-21-39382735, E-mail: matheushenrique.gs@hotmail.com

How to cite this article: do Espirito Santo PA, de Souza MHG, Silva IKG, Kawaguti FS, Maluf-Filho F, Lenz L. Underwater Resection of Neuroendocrine Tumors of the Gastrointestinal Tract: A Systematic Review. *J Transl Gastroenterol* 2023;1(1):13–21. doi: 10.14218/JTG.2023.00031.




Fig. 1. UEMR technique. Initially the luminal gas is aspirated (a) and water is infused (b) until the mucosal lesion floats in a completely water-filled lumen to lift the lesion and create a pseudopedicle (c), making it feasible to use a snare (d).

submucosal dissection (ESD), and full-thickness resection.

Although ESD appears to be a good alternative as it allows *en bloc* resection of large lesions, this technique requires advanced skills and is not widely available, and it is also prone to postoperative adverse events such as bleeding or perforation.⁴⁻⁶ In comparison, EMR may fail to completely remove submucosal tumors.⁵

Binmoeller *et al.*⁷ also developed an alternative method of lesion removal without submucosal injection. Underwater endoscopic mucosal resection (UEMR) is a new method that presents as an alternative to the conventional EMR. UEMR is performed without lifting the lesion with submucosal injection, but rather using the ability of water to lift the lesion. Filling the lumen with water allows the lesion to be lifted and the mucosa and submucosal tumor separate from the muscularis propria, creating a pseudopedicle that facilitates the use of a snare (Fig. 1).^{5,8-11} Many articles have been published showing reasonable technical success rates, with a low incidence of adverse events with UEMR. However, the vast majority of these studies have been of UEMR for the resection of colorectal adenomas.^{7,12-18} Since the first description by Kawaguti *et al.*,¹⁹ only a few studies of UEMR in GI NETs have been described.

Methods

Protocol registration

This systematic review was registered at the International Prospective Register of Systematic Reviews (PROSPERO) under protocol CRD42023451193. This study was designed according to the guidelines of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement guidelines.²⁰

Search strategy and study selection

We systematically searched PubMed, EMBASE, and Cochrane Library databases to identify studies (published up to August 8, 2023) reporting the use of UEMR technique in GI NETs. The search strategy was (underwater OR UEMR) AND (neuroendocrine OR NET OR carcinoid) (tumor OR tumors OR lesion OR lesions) AND (gastrointestinal OR (esophagus OR oesophagus) OR (gastric OR stomach) OR (small bowel OR (duodenal OR duodenum) OR (ileum OR ileal) OR (colon* OR colorectal) OR (rectum OR rectal).

Two authors (PAES and MHGS) independently screened all studies identified by the database search, initially based on titles and abstracts. Subsequently, full texts of potentially relevant articles were reviewed for inclusion if they met the predefined criteria. We also analyzed references from relevant original articles and literature reviews to identify additional studies that were not encompassed in the initial database searches. Disagreements were resolved through consensus. Studies meeting the following inclusion criteria were included: (1) observational study (including case report, case series, cohort studies and RCTs); (2) comparing UEMR with other ER technique; (3) in patients with GI NET; (4) reporting outcomes of interest. Abstracts, editorials, reviews, and meta-analyses were excluded.

Data extraction and quality assessment

For all eligible studies, we extracted data including (1) study characteristics: first author, study design, publication year, country; (2) study population: number of cases, age, sex, NET site and tumor characteristics (location, size, and grade); (3) outcomes of interest. Primary outcomes were *en bloc* resection, R0 resection, complete resection, and complications. Secondary outcomes were procedure time and cost. *En bloc* resection was defined as the removal of a lesion in a single piece confirmed by endoscopy. R0 resection was defined as *en bloc* resection with both horizontal and vertical margins free of any adenomatous tissue, as confirmed by histological analysis. Complete resection was defined as the lack of visible lesions on a macroscopic scale on endoscopic examination. Complications were defined as any adverse event during or after the procedure.

The quality assessment of included studies was independently performed by two authors (PAES and MHGS) using modified versions of the Joanna Briggs Institute (JBI) critical appraisal tool to evaluate the risk of bias in case reports, case series, and cohort studies.^{21,22} The JBI critical appraisal checklist consists of confirmatory questions about study characteristics that are scored as yes/no, unclear or not applicable (Supplementary Table 1). Disagreements were resolved by consensus.

Data synthesis and analysis

We provided a narrative analysis of the results of the included studies. For didactic purposes, the topics were divided into specific organs. We summarized the main reports on the use of the UEMR




Fig. 2. PRISMA flow diagram of study screening and selection. PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

technique in each GI NET site and assessed the feasibility of this technique from a comprehensive perspective. Reference clinical management options recommended by the European Neuroendocrine Tumor Society (ENETS) and the North American Neuroendocrine Tumor Society (NANETS) guidelines were also highlighted. Data from each eligible article were synthesized, described qualitatively, and presented in tabular form. Categorical variables were reported as raw numbers (percentages). Continuous variables were reported as means \pm SD, medians (range), or medians (IQR).

Results

The literature search initially identified 64 records. After removing 18 duplicate records and 25 unrelated studies based on the titles and abstracts, we identified 21 potentially eligible studies that were screened for inclusion and exclusion criteria. Marui *et al.*²³ were also identified by citation searching. Ultimately, 12 studies were included in this review. This process is illustrated in the PRISMA flow diagram (Fig. 2).

Characteristics of included studies

Overall, these studies aggregate data from 98 cases of the UEMR technique in GI NETs. Of these, two cases were in gastric NETs, 11 cases were in duodenal NETs, one case was in ileum NET, and 84 cases were in rectal NETs. Seven studies were conducted in Asia (South Korea, Japan, and China), three studies were conducted in Brazil, and two studies were conducted in Europe (Italy

and Spain). Table 1 summarizes the main characteristics of the included studies. As reported in Table 2,^{5,9,19,23-31} based on the JBI critical appraisal checklist assessment tool, the quality assessment ranged from 7/11 to 11/11, suggesting a moderate overall risk of bias.

UEMR in esophageal NETs

NETs are extremely rare in the esophagus, accounting for less than 0.04% of all NETs. They are characterized by early dissemination, aggressive behavior, and poor prognosis. Due to their rarity, no standardized guidelines have been proposed for the treatment of esophageal NETs (eNETs), although general recommendations suggest surgery with adjuvant chemoradiotherapy as the treatment of choice.³²⁻³⁴

Few cases of endoscopic removal of eNETs have been described, usually by EMR or ESD and more recently by submucosal tunneling ER.^{35,36}

We found only three studies of underwater esophageal resection. One case of UEMR for a protruding well-differentiated intramucosal adenocarcinoma, one report of UEMR for a granular cell tumor and one series of nine patients with superficial esophageal tumors treated with underwater ESD.³⁷⁻³⁹ However, our research did not find any UEMR results for eNETs.

UEMR in gastric NETs

Gastric NETs (gNETs) can be divided into three types, type 1 and 2, which are associated with chronic atrophic gastritis and

Table 1. Characteristics of included studies

NET site	Study characteristics				Study population				Outcomes			
	First author (country, year)	Study design	Number of cases	Sex, female/male	Age, years	Location	Size, mm	Grade, G1/G2	En bloc resection, n (%)	R0 resection, n (%)	Complete resection, n (%)	Complication
Gastric	Kim (South Korea, 2022) ²⁴	Case series	2	1/1	65 and 68	Greater curvature of stomach: 2	11 and 5	2/0	2 (100)	2 (100)	2 (100)	No
Duodenal	Anderloni (Italy, 2016) ²⁵	Case report	1	0/1	68	Anterior wall of the duodenal bulb	15	0/1	1 (100)	1 (100)	1 (100)	No
	Kim (South Korea, 2022) ²⁴	Case series	7	3/4	Median (range): 60 (36–67)	Duodenal bulb: 5; superior descending angle: 1; near ampulla: 1	Median (range): 6 (3–15)	7/0	7 (100)	7 (100)	7 (100)	No
	Uchima (Spain, 2022) ²⁶	Case report	1	0/1	64	Duodenal bulb	13	1/0	1 (100)	1 (100)	1 (100)	No
	Lee (South Korea, 2022) ²⁷	Case series	2	0/2	36 and 61	Duodenal bulb: 2	3 and 6	2/0	2 (100)	1 (50)	NA	No
Ileum	Rezende (Brazil, 2021) ²⁸	Case report	1	0/1	64	Distal ileum	10	1/0	1 (100)	1 (100)	1 (100)	No
Rectal	Kawaguti (Brazil, 2015) ¹⁹	Case report	1	1/0	51	Distal rectum	10	1/0	1 (100)	1 (100)	1 (100)	No
	Marui (Japan, 2016) ²³	Case report	1	0/1	63	Lower colorectal area	8	1/0	1 (100)	1 (100)	1 (100)	Minor bleeding
	Yamashina (Japan, 2018) ³¹	Case series	6	3/3	Median (range): 55.5 (47–86)	Rectum above the peritoneal reflection: 2; rectum below the peritoneal reflection: 4;	Median (range): 5.5 (4–10)	6/0	6 (100)	5 (83.3)	NA	No
	Coutinho (Brazil, 2021) ³⁰	Case series	11	9/2	Median (range): 55 (30–73)	Rectum	Median (range): 6 (3–12)	9/2	11 (100)	9 (81.8)	NA	No
	Park (South Korea, 2020) ⁹	Retrospective cohort	36	14/22	Median (range): 44 (23–76)	Distance from anal verge (cm), median (range): 5 (3–15)	Median (range): 5 (2–10)	36/0	≥31 (86.1)*	31 (86.1)	NA	No
	Shi (China, 2022) ⁵	Retrospective cohort	22	8/14	Mean (SD): 46.4 (9.3)	Distance from the anal margin (cm), mean (SD): 7.7 (1.6)	Mean (SD): 7.2 (1.6)	NA	NA	NA	21 (95.5)	No
	Okada (Japan, 2022) ²⁹	Retrospective cohort	7	3/4	Median (IQR): 65 (50.5–67.5)	Rectum	Median (IQR): 4 (3–5)	7/0	7 (100)	7 (100)	NA	No

IQR, interquartile range; NA, not available; NET, neuroendocrine tumor; SD, standard deviation. *Park et al. (8) does not provide the values of en bloc resection; however, we assume that it is greater than or equal to the value of R0 resection.

Table 2. Quality assessment of included studies according to JBI critical appraisal tool

First author, year	Study design	Q1	Q2	Q3	Q4	Q5	Q6	Q7	Q8	Q9	Q10	Q11	Total
Anderloni, 2016 ²⁵	Case report	Y	Y	Y	Y	Y	Y	Y	Y	–	–	–	8/8
Uchima, 2022 ²⁶	Case report	Y	Y	Y	Y	Y	Y	Y	Y	–	–	–	8/8
Rezende, 2021 ²⁸	Case report	Y	Y	Y	Y	Y	Y	Y	Y	–	–	–	8/8
Kawaguti, 2015 ¹⁹	Case report	Y	Y	Y	Y	Y	Y	Y	Y	–	–	–	8/8
Marui, 2016 ²³	Case report	Y	Y	Y	Y	Y	Y	Y	Y	–	–	–	8/8
Yamashina, 2018 ³¹	Case series	Y	Y	Y	Y	Y	Y	Y	Y	Y	U	–	9/10
Kim, 2022 ²⁴	Case series	Y	Y	Y	Y	U	Y	Y	Y	Y	U	–	8/10
Lee, 2022 ²⁷	Case series	Y	Y	Y	Y	U	Y	Y	Y	Y	U	–	8/10
Coutinho, 2021 ³⁰	Case series	Y	Y	Y	Y	Y	Y	Y	Y	Y	U	–	9/10
Park, 2020 ⁹	Retrospective	U	Y	Y	Y	N	Y	Y	Y	U	N	Y	7/11
Shi, 2022 ⁵	Retrospective	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	11/11
Okada, 2022 ²⁹	Retrospective	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	11/11

JBI, Joanna Briggs Institute. N, no; NA, not applicable; Q, question; U, unclear; Y, yes; –, respective question does not exist.

Zollinger-Ellison syndrome, respectively, and type 3, which is rare but aggressive.⁴⁰ Surgical resection with lymphadenectomy is recommended for type 3 tumors due to their high-grade malignancy. In contrast, conservative management strategies are preferred for types 1 and 2 gNETs, with ER being recommended for tumors ≥ 10 mm.⁴⁰ However, ER is not indicated for type 3 gNETs. Recommendations for ER differ between the ENETS and NANETS guidelines. ENETS recommends surgical removal for all lesions ≥ 10 mm without muscularis propria invasion, whereas NANETS recommends surgical removal for lesions >20 mm and for lesions measuring 10–20 mm in size if *muscularis propria* invasion is present.^{40,41} The first and only UEMR procedure for gNETs was performed and reported by Kim *et al.*²⁴ in 2022 in a case series pilot study involving two patients with G1 NETs (5 mm and 11 mm lesion size) located in the greater curvature of the stomach. In both cases, the tumors were R0 and completely resected without complications.

UEMR in duodenal NETs

ER of duodenal NETs (dNETs) is indicated for lesions ≤ 10 mm confined to the submucosal layer, without lymph node involvement or metastases. For intermediate lesions (10–20 mm), ER or surgery may be considered. Surgical resection is recommended for large G1/G2 dNETs (≥ 20 mm) without metastases and for dNETs with lymph node metastases.⁴⁰ The first UEMR for dNETs procedure was described in 2016 by Anderloni *et al.*²⁵ in a case of a 15 mm lesion in the anterior wall of the duodenal bulb, which was classified as a G2 NET. Following this case, other authors have also described case reports of UEMR for dNETs.^{26,27} Kim *et al.*²⁴ also published a series including seven patients with lesions ranging from 3 to 15 mm. NETs were classified as G1, predominantly located in the duodenal bulb (duodenal bulb: five; superior descending angle: one; near the ampulla: one). All lesions were R0 resected without complications. These publications (11 cases in total) reported technical success in all procedures, without adverse events.^{24,26,27} One case of water intoxication was reported after an infusion of approximately 5 L of water during a prolonged UEMR for resection of nonampullary duodenal adenoma.⁴² Although this report was not for NET resection, one should be aware of the risk

of this adverse event after UEMR in the upper digestive tract regardless of the type of lesion.

UEMR in ileum NETs

The distal ileum is the most common site of origin for small bowel NETs.⁴³ Ileal NETs (iNETs) are characterized by a worse prognosis than other sites, with an overall median survival of 88 months (vs. stomach: 124 months; duodenum: 99 months; rectal: 240 months).⁴⁴ Rezende *et al.*²⁸ in 2021 described, for the first and only time, a case of a 10 mm lesion in the distal ileum, classified as a G1 NET and resected by the UEMR procedure. This was an outlier case in a patient who had previously undergone a colectomy with ileorectal anastomosis, and given the difficulties of surgical approach, ER presented as a less invasive treatment. Whenever feasible, surgical resection with lymphadenectomy should be considered the treatment of choice for small bowel NETs.

UEMR in rectal NETs

ER of rectal NETs (rNETs) is indicated for lesions 5–20 mm in size and classified as G1/G2.^{2,29,30,45,46} In this setting, UEMR has recently been introduced as a viable alternative to the EMR technique (Fig. 3). The first case of UEMR in rNETs was described in 2015 by Kawaguti *et al.*¹⁹ in a patient with a 10 mm lesion in the distal rectum, classified as a G1 NET, that was completely resected without complications. One year later, Marui *et al.*²³ reported an *en bloc* resection without complications of an 8 mm diameter carcinoid rectal tumor classified as a G1 NET. Another case report study of six patients with rectal G1 NETs ≤ 10 mm in diameter reported *en bloc* resection and R0 resection in 100% and 83% of cases, respectively, with no adverse events.⁴⁵ In addition, a retrospective study in Brazil evaluated the efficacy and safety of the UEMR technique in 11 patients with rNETs ≤ 12 mm (G1: 9; G2: 2). This study achieved a 100% *en bloc* resection rate and an 81% R0 resection rate with no adverse events.³⁰ The utility of the UEMR technique in rNETs has also been compared with ESD. Shi *et al.*⁵ also compared both techniques in 78 patients (UEMR: 22; ESD: 56) with ≤ 10 mm rectal G1/G2 NETs and found a slightly lower complete resection rate in the UEMR group (UEMR: 95.5% vs. ESD: 96.4%, $p = 0.840$), with one complication in the ESD group and no complications in the UEMR




Fig. 3. UEMR procedure for rectal NET. After switching from the endoscopic view with air (a) to the underwater view (b), the lesion is captured (c) and subsequently presented with that view after resection (d). We declare that the patient has provided consent for the procedure, and it should be noted that the image of the technique is purely illustrative. No patient-identifiable information is presented. NET, neuroendocrine tumor; UEMR, underwater endoscopic mucosal resection.

group. Another study also comparing UEMR with ESD in 115 patients (UEMR: 36; ESD: 79) with rectal G1 NETs ≤ 10 mm in size, found similar R0 resection rates (UEMR: 86.1% vs. ESD: 86.1%, $p = 0.996$) with 2.5% (2/79) of adverse events in the ESD group and no adverse events in UEMR group. The procedure time was significantly longer in the ESD group than in the UEMR group (mean 26.6 ± 13.4 m vs. 5.8 ± 2.9 m, $p < 0.0001$). Although not included in the title or primary objectives of this study, Park *et al.*⁹ also compared UEMR with precut EMR (EMR-P) or cap-assisted EMR (EMR-C). The prevalence of lateral margin positivity was significantly lower in the UEMR group (UEMR vs. EMR-P/EMR-C, 5.6% vs. 20.8%, $p = 0.039$). R0 resection rates were higher in the UEMR group, but the difference did not reach statistical significance (UEMR vs. EMR-P/EMR-C, 86.1% vs. 72.7%, $p = 0.115$). The procedure time was significantly shorter in the UEMR group (UEMR vs. EMR-P/EMR-C, mean 5.8 ± 2.9 m vs. 18.5 ± 11.2 m, $p < 0.001$). Finally, a retrospective study compared UEMR and endoscopic submucosal resection with a ligation device (ESMR-L) in 32 patients (UEMR: 7; ESMR-L: 25) with predominantly G1 NETs < 10 mm. This study achieved 100% *en bloc* resection and R0 resection rates in both groups, with two adverse events in the ESMR-L group and no adverse events in the UEMR group.²⁹ A summary of comparative studies of UEMR with other endoscopic techniques for resection of rNETs is shown in Table 3.^{5,9,29}

Discussion

ER plays an important role in the management of NETs. However,

the best technique for resection remains controversial, and clear recommendations have yet to be made. As most NETs involve the submucosa, conventional hot snare polypectomy or EMR has been reported inadequate for R0 resection and rarely result in free vertical margins.^{2,29,46} ESD has been shown to be a good alternative, achieving R0 resection in 80.6–100% in some studies,⁹ but this method requires greater technical skill, has a longer procedure time, a high risk of adverse events, and is not widely used worldwide.¹¹ Endoscopic full-thickness resection has also been described as an option for endoscopic treatment of rectal NETs. However, it has specific adverse events such as stenosis, adhesion, or damage to extraluminal structures, and the potential to interfere with subsequent surgery if required.⁴⁷ Given these limitations, other methods have been described that attempt to ensure complete histological resection without adding disadvantages to the endoscopic management of these lesions. EMR-C has shown promising results in the removal of submucosal subepithelial lesions, but the size of the cap limits its functionality to the removal of tumors smaller than 10 mm.²⁴ Another technique described is ESMR-L, which is commonly used in Japan for resection of small rectal NETs and is performed using an elastic band to excise the deep submucosal layer. ESMR-L results in better R0 resection rates than EMR and comparable rates to ESD.²⁹

UEMR has some notable strengths. In particular, it has shown positive results, especially for ER of rNETs, with a high rate of technical success and a high rate of R0 resection. The UEMR is also a very safe procedure, with a low incidence of adverse events. And what is most appealing about this technique is its simplicity. In addition, it is inexpensive, easy to master (can be performed by begin-

Table 3. Summary of comparative studies of UEMR with other endoscopic techniques for resection of rNETs

First author, year	Endoscopy type		p-value
Park, 2020 ⁹	UEMR	ESD	
R0 resection	86.1%	86.1%	0.996 ^{a,b}
Complications	0%	2.5%	1.0 ^a
Procedure time in m, mean ± SD	5.8 ± 2.9	26.6 ± 13.4	<0.001
Park, 2020 ⁹	UEMR	EMR	
R0 resection	86.1%	72.7%	0.115 ^a
Complications	0%	0%	–
Procedure time in m, mean ± SD	5.8 ± 2.9	18.5 ± 11.2	<0.001 ^b
Shi, 2022 ⁵	UEMR	ESD	
Complete resection	95.5%	96.4%	0.840 ^c
Complications	0%	1.8%	–
Procedure time in m, median/IQR	5.0 ± 1.4	26.6 ± 8.1	<0.001 ^b
Total cost in USD, mean ± SD	\$328 ± 20.6	\$1,759.0 ± 72.6	<0.001 ^b
Okada, 2022 ²⁹	UEMR	ESMR-L	
R0 resection	100%	100%	1.0 ^a
Complications	0%	8%	1.0 ^a
Procedure time in m (median/IQR)	6/5–8	12/9–14	0.002 ^d
Total cost in USD, median/IQR	\$90.45/83.64–108.41	\$274.73/265.86–292.45	<0.001 ^d

ESD, endoscopic submucosal dissection; ESGM-L, endoscopic submucosal resection with a ligation device; IQR, interquartile range; UEMR, underwater endoscopic mucosal resection. SD, standard deviation. USD, United States dollar. ^aFisher exact test; ^bStudent's *t*-test; ^cchi-square test; ^dMann-Whitney *U*-test.

ners without a learning curve), requires no special accessories, and has a short procedure time. For these reasons, UEMR can even be considered the first-line technique for rNETs resection.^{4,5,9,11,17,30,31}

While UEMR offers some advantages, several technical and safety considerations warrant attention. In the upper gastrointestinal tract, the risk of bronchoaspiration exists, which may be mitigated by using an overtube or endotracheal intubation.⁴⁷ Additionally, the stiffer submucosal layer of the stomach may limit optimal lesion lifting, particularly for tumors not confined to the mucosa.⁴⁸ Although rare, hyponatremia due to fluid absorption remains a potential risk, especially during prolonged procedures; this may be reduced by infusing saline instead of water.⁴² Finally, UEMR—like all endoscopic approaches—plays a very limited role in the management of aggressive neuroendocrine tumors (e.g., esophageal NETs) and should be reserved for select, non-surgical candidates for ileal NETs.^{28,33,36}

This systematic review has several limitations. First, the included studies are predominantly case reports and small case series, with a limited number of comparative trials. Second, the evidence for UEMR in upper GI NETs (esophagus, stomach, duodenum) is scarce, with only a few reported cases. Third, heterogeneity in study designs and outcome reporting precluded a meta-analysis. Nonetheless, this review provides a comprehensive overview of the current evidence and highlights the need for larger, prospective studies to validate the role of UEMR across different GI NET locations.

Conclusions

This systematic review synthesizes the current, albeit limited, evi-

dence on UEMR for GI NETs. The findings suggest that UEMR could be a technically feasible option for well-selected cases, most notably for small (≤ 20 mm), low-grade (G1/G2) rectal NETs, where it appears comparable to conventional EMR or ESD in achieving complete resection, but with potentially shorter procedure times and lower technical demands. For NETs located in the stomach, duodenum, or ileum, the evidence remains preliminary and derived from isolated case reports, precluding definitive recommendations. In these upper GI locations, UEMR should currently be considered only within specialized centers and for patients who are not candidates for surgery, while acknowledging risks such as aspiration. Ultimately, UEMR does not replace but may complement the existing endoscopic armamentarium. Its definitive role in clinical practice, particularly outside the rectum, awaits validation through prospective, comparative studies with standardized protocols and longer follow-up.

Supporting information

Supplementary material for this article is available at <https://doi.org/10.14218/JTG.2023.00031>.

Supplementary Table 1. Dummy caption Dummy caption Dummy caption Dummy caption Dummy caption Dummy caption.

Acknowledgments

None.

Funding

None.

Conflict of interest

The authors PAES, MHGS, IKGSS, FSK, and LL do not have any relevant competing interests to disclose regarding the content of this article. FMF discloses that he has served on advisory boards for Boston Scientific, Olympus, Medtronic, and Cook in the last 3 years.

Author contributions

Contributed to study concept and design (LL), acquisition of the data (PAES, MHGS, and IKGSS), data analysis and drafting of the manuscript (PAES, MHGS, IKGSS, and LL), critical revision of the manuscript (FSK, FMF, LL), and supervision (FMF, LL)

Data sharing statement

No additional data are available.

References

- labichino G, Di Leo M, Arena M, Rubis Passoni GG, Morandi E, Turpini F, *et al*. Diagnosis, treatment, and current concepts in the endoscopic management of gastroenteropancreatic neuroendocrine neoplasms. *World J Gastroenterol* 2022;28(34):4943–4958. doi:10.3748/wjg.v28.i34.4943, PMID:36160644.
- Carvão J, Dinis-Ribeiro M, Pimentel-Nunes P, Libânio D. Neuroendocrine Tumors of the Gastrointestinal Tract: A Focused Review and Practical Approach for Gastroenterologists. *GE Port J Gastroenterol* 2021;28(5):336–348. doi:10.1159/000512089, PMID:34604465.
- Pimentel-Nunes P, Ortigão R, Afonso LP, Bastos RP, Libânio D, Dinis-Ribeiro M. Endoscopic Resection of Gastrointestinal Neuroendocrine Tumors: Long-Term Outcomes and Comparison of Endoscopic Techniques. *GE Port J Gastroenterol* 2022;30(2):98–106. doi:10.1159/000521654, PMID:37008523.
- Lenz L, Maluf-Filho F. Response to Xiu-He *et al*. Nonpedunculated colorectal lesions: underwater or conventional EMR? *Gastrointest Endosc* 2023;97(4):812. doi:10.1016/j.gie.2022.11.025.
- Shi H, Wang C, Wu J, Qin B, Jiang J, Liu N, *et al*. Underwater endoscopic mucosal resection for rectal neuroendocrine tumors (with videos): a single center retrospective study. *BMC Gastroenterol* 2022;22(1):276. doi:10.1186/s12876-022-02350-6, PMID:35655173.
- Yamashina T, Uedo N, Akasaka T, Iwatsubo T, Nakatani Y, Akamatsu T, *et al*. Comparison of Underwater vs Conventional Endoscopic Mucosal Resection of Intermediate-Size Colorectal Polyps. *Gastroenterology* 2019;157(2):451–461.e2. doi:10.1053/j.gastro.2019.04.005, PMID:30981791.
- Binmoeller KF, Weiler F, Shah J, Bhat Y, Kane S. "Underwater" EMR without submucosal injection for large sessile colorectal polyps (with video). *Gastrointest Endosc* 2012;75(5):1086–1091. doi:10.1016/j.gie.2011.12.022, PMID:22365184.
- Lenz L, Martins B, Kawaguti FS, Tellian A, Pennachi CMPS, Sorbello M, *et al*. UNDERWATER ENDOSCOPIC MUCOSAL RESECTION FOR NON-PEDUNCULATED COLORECTAL LESIONS. A PROSPECTIVE SINGLE-ARM STUDY. *Arq Gastroenterol* 2020;57(2):193–197. doi:10.1590/S0004-2803.202000000-37, PMID:32609162.
- Park SS, Han KS, Kim B, Chang Kim B, Hong CW, Sohn DK, *et al*. Comparison of underwater endoscopic mucosal resection and endoscopic submucosal dissection of rectal neuroendocrine tumors (with videos). *Gastrointest Endosc* 2020;91(5):1164–1171.e2. doi:10.1016/j.gie.2019.12.039, PMID:31904380.
- Rodríguez Sánchez J, Alvarez-Gonzalez MA, Pellisé M, Coto-Ugarte D, Uchima H, Aranda-Hernández J, *et al*. Underwater versus conventional EMR of large nonpedunculated colorectal lesions: a multicenter randomized controlled trial. *Gastrointest Endosc* 2023;97(5):941–951.e2. doi:10.1016/j.gie.2022.12.013, PMID:36572129.
- Lenz L, Maluf-Filho F. Response. *Gastrointest Endosc* 2023;97(4):813. doi:10.1016/j.gie.2022.11.026, PMID:36958927.
- Kawamura T, Sakai H, Ogawa T, Sakiyama N, Ueda Y, Shirakawa A, *et al*. Feasibility of Underwater Endoscopic Mucosal Resection for Colorectal Lesions: A Single Center Study in Japan. *Gastroenterology Res* 2018;11(4):274–279. doi:10.14740/gr1021w, PMID:30116426.
- Chien HC, Uedo N, Hsieh PH. Comparison of underwater and conventional endoscopic mucosal resection for removing sessile colorectal polyps: a propensity-score matched cohort study. *Endosc Int Open* 2019;7(11):E1528–E1536. doi:10.1055/a-1007-1578, PMID:31681832.
- Amato A, Radaelli F, Spinzi G. Underwater endoscopic mucosal resection: The third way for en bloc resection of colonic lesions? *United European Gastroenterol J* 2016;4(4):595–598. doi:10.1177/2050640615617635, PMID:27536370.
- Nomura H, Tsuji S, Utsunomiya M, Kawasaki A, Tsuji K, Yoshida N, *et al*. Resection depth and layer of underwater versus conventional endoscopic mucosal resection of intermediate-sized colorectal polyps: A pilot study. *Endosc Int Open* 2022;10(8):E1037–E1044. doi:10.1055/a-1864-6452, PMID:35979030.
- Uedo N, Nemeth A, Johansson GW, Toth E, Thorlacius H. Underwater endoscopic mucosal resection of large colorectal lesions. *Endoscopy* 2015;47(2):172–174. doi:10.1055/s-0034-1390749, PMID:25314326.
- Lenz L, Martins B, Andrade de Paulo G, Kawaguti FS, Baba ER, Uemura RS, *et al*. Underwater versus conventional EMR for nonpedunculated colorectal lesions: a randomized clinical trial. *Gastrointest Endosc* 2023;97(3):549–558. doi:10.1016/j.gie.2022.10.033, PMID:36309072.
- Spadaccini M, Fuccio L, Lamonaca L, Frazzoni L, Maselli R, Di Leo M, *et al*. Underwater EMR for colorectal lesions: a systematic review with meta-analysis (with video). *Gastrointest Endosc* 2019;89(6):1109–1116.e4. doi:10.1016/j.gie.2018.10.023, PMID:30862352.
- Kawaguti FS, de Oliveira JF, da Costa Martins B, Sorbello MP, Retes FA, Ribeiro U, *et al*. Underwater endoscopic resection of a neuroendocrine rectal tumor. *Endoscopy* 2015;47 Suppl 1 UCTN:E513–E514. doi:10.1055/s-0034-1393224, PMID:26492301.
- Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, *et al*. The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *Int J Surg* 2021;88(March). doi:10.31222/osc.io/v7gm2.
- Moola S, Munn Z, Tufanaru C, Aromataris E, Sears K, Sfetcu R, *et al*. Chapter 7: Systematic reviews of etiology and risk. In: Aromataris E, Munn Z (eds). *JBIM Manual for Evidence Synthesis*. JBI; 2020. doi:10.46658/JBIMES-20-08.
- Munn Z, Barker TH, Moola S, Tufanaru C, Stern C, McArthur A, *et al*. Methodological quality of case series studies: an introduction to the JBI critical appraisal tool. *JBI Evid Synth* 2020;18(10):2127–2133. doi:10.11124/JBISRR-D-19-00099, PMID:33038125.
- Marui S, Yamashina T, Osaki Y. A New Endoscopic Resection Method for Rectal Carcinoid Tumor: Underwater Endoscopic Mucosal Resection. *Glob J Gastroenterol Hepatol* 2016;4(2):24–5. doi:10.12970/2308-6483.2016.04.02.1.
- Kim SJ, Kim TU, Choi CW, Kim HW, Park SB, Ryu DG. Underwater endoscopic mucosal resection of upper gastrointestinal subepithelial tumors: A case series pilot study (with video). *Medicine (Baltimore)* 2022;101(41):e31072. doi:10.1097/MD.00000000000031072, PMID:36254017.
- Anderloni A, Murino A, Jovani M, Battista S, Repici A. Underwater endoscopic mucosal resection of a duodenal neuroendocrine tumor. *Gastrointest Endosc* 2016;83(1):259–260. doi:10.1016/j.gie.2015.08.001, PMID:26264433.
- Uchima H, Diez-Caballero A, Capdevila J, Rosinach M, Mata A, Turró R, *et al*. Underwater endoscopic mucosal resection for en bloc resection of a neuroendocrine tumor in the duodenal bulb. *Endoscopy* 2022;54(6):E264–E265. doi:10.1055/a-1512-8954, PMID:34144620.
- Lee JW, Kim SJ, Choi CW, Ryu DG. Underwater Endoscopic Mucosal Resections of Non-ampullary Small Duodenal Tumors. *Korean J Helicobacter Up Gastrointest Res* 2022;22(2):152–156. doi:10.7704/kjhgur.2022.0009.

- [28] Rezende DT, Kawaguti FS, Safatle-Ribeiro AV, Tolentino LHL, Ribeiro Junior U, Maluf-Filho F. Underwater endoscopic resection of an ileal neuroendocrine tumor. *Endoscopy* 2021;53(2):E48–E49. doi:10.1055/a-1178-0143, PMID:32503073.
- [29] Okada M, Shinozaki S, Ikeda E, Hayashi Y, Takezawa T, Fukuda H, *et al*. Underwater Endoscopic Mucosal Resection of Small Rectal Neuroendocrine Tumors. *Front Med (Lausanne)* 2022;9:835013. doi:10.3389/fmed.2022.835013, PMID:35514753.
- [30] Coutinho LMA, Lenz L, Kawaguti FS, Martins BC, Baba E, Gusmon C, *et al*. UNDERWATER ENDOSCOPIC MUCOSAL RESECTION FOR SMALL RECTAL NEUROENDOCRINE TUMORS. *Arq Gastroenterol* 2021;58(2):210–213. doi:10.1590/S0004-2803.202100000-37, PMID:34133614.
- [31] Yamashina T, Tumura T, Maruo T, Matsumae T, Yoshida H, Tanke G, *et al*. Underwater endoscopic mucosal resection: a new endoscopic method for resection of rectal neuroendocrine tumor grade 1 (carcinoid) ≤ 10 mm in diameter. *Endosc Int Open* 2018;6(1):E111–E114. doi:10.1055/s-0043-123467, PMID:29344571.
- [32] Awada H, Hajj Ali A, Bakhshwin A, Daw H. High-grade large cell neuroendocrine carcinoma of the esophagus: a case report and review of the literature. *J Med Case Rep* 2023;17(1):144. doi:10.1186/s13256-023-03879-0, PMID:37024958.
- [33] Giannetta E, Guarnotta V, Rota F, de Cicco F, Grillo F, Colao A, *et al*. A rare rarity: Neuroendocrine tumor of the esophagus. *Crit Rev Oncol Hematol* 2019;137:92–107. doi:10.1016/j.critrevonc.2019.02.012, PMID:31014519.
- [34] Tustumi F, Takeda FR, Uema RH, Pereira GL, Sallum RA, Cecconello I. Primary neuroendocrine neoplasm of the esophagus - Report of 14 cases from a single institute and review of the literature. *Arq Gastroenterol* 2017;54(1):4–10. doi:10.1590/S0004-2803.2017v54n1-01, PMID:28079231.
- [35] Ma Z, Cai H, Cui Y. Progress in the treatment of esophageal neuroendocrine carcinoma. *Tumour Biol* 2017;39(6):1010428317711313. doi:10.1177/1010428317711313, PMID:28653897.
- [36] Cai MY, Cardoso MF, Zhu Y, Zhong YS, Zhou PH. An esophageal submucosal tumor treated with submucosal tunneling endoscopic resection: an unexpected result. *Gastroenterol Rep (Oxf)* 2021;9(3):269–271. doi:10.1093/gastro/goaa073, PMID:34316377.
- [37] Doumbe-Mandengue P, Geyl S, Guyot A, Pioche M, Rodrigues R, Albouys J, *et al*. Underwater endoscopic mucosal resection for a protruding lesion in the esophagus. *Endoscopy* 2022;54(S 02):E869–E870. doi:10.1055/a-1847-7235, PMID:35750074.
- [38] Hwang CS, Kim SJ, Choi CW. Underwater endoscopic mucosal resection for esophageal granular cell tumor sized more than 1 cm. *Dig Endosc* 2020;32(6):995. doi:10.1111/den.13764, PMID:32512623.
- [39] Akasaka T, Takeuchi Y, Uedo N, Ishihara R, Iishi H. "Underwater" endoscopic submucosal dissection for superficial esophageal neoplasms. *Gastrointest Endosc* 2017;85(1):251–252. doi:10.1016/j.gie.2016.07.018, PMID:27449194.
- [40] Delle Fave G, O'Toole D, Sundin A, Taal B, Ferolla P, Ramage JK, *et al*. ENETS Consensus Guidelines Update for Gastroduodenal Neuroendocrine Neoplasms. *Neuroendocrinology* 2016;103(2):119–124. doi:10.1159/000443168, PMID:26784901.
- [41] Kunz PL, Reidy-Lagunes D, Anthony LB, Bertino EM, Brendtro K, Chan JA, *et al*. Consensus guidelines for the management and treatment of neuroendocrine tumors. *Pancreas* 2013;42(4):557–577. doi:10.1097/MPA.0b013e31828e34a4, PMID:23591432.
- [42] Binmoeller KF, Shah JN, Bhat YM, Kane SD. "Underwater" EMR of sporadic laterally spreading nonampullary duodenal adenomas (with video). *Gastrointest Endosc* 2013;78(3):496–502. doi:10.1016/j.gie.2013.03.1330, PMID:23642790.
- [43] Wang R, Zheng-Pywell R, Chen HA, Bibb JA, Chen H, Rose JB. Management of Gastrointestinal Neuroendocrine Tumors. *Clin Med Insights Endocrinol Diabetes* 2019;12:1179551419884058. doi:10.1177/1179551419884058, PMID:31695546.
- [44] Yao JC, Hassan M, Phan A, Dagohoy C, Leary C, Mares JE, *et al*. One hundred years after "carcinoid": epidemiology of and prognostic factors for neuroendocrine tumors in 35,825 cases in the United States. *J Clin Oncol* 2008;26(18):3063–3072. doi:10.1200/JCO.2007.15.4377, PMID:18565894.
- [45] Sekiguchi M, Sekine S, Sakamoto T, Otake Y, Nakajima T, Matsuda T, *et al*. Excellent prognosis following endoscopic resection of patients with rectal neuroendocrine tumors despite the frequent presence of lymphovascular invasion. *J Gastroenterol* 2015;50(12):1184–1189. doi:10.1007/s00535-015-1079-7, PMID:25936647.
- [46] Fine C, Roquin G, Terrebbonne E, Lecomte T, Coriat R, Do Cao C, *et al*. Endoscopic management of 345 small rectal neuroendocrine tumours: A national study from the French group of endocrine tumours (GTE). *United European Gastroenterol J* 2019;7(8):1102–1112. doi:10.1177/2050640619861883, PMID:31662867.
- [47] Deprez PH, Moons LMG, O'Toole D, Gincul R, Seicean A, Pimentel-Nunes P, *et al*. Endoscopic management of subepithelial lesions including neuroendocrine neoplasms: European Society of Gastrointestinal Endoscopy (ESGE) Guideline. *Endoscopy* 2022;54(4):412–429. doi:10.1055/a-1751-5742, PMID:35180797.
- [48] Iwagami H, Kanesaka T, Ishihara R, Uedo N. Underwater endoscopic mucosal resection for remaining early gastric cancer after endoscopic submucosal dissection. *Endoscopy* 2019;51(8):E229–E230. doi:10.1055/a-0875-3429, PMID:30995684.