



Guideline

Chinese Guidelines for the Diagnosis and Treatment of Autoimmune Pancreatitis (Shanghai, 2023)



Lei Xin¹, Yili Cai², Lianghao Hu¹, Hongyu Li³, Dong Wu⁴, Zhaoshen Li^{1*} , Xun Li^{5*} , Xiaozhong Guo^{3*} , Zhuan Liao^{1*} and on behalf of the National Clinical Research Center for Digestive Diseases (Shanghai); National Key Laboratory of Immunity and Inflammation; Professional Committee of Pancreatic Disease, Chinese Medical Doctor Association; Pancreas Study Group, Chinese Society of Gastroenterology, Chinese Medical Association; Editorial Board of *Chinese Journal of Pancreatology*

¹Department of Gastroenterology, The First Affiliated Hospital of Naval Medical University, Shanghai, China; ²Department of Gastroenterology, Naval Medical Center, Naval Medical University, Shanghai, China; ³Department of Gastroenterology, General Hospital of Northern Theater Command, Shenyang, Liaoning, China; ⁴Department of Gastroenterology, Peking Union Medical College Hospital, Beijing, China; ⁵Department of General Surgery, The First Hospital of Lanzhou University, Lanzhou, Gansu, China

Received: April 14, 2026 | Revised: May 19, 2026 | Accepted: May 25, 2026 | Published online: June 23, 2026

Abstract

Autoimmune pancreatitis (AIP) is a rare immune-mediated form of chronic pancreatitis. It may affect multiple organs, and its heterogeneous clinical manifestations complicate diagnosis and management. Based on the *Chinese Guidelines for the Diagnosis and Management of Autoimmune Pancreatitis (Shanghai 2012 Draft)*, together with the latest domestic and international guidelines and research advances, the present guideline provides 20 recommendations covering four aspects: diagnosis, treatment, follow-up, and prognosis. The aim is to improve the diagnosis and management of AIP in China and ultimately improve patient outcomes.

Introduction

Autoimmune pancreatitis (AIP) is a distinct form of chronic pancreatitis that typically presents with obstructive jaundice, abdominal discomfort, and other clinical manifestations. It is immune-mediated and marked by lymphocytic and plasmacytic infiltration, pancreatic fibrosis, pancreatic dysfunction, and possible extrapancreatic organ involvement, and it generally responds well to glucocorticoid therapy. In 2012, the Chinese Journal of Pancreatology organized the development of the *Chinese Guidelines*

for the Diagnosis and Management of Autoimmune Pancreatitis (Shanghai, 2012 Draft). After 10 years of application in clinical practice, these guidelines have played an important role in improving the diagnosis and management of AIP in China. In recent years, substantial progress has been made in AIP research both in China and internationally, with a deeper understanding of the disease and increasingly standardized clinical practice, although some issues remain controversial. To better guide the clinical diagnosis and management of AIP, the National Clinical Research Center for Digestive Diseases (Shanghai), the National Key Laboratory of Immunity and Inflammation, the Professional Committee of Pancreatic Disease of the Chinese Medical Doctor Association, the Pancreas Study Group of the Chinese Society of Gastroenterology of the Chinese Medical Association, and the Editorial Board of the Chinese Journal of Pancreatology jointly established a multidisciplinary working group. This group included experts in gastroenterology, rheumatology, surgery, radiology, pathology, laboratory medicine, and evidence-based medicine, and revised the previous guidelines to form the present document. The present guidelines were registered on the Practice Guideline Registration for Transparency platform (PREPARE-2022CN812). The design and development followed the WHO Handbook for Guideline Development issued by the World Health Organization in 2014 and the Chinese guiding principles for the development/revision

Keywords: Autoimmune pancreatitis; IgG4-related disease; IgG4-related cholangitis; Guidelines; Glucocorticoid therapy; Multidisciplinary management.

***Correspondence to:** Zhaoshen Li, Department of Gastroenterology, The First Affiliated Hospital of Naval Medical University, Shanghai 200433, China. ORCID: <https://orcid.org/0000-0003-0404-242X>. Tel: +86-021-81873270, E-mail: zhsl@vip.163.com; Xun Li, Department of General Surgery, The First Hospital of Lanzhou University, Lanzhou, Gansu 730000, China. ORCID: <https://orcid.org/0000-0002-1452-613X>. Tel: +86-0931-8912169, E-mail: lxdr21@126.com; Xiaozhong Guo, Department of Gastroenterology, General Hospital of Northern Theater Command, Shenyang, Liaoning 110840, China. ORCID: <https://orcid.org/0000-0002-6397-0501>. Tel: +86-024-28897603, E-mail: guoxiaozhong1962@163.com; Zhuan Liao, Department of Gastroenterology, The First Affiliated Hospital of Naval Medical University, Shanghai 200433, China. ORCID: <https://orcid.org/0000-0001-8506-8159>. Tel: +86-021-31161012, E-mail: zhuanleo@126.com

How to cite this article: Xin L, Cai Y, Hu L, Li H, Wu D, Li Z, *et al.* Chinese Guidelines for the Diagnosis and Treatment of Autoimmune Pancreatitis (Shanghai, 2023). *J Transl Gastroenterol* 2026;4(2):98–111.

of clinical practice guidelines issued by the Chinese Medical Association in 2022,¹ with reference to the Appraisal of Guidelines for Research & Evaluation II and the Reporting Items for Practice Guidelines in Healthcare Statement.^{2,3} Through a systematic review of the published literature and interviews with selected experts, 20 recommendations were formulated regarding diagnosis, treatment, follow-up, and prognosis. After evidence retrieval and methodological quality assessment, the GRADE approach (Grading of Recommendations Assessment, Development and Evaluation) was used to rate the quality of evidence (high, moderate, low, or very low) and the strength of recommendations (strong or weak).⁴ Recommendations are presented as declarative statements (Table 1). Three rounds of Delphi surveys were conducted, and after further refinement based on expert feedback, consensus was reached for all recommendations (all consensus rates > 96.7%).

Diagnosis

I. Clinical manifestations

Recommendation 1. Patients with AIP commonly present with non-specific symptoms such as obstructive jaundice and abdominal discomfort, and some patients are asymptomatic. Clinical manifestations of extrapancreatic organ involvement should be carefully assessed. (Quality of evidence: High; Strength of recommendation: Strong; Level of consensus: 100%)

The mean age at diagnosis of type 1 AIP is approximately 60 years, and the incidence in men is about three times that in women.⁵ The clinical manifestations of type 1 AIP can be divided into pancreatic and extrapancreatic manifestations. Pancreatic manifestations are non-specific, and common symptoms include obstructive jaundice (51.9–75.0%), abdominal discomfort or abdominal pain (41.0–65.4%), and weight loss (40.0–45.1%). Some patients present with polydipsia, polyuria, or fatigue, whereas others have no clinical manifestations and seek medical attention only after pancreatic enlargement is detected during a health examination.^{6–10} These clinical manifestations resemble those of pancreatic cancer and may therefore lead to misdiagnosis.

A substantial proportion of patients with type 1 AIP have extrapancreatic manifestations, which essentially represent organ involvement in IgG4-related disease (IgG4-RD) affecting different organs. Such involvement does not necessarily parallel the severity of pancreatic lesions and may occur before, concurrently with, or after pancreatic manifestations. Some patients have corresponding physical signs, which should be carefully assessed on examination. Common extrapancreatic organ involvement includes sclerosing cholangitis, Mikulicz disease (dacryoadenitis and sialadenitis), retroperitoneal fibrosis, interstitial nephritis, superficial and deep lymphadenopathy (hilar, mediastinal, and abdominal), and interstitial pneumonia.¹¹ One study including 100 Chinese patients with AIP found that 77 patients (77.0%) had 81 extrapancreatic lesions,¹² with biliary lesions being the most common (64 lesions), including 42 strictures of the distal common bile duct and 22 strictures of the common hepatic duct or intrahepatic bile ducts. In addition, approximately 51.5% of patients with type 1 AIP had peripancreatic vascular involvement.¹³

The mean age at diagnosis of type 2 AIP is lower than that of type 1 AIP. One study reported a mean age of 43 years, and the disease generally occurs between 30 and 50 years of age, with no sta-

tistically significant sex difference in incidence.^{14,15} Patients with type 2 AIP often present with abdominal pain resembling acute pancreatitis.¹⁶ In addition, compared with type 1 AIP, type 2 AIP is more likely to be associated with inflammatory bowel disease (IBD).¹⁷ In recent years, increasing attention has been paid to the relationship between AIP and IBD, and 6–27% of patients with AIP are reported to have concomitant IBD.^{18,19}

II. Imaging examinations

Recommendation 2. Contrast-enhanced computed tomography (CT) is recommended as the first-line imaging modality for the diagnosis of AIP. (Quality of evidence: Moderate; Strength of recommendation: Strong; Level of consensus: 100%)

On contrast-enhanced CT, AIP commonly presents as diffuse enlargement of the pancreas. A hypoenhancing capsule-like rim may be seen during the pancreatic parenchymal phase, producing a “sausage-like” appearance; punctate enhancement may be present within the pancreatic parenchyma, followed by delayed enhancement on later-phase images. AIP may also present as a focal pancreatic mass, and a small number of cases show multifocal lesions.^{20,21} The International Consensus Diagnostic Criteria (ICDC) for AIP classify pancreatic parenchymal CT findings into two levels: typical and indeterminate. Typical findings include diffuse pancreatic enlargement with delayed enhancement, whereas indeterminate findings include focal or multifocal enlargement with delayed enhancement.¹⁶ A capsule-like rim around the pancreas is uncommon in other diseases and is therefore a relatively specific imaging sign of AIP. Delayed enhancement is also characteristic of AIP, although it is less common in cases with less marked fibrosis.²²

CT can differentiate AIP from chronic pancreatitis. The features of typical chronic pancreatitis include pancreatic atrophy, parenchymal or intraductal calcification, irregular ductal dilatation with side-branch ectasia, and pseudocyst formation. However, recent studies have shown that mild pancreatic ductal dilatation is not rare in AIP. Approximately 10% of patients with a longer disease course may develop pancreatic atrophy, pancreatic duct stones, and related changes,²³ and cases complicated by pseudocysts have also been reported.²⁴

CT is of major value in differentiating AIP from pancreatic cancer. A meta-analysis showed that the sensitivity and specificity of contrast-enhanced CT for distinguishing AIP from pancreatic cancer were 59% (95% confidence interval (CI), 0.410–0.750) and 99% (95% CI, 0.880–1.000), respectively.²⁵ The typical CT appearance of AIP is diffuse pancreatic enlargement, whereas pancreatic cancer typically appears as a hypovascular focal mass. It should be noted that pancreatic enlargement alone cannot accurately distinguish AIP from pancreatic cancer or other malignant tumors. In a study of 245 patients with diffuse pancreatic enlargement on contrast-enhanced CT, only 54.7% had AIP; other malignant lesions included pancreatic ductal adenocarcinoma (35.9%), pancreatic neuroendocrine tumor (4.5%), lymphoma (1.6%), metastasis (1.6%), solid pseudopapillary tumor (1.2%), and acinar cell carcinoma (0.4%). The presence of residual normal pancreatic parenchyma, a longer short-axis diameter (cutoff, 3.15 cm), and a lower delayed-phase lesion-to-aorta enhancement ratio (cutoff, 0.75) suggested a higher likelihood of malignancy.²¹ In practice, multiple imaging signs should be integrated when differentiating these two diseases. CT features suggestive

Table 1. Summary of recommendations, evidence quality, recommendation strength, and consensus level

No.	Recommendation	Evidence/Strength	Consensus
1	Patients with autoimmune pancreatitis (AIP) commonly present with non-specific symptoms such as obstructive jaundice and abdominal discomfort, and some patients are asymptomatic. Clinical manifestations of extrapancreatic organ involvement should be carefully assessed	High/Strong	100%
2	Contrast-enhanced computed tomography (CT) is recommended as the first-line imaging modality for the diagnosis of AIP	Moderate/Strong	100%
3	Contrast-enhanced magnetic resonance imaging (MRI) is suggested as an important imaging modality for the diagnosis of AIP. Magnetic resonance cholangiopancreatography is suggested for the assessment of pancreatic ductal and biliary changes in AIP	Low/Weak	100%
4	In patients with indeterminate CT or MRI findings, positron emission tomography-CT may be used as an adjunct for the diagnosis and differential diagnosis of AIP	Low/Weak	100%
5	Endoscopic ultrasound (EUS) can identify characteristic changes in the pancreatic parenchyma and the pancreaticobiliary ducts in AIP, and can also provide histologic or cytologic specimens for diagnosis and differential diagnosis. It is therefore recommended as an important diagnostic modality for AIP	Moderate/Strong	100%
6	Endoscopic retrograde cholangiopancreatography can identify characteristic pancreaticobiliary ductal changes in AIP; however, it is not recommended as a routine diagnostic modality because of its invasiveness	Low/Weak	100%
7	Serum IgG4 is suggested as the preferred laboratory test for the diagnosis of type 1 AIP, assessment of treatment response, and monitoring of disease activity	Low/Weak	96.7%
8	Characteristic pathological changes are important diagnostic evidence for AIP. Tissue acquisition and pathologic diagnosis should therefore be emphasized, and EUS-guided fine-needle biopsy is recommended as the preferred method	Moderate/Weak	100%
9	The diagnosis of AIP is recommended to be based on an integrated assessment of symptoms, signs, imaging, EUS findings, serum IgG4, pathological findings, and extrapancreatic organ involvement, with reference to the response to glucocorticoid therapy	Moderate/Strong	100%
10	In patients with suspected AIP, differential diagnosis from pancreatic cancer and other diseases should be emphasized	High/Strong	100%
11	In patients with established type 1 AIP, dynamic assessment of disease activity by imaging and serum IgG4 during the disease course is suggested	Low/Weak	96.7%
12	Induction therapy is suggested for patients with AIP who have symptoms or laboratory/imaging evidence of active disease	Moderate/Weak	100%
13	Oral glucocorticoids are recommended as the first-line induction therapy for AIP. Prednisone 30–40 mg/day, or approximately 0.6 mg·kg ⁻¹ ·day ⁻¹ , may be used initially and tapered gradually after remission is achieved	Moderate/Strong	100%
14	In patients with diffuse pancreatic enlargement, slow radiologic remission after treatment, or serum IgG4 levels remaining more than twice the upper limit of normal, low-dose glucocorticoid maintenance therapy is recommended to reduce the risk of relapse	High/Strong	96.7%
15	Re-treatment with glucocorticoids is suggested for patients with relapsed AIP	Low/Weak	100%
16	In patients with AIP who are refractory to glucocorticoids or experience repeated relapses during tapering or after withdrawal, combination treatment with immunosuppressive agents or switching to biologic therapy is suggested	Moderate/Weak	100%
17	Pancreatic enzyme replacement therapy is suggested for patients with AIP who have pancreatic exocrine insufficiency	Low/Weak	100%
18	Routine biliary drainage is not recommended for patients with AIP complicated by obstructive jaundice	Moderate/Strong	100%
19	Because AIP may involve multiple organs, strengthening multidisciplinary collaboration is suggested during diagnosis and management	Low/Weak	100%
20	Long-term follow-up should be emphasized in patients with AIP, with attention to the risks of relapse, pancreatic endocrine and exocrine insufficiency, pancreatic duct stones, and malignancy	Low/Weak	100%

of pancreatic cancer include absence of diffuse pancreatic enlargement, a hypovascular mass, absence of renal involvement, invasion of the pancreatic or bile ducts, invasion of adjacent vessels, and enlarged lymph nodes.²⁶ In a radiomics study including 42 cases of focal AIP and 334 cases of pancreatic cancer, the radiomics model yielded a sensitivity of 95.24%, specificity of 92.73%, and accuracy of 94.00% for the diagnosis of focal AIP.²⁷ Similarly, the use of relative CT values of the adjacent pancreatic parenchyma has also shown high feasibility for differentiating focal AIP from pancreatic cancer.²⁸

Recommendation 3. Contrast-enhanced magnetic resonance imaging (MRI) is suggested as an important imaging modality for the diagnosis of AIP. Magnetic resonance cholangiopancreatography (MRCP) is suggested for the assessment of pancreatic ductal and biliary changes in AIP. (Quality of evidence: Low; Strength of recommendation: Weak; Level of consensus: 100%)

Typical MRI findings of AIP include diffuse pancreatic enlargement or a focal mass accompanied by abnormal signal intensity, usually with slightly low signal on T1-weighted imaging and slightly high signal on T2-weighted imaging; a capsule-like rim around the pancreas may be seen on both T1- and T2-weighted images.^{29,30} Compared with contrast-enhanced CT, MRI provides better soft-tissue contrast, although its cost and technical requirements are higher; therefore, it may serve as an adjunct to CT. The main MRI features used to distinguish AIP from pancreatic cancer include multiple strictures of the main pancreatic duct, absence of marked upstream main pancreatic duct dilatation, a capsule-like rim around the pancreas, and the duct-penetrating sign. Among these, the absence of significant upstream ductal dilatation has the highest sensitivity, whereas the capsule-like rim has the highest specificity.³¹ In addition, MR elastography may be useful for differentiating AIP from pancreatic cancer and holds promise as a quantitative imaging modality for both differential diagnosis and treatment monitoring.³²

Typical MRCP findings in AIP are long-segment and multiple pancreatic duct strictures. The ICDC for AIP also classifies main pancreatic duct abnormalities as typical or indeterminate.¹⁶ Typical findings include long strictures involving more than one-third of the main pancreatic duct or multiple strictures without significant upstream dilatation, whereas indeterminate findings include focal or multifocal strictures without significant upstream dilatation (pancreatic duct diameter < 5 mm). After glucocorticoid therapy, the pancreatic duct diameter may return to normal; therefore, MRCP may also be used to assess treatment response in AIP.³³ With continued progress in recent years, MRCP has gradually become the preferred imaging modality for evaluating the pancreatic duct.³⁴

MRI and MRCP are also useful for differentiating AIP from ordinary chronic pancreatitis. On MRI, ordinary chronic pancreatitis typically presents with pancreatic atrophy or irregular pancreatic contours, heterogeneous parenchymal signal intensity, parenchymal or intraductal stones appearing as signal voids, pseudocysts, and irregular dilatation of the main pancreatic duct with side-branch ectasia. On MRCP, long-segment or multiple strictures of the main pancreatic duct without significant upstream ductal dilatation are indicative of AIP, whereas irregular ductal dilatation, ductal obstruction by stones, side-branch ectasia, and pseudocyst formation are more suggestive of ordinary chronic pancreatitis.

Recommendation 4. In patients with indeterminate CT or MRI findings, positron emission tomography-computed tomography (PET-CT) may be used as an adjunct for the diagnosis and differential diagnosis of AIP. (Quality of evidence: Low; Strength of recommendation: Weak; Level of consensus: 100%)

Typical PET-CT findings in AIP include diffusely increased fluorodeoxyglucose (FDG) uptake in the pancreas with heterogeneous intensity. Similar uptake patterns may also be seen in involved extrapancreatic organs,^{35,36} which can facilitate differentiation from pancreatic cancer.³⁷ A study comparing PET-CT findings between AIP and pancreatic cancer found that a maximum standardized uptake value (SUVmax) >7.08 and focal or multifocal homogeneous FDG uptake were independent predictors of pancreatic cancer ($P < 0.05$).³⁸ FDG uptake decreases after glucocorticoid therapy in patients with AIP; therefore, PET-CT may also be used to evaluate treatment response, although its cost should be taken into account. In addition, dual-time-point PET-CT and quantitative radiomics prediction models may also be helpful in differentiating AIP from pancreatic cancer.^{39,40}

III. Digestive endoscopy

Recommendation 5. Endoscopic ultrasound (EUS) can identify characteristic changes in the pancreatic parenchyma and pancreaticobiliary ducts in AIP and can also provide histologic or cytologic specimens for diagnosis and differential diagnosis. It is therefore recommended as an important diagnostic modality for AIP. (Quality of evidence: Moderate; Strength of recommendation: Strong; Level of consensus: 100%)

Typical EUS findings in AIP include diffuse pancreatic enlargement or a localized mass, with predominantly hypoechoic pancreatic parenchyma containing internal hyperechoic foci (linear, patchy, or heterogeneous changes), wavy margins, and irregular narrowing of the main pancreatic duct accompanied by ductal wall thickening.^{41,42} In patients with biliary involvement, thickening and hypoechogenicity of the bile duct wall may be seen. The outer and inner layers of the bile duct wall may show hyperechoic bands, resulting in a “sandwich-like” or “onion-skin-like” appearance, whereas cholangiocarcinoma usually appears as a heterogeneously hyperechoic mass.⁴³

Enhanced EUS imaging techniques may facilitate the diagnosis of AIP. On contrast-enhanced EUS, AIP shows a hypervascular pattern relative to the surrounding normal pancreatic parenchyma, which differs from the hypovascular appearance of pancreatic cancer; in addition, arborizing vessels are characteristic of AIP.^{44,45} On contrast-enhanced harmonic EUS, AIP shows iso-enhancement with a homogeneous echo distribution of the contrast agent, whereas pancreatic cancer shows hypo-enhancement with heterogeneous echo distribution.⁴⁶ However, studies of these techniques have generally included small sample sizes and non-randomized designs, and further research is needed to confirm the diagnostic value of EUS.

Studies of artificial intelligence-assisted EUS diagnosis may further improve diagnostic accuracy for AIP. In one study, EUS images from 81 patients with AIP and 100 patients with chronic pancreatitis were analyzed, and 14 of 115 textural parameters

in the region of interest were selected for discrimination, yielding a sensitivity of 84.1%, specificity of 92.5%, and accuracy of 89.3%.⁴⁷ Another study based on a convolutional neural network model using both static EUS images and dynamic videos reported a sensitivity of 94% and specificity of 71% for differentiating AIP from chronic pancreatitis, and a sensitivity of 90% and specificity of 93% for differentiating AIP from pancreatic cancer.⁴⁸

In patients whose diagnosis of AIP cannot be established by typical imaging and laboratory findings, EUS-guided acquisition of pancreatic histologic or cytologic specimens may be used either to diagnose AIP or to exclude malignancy. It was previously thought that specimens obtained by EUS-guided fine-needle aspiration (EUS-FNA) were limited in quantity and had relatively low sensitivity but high specificity for the diagnosis of AIP and were therefore mainly used for differential diagnosis from pancreatic cancer. In recent years, with improvements in needle design, increasing numbers of studies have reported histologic diagnosis of AIP using EUS-FNA or EUS-guided fine-needle biopsy (EUS-FNB). A Chinese study including 27 patients with AIP who underwent EUS-FNA using a 22G needle found that samples from 63% of patients met the histologic criteria proposed in the ICDC for AIP (grade 1 in 5 cases and grade 2 in 12 cases).⁴⁹ A meta-analysis including nine studies of EUS-FNA in 309 patients with AIP and seven studies of EUS-FNB in 131 patients with AIP,⁵⁰ using grade 1 and grade 2 histologic diagnostic evidence as endpoints, showed that the pooled diagnostic yield was 55.8% (95% CI 0.370–0.739) for FNA and 87.2% (95% CI 0.688–0.981) for FNB, with a statistically significant difference ($P = 0.030$). The histologic acquisition rate and adverse event rate were similar between FNA and FNB, and 19G needles had a higher diagnostic yield than 22G needles. For the use of EUS-FNA/FNB-obtained tissue in the diagnosis of AIP, the Japanese guidelines for the histological diagnosis of AIP may be referenced.⁵¹

Recommendation 6. Endoscopic retrograde cholangiopancreatography (ERCP) can identify characteristic pancreaticobiliary ductal changes in AIP; however, it is not recommended as a routine diagnostic modality because of its invasiveness. (Quality of evidence: Low; Strength of recommendation: Weak; Level of consensus: 100%)

Typical ERCP findings of AIP include (i) long ($>1/3$ the length of the pancreatic duct) stricture; (ii) lack of upstream dilatation from the stricture (<5 mm); (iii) multiple strictures; and (iv) side branches arising from a strictured segment.⁵² In contrast, pancreatic cancer more commonly presents with focal ductal stenosis and upstream dilatation.

In some patients with AIP, ERCP cholangiography may show bile duct stricture or features of sclerosing cholangitis. Brush cytology and bile cytology at the site of biliary stricture can aid differential diagnosis.⁵³ Intraductal ultrasonography is useful for differentiating IgG4-related sclerosing cholangitis (IgG4-SC), primary sclerosing cholangitis (PSC), and cholangiocarcinoma. Typical intraductal ultrasonography findings in IgG4-SC include preservation of the three-layer structure of the bile duct wall and uniform, symmetric wall thickening, which may also be seen in non-stenotic segments. PSC often shows destruction of the three-layer bile duct wall structure, asymmetric wall thickening, irregular inner margins, interruption of the outer margin, and diverticulum-like outpouchings. Cholangiocarcinoma typically presents as asymmetric wall thickening or intraluminal protrusion, with irregular inner margins

and interruption of the outer margin, without wall thickening in non-stenotic segments.⁹ With the development of peroral cholangioscopy-assisted ERCP techniques, biopsy performed under direct vision can diagnose IgG4-SC,⁵⁴ although no reports are yet available on pancreatoscopy-guided biopsy via ERCP for the diagnosis of AIP.

In some patients with AIP, the duodenal papilla may also be involved; therefore, concurrent biopsy during ERCP may provide additional diagnostic evidence.^{55,56}

The ICDC for AIP and the Japanese Clinical Diagnostic Criteria (2011) for AIP regarded pancreatography as important diagnostic evidence for AIP, whereas MRCP was not included among the diagnostic criteria. With advances in MRCP, the 2018 Japanese diagnostic criteria for AIP placed MRCP alongside ERCP as the preferred imaging modality for assessing the pancreatic duct.³⁴ Considering the invasiveness and medical costs of ERCP, the present guidelines do not recommend its use as a routine diagnostic method for AIP.

IV. Laboratory tests

Recommendation 7. Serum IgG4 is suggested as the preferred laboratory test for the diagnosis of type 1 AIP, assessment of treatment response, and monitoring of disease activity. (Quality of evidence: Low; Strength of recommendation: Weak; Level of consensus: 96.7%)

Serum IgG4 levels are elevated in 60–90% of patients with type 1 AIP but are also elevated in 7–10% of patients with pancreatic cancer. Using an IgG4 level greater than twice the upper limit of normal as a diagnostic criterion can significantly improve specificity.^{57,58} Some patients with AIP also have hypergammaglobulinemia, elevated IgG and IgE, increased erythrocyte sedimentation rate, elevated C-reactive protein, or positive autoantibodies such as antinuclear antibodies and rheumatoid factor; however, these markers are only suggestive.⁵⁹ In type 2 AIP, serum IgG4 is generally not elevated and autoantibodies are usually negative.

A meta-analysis including 11 studies, 594 patients with AIP, and 958 patients with pancreatic cancer found that elevated serum IgG4 had a sensitivity of 72% (95% CI, 0.680–0.750) and a specificity of 93% (95% CI, 0.920–0.950) for distinguishing AIP from pancreatic cancer.⁶⁰ When serum IgG4 levels greater than twice the upper limit of normal were used to differentiate AIP from pancreatic cancer, the sensitivity was 43% (95% CI, 0.380–0.490) and the specificity was 98% (95% CI, 0.970–0.990).

Some studies have suggested that a decrease in serum IgG4 after treatment for type 1 AIP reflects control of inflammation, whereas persistent elevation above the normal range after treatment predicts relapse. A systematic review including 10 studies and 850 patients with AIP found that the baseline serum IgG4 level at diagnosis was a risk factor for relapse after glucocorticoid induction therapy ($\beta = -0.001$, $P = 0.009$).⁶¹ In addition, both a high baseline serum IgG4 level and re-elevation of serum IgG4 during follow-up are risk factors for relapse.⁶² A nationwide survey in Japan found no statistically significant difference in serum IgG4 levels at diagnosis between patients who relapsed and those who did not, but the lowest post-treatment IgG4 level was higher in patients who relapsed.¹¹

V. Pathological findings

Recommendation 8. Characteristic pathological changes are important diagnostic evidence for AIP. Tissue acquisition

and pathologic diagnosis should therefore be emphasized, and EUS-FNB is recommended as the preferred method. (Quality of evidence: Moderate; Strength of recommendation: Weak; Level of consensus: 100%)

Both type 1 and type 2 AIP show lymphocytic and plasmacytic infiltration around pancreatic ducts and inflammatory changes in the pancreatic stroma, whereas the pathological features of ordinary chronic pancreatitis, such as intraductal protein plugs, stones, and calcification, are less common.

Typical pathological findings of type 1 AIP include dense lymphoplasmacytic infiltration around pancreatic ducts, storiform fibrosis of the parenchyma, obliterative phlebitis, and abundant IgG4-positive plasma cells. Inflammatory cell infiltration within the pancreatic lobules is often prominent, frequently causing acinar atrophy, and fibrosis in the interlobular septa or around the pancreas is more obvious.⁶³ The ICDC statement on the pathology of IgG4-RD recommends that, for diagnosing pancreatic lesions of IgG4-RD, biopsy specimens should contain >10 IgG4-positive plasma cells per high-power field (40×, 0.2 mm²), and surgical resection specimens should contain >50 IgG4-positive plasma cells per high-power field. An IgG4-positive/IgG-positive plasma cell ratio >40% is also an important diagnostic criterion for IgG4-RD. However, it should be noted that the number of IgG4-positive plasma cells or the IgG4/IgG ratio alone cannot directly establish the diagnosis of AIP, because both pancreatic ductal adenocarcinoma tissue and peritumoral tissue may also show varying degrees of IgG4-positive plasma cell infiltration.⁶⁴

Typical pathological findings of type 2 AIP include marked neutrophilic infiltration within the lumen and epithelium of interlobular ducts, resulting in granulocytic epithelial lesions, whereas storiform fibrosis, obliterative phlebitis, and IgG4-positive plasma cell infiltration are uncommon.⁶⁵

As a minimally invasive diagnostic method, EUS-FNB has a clearly higher diagnostic yield for AIP than EUS-FNA.⁵⁰ In patients with suspected AIP based on imaging and laboratory findings, EUS-FNB should be considered first for obtaining histopathologic specimens to confirm the diagnosis.

VI. Diagnosis and differential diagnosis

Recommendation 9. The diagnosis of AIP is recommended to be based on an integrated assessment of symptoms, signs, imaging, EUS findings, serum IgG4, pathological findings, and extrapancreatic organ involvement, with reference to the response to glucocorticoid therapy. (Quality of evidence: Moderate; Strength of recommendation: Strong; Level of consensus: 100%)

At present, several clinical diagnostic criteria for AIP have been proposed by experts in China and abroad. The most widely used are the ICDC for AIP and the Japanese Clinical Diagnostic Criteria for AIP.^{9,16} In addition, because type 1 AIP is a form of IgG4-RD, diagnosis may also refer to the Japanese criteria and American/European classification criteria for IgG4-RD.^{66,67}

The *Chinese Guidelines for the Diagnosis and Management of Autoimmune Pancreatitis (Shanghai, 2012 Draft)* proposed practical diagnostic criteria for AIP tailored to Chinese clinical practice. Based on subsequent research advances in China and abroad, the preliminary diagnostic criteria for AIP in current clinical practice

are adjusted as follows:

1. Imaging: CT or MRI shows typical findings such as diffuse or focal enlargement of the pancreas and a capsule-like rim; MRCP shows irregular narrowing of the main pancreatic duct.
2. Laboratory tests: serum IgG4 is significantly elevated.
3. Histopathology:
 - a. the pancreas shows marked lymphocytic and plasmacytic infiltration with storiform fibrosis and obliterative phlebitis, and immunohistochemistry demonstrates >10 IgG4-positive plasma cells per high-power field;
 - b. there is abundant neutrophilic infiltration around the pancreatic ducts causing ductal epithelial injury;
 - c. EUS-guided cytology or histology excludes pancreatic and biliary malignancies.
4. Extrapancreatic organ involvement: hilar or intrahepatic bile duct strictures, enlargement of the salivary or lacrimal glands, retroperitoneal fibrosis, renal involvement, and related lesions.
5. Response to glucocorticoids: after diagnostic glucocorticoid therapy, pancreatic and/or extrapancreatic manifestations improve significantly.

A preliminary diagnosis of AIP may be made if any one of the following groups is met:

- Group A: 1 + (2 and/or 4);
- Group B: 3a (type 1 AIP) or 3b (type 2 AIP);
- Group C: (1 or 2 or 4) + 3c + 5.

In clinical practice, patients suspected of AIP should first undergo imaging examinations. If typical imaging findings are present and are supported by laboratory findings and/or extrapancreatic organ involvement, a preliminary diagnosis of AIP may be made (Group A). If only one of the following is present—typical imaging findings, laboratory findings, or extrapancreatic organ involvement—EUS-FNB/FNA is recommended to obtain tissue samples for diagnosing AIP (Group B); alternatively, after pancreatic malignancy has been excluded, diagnostic glucocorticoid therapy may be given, and a marked response supports the diagnosis of AIP (Group C). The duration of diagnostic glucocorticoid therapy should not exceed 2 weeks,⁶⁸ after which imaging should be repeated to assess improvement in pancreatic or extrapancreatic lesions. In patients who undergo surgery, a diagnosis of AIP may also be established if postoperative pathology shows typical features of AIP (Group B).

Recommendation 10. In patients with suspected AIP, differential diagnosis from pancreatic cancer and other diseases should be emphasized. (Quality of evidence: High; Strength of recommendation: Strong; Level of consensus: 100%)

AIP can easily be confused with pancreatic malignancy, and cases complicated by sclerosing cholangitis are readily misdiagnosed as cholangiocarcinoma. Differentiation between AIP and pancreaticobiliary malignancy requires integration of imaging examinations, digestive endoscopy, laboratory testing, extrapancreatic manifestations, pathology, and response to glucocorticoid therapy. The key points for differentiation are detailed in the foregoing recommendations. It should be noted that CA19-9, a tumor marker commonly used in the diagnosis of pancreatic cancer, may also be elevated in some patients with AIP.⁸ In addition, although histopathology is the gold standard for the diagnosis and differentiation of AIP and pancreatic cancer, a case report has suggested that AIP may occur as a peritumoral manifestation of pancreatic cancer,⁶⁹ and this possibility should be kept in mind in clinical

practice.

AIP should also be distinguished from chronic pancreatitis, particularly in patients with a long-standing or relapsing disease course. Imaging features favoring AIP include pancreatic enlargement, a capsule-like rim, delayed enhancement, and long or multiple ductal strictures without marked upstream dilatation, whereas chronic pancreatitis more often shows pancreatic atrophy, parenchymal or intraductal calcification, irregular ductal dilatation with side-branch ectasia, and pseudocyst formation. Given that imaging findings in certain advanced-stage AIP may overlap with those of chronic pancreatitis, the diagnosis should be based on a comprehensive clinicopathologic evaluation rather than imaging alone.

AIP is often accompanied by IgG4-SC. When pancreatic lesions are not prominent, it must also be differentiated from PSC and related conditions. In general, patients with PSC are younger, have a higher rate of coexisting IBD, respond poorly to glucocorticoid therapy, and show a “pruned-tree” appearance of the intrahepatic bile ducts or pseudodiverticulum-like changes of the bile ducts on imaging. Korean investigators proposed a scoring system to distinguish IgG4-SC from PSC, which includes three aspects: age (<30 years, 0 points; 30–40 years, 1 point; 40–50 years, 2 points; 50–60 years, 3 points; ≥60 years, 4 points), other organ involvement (present, 3 points; absent, 0 points), and beaded changes on cholangiography (present, 2 points; absent, 0 points). A total score of 0–4 suggests PSC, 5–6 suggests a diagnostic trial of glucocorticoid therapy, and 7–9 suggests IgG4-SC.⁷⁰

Recommendation 11. In patients with established type 1 AIP, dynamic assessment of disease activity by imaging and serum IgG4 during the disease course is suggested. (Quality of evidence: Low; Strength of recommendation: Weak; Level of consensus: 96.7%)

Type 1 AIP is a multiorgan disease; therefore, once the diagnosis is established, patients should undergo comprehensive assessment including clinical manifestations, extrapancreatic organ involvement, laboratory tests, and imaging examinations. Laboratory evaluation mainly includes complete blood count, urinalysis, liver and renal function, blood glucose, blood lipids, electrolytes, erythrocyte sedimentation rate, C-reactive protein, immunoglobulins, and IgG subclasses. In addition to evaluating the severity of pancreatic lesions, imaging should also assess the extent of involved extrapancreatic organs. Disease assessment may refer to the IgG4-RD Responder Index.⁷¹ This index evaluates disease status over the previous 28 days and assigns 0–3 points according to the degree of involvement in each organ, with the total score being the sum of organ-specific scores. When an involved vital organ is affected urgently and requires active treatment, the score for that organ is doubled. During treatment, patients should also be assessed regularly for disease remission and drug-related adverse effects. An IgG4-RD Responder Index ≥9 is an independent risk factor for relapse of AIP.⁷²

Treatment

Treatment of AIP should be individualized. The goals are to reduce inflammation, maintain remission, preserve organ function, and minimize treatment-related adverse effects. Treatment of AIP includes induction therapy, maintenance therapy, and management after relapse.

1. Glucocorticoid therapy

Recommendation 12. Induction therapy is suggested for patients with AIP who have symptoms or laboratory/imaging evidence of active disease. (Quality of evidence: Moderate; Strength of recommendation: Weak; Level of consensus: 100%)

AIP is an autoimmune disease, and 10–25% of patients may undergo spontaneous remission. However, active drug therapy is generally considered appropriate to prevent progression of inflammation and fibrosis and to delay or avoid irreversible organ damage. Several international guidelines are broadly consistent regarding indications for induction therapy^{9,73,74}: symptomatic patients with AIP, such as those with obstructive jaundice, abdominal pain, or back pain, should be actively considered for induction therapy; asymptomatic patients should also be treated if they have abnormal liver function or imaging evidence of active lesions. No clinical studies have supported treatment of asymptomatic AIP solely to delay pancreatic endocrine or exocrine insufficiency, and clinical evidence on the long-term prognosis of untreated asymptomatic patients remains limited.

Recommendation 13. Oral glucocorticoids are recommended as first-line induction therapy for AIP. Prednisone 30–40 mg/day, or approximately 0.6 mg·kg⁻¹·day⁻¹, may be used initially and tapered gradually after remission is achieved. (Quality of evidence: Moderate; Strength of recommendation: Strong; Level of consensus: 100%)

Oral glucocorticoids are the cornerstone of induction therapy for AIP, with the goal of rapidly controlling active disease. Both type 1 and type 2 AIP generally respond well to glucocorticoids. A Chinese systematic review including 26 studies showed that 78.4% of patients received glucocorticoid therapy, with a pooled remission rate of 96.2%.⁸ Before glucocorticoid treatment, pancreaticobiliary tumors should be carefully excluded; if the response to glucocorticoids is poor, the diagnosis should be reconsidered.

An initial dose of prednisone 30–40 mg/day or 0.6–1.0 mg·kg⁻¹·day⁻¹ is generally recommended. A nationwide Japanese survey of 1,474 patients with AIP showed that 84.4% received initial glucocorticoid therapy, most commonly prednisolone 30 mg/day (63.9%) or 40 mg/day (21.0%); when the initial dose was converted to 0.45 mg/kg, the induction remission rate remained high (202/205, 98.5%).¹¹ A Dutch study including 65 patients categorized into low-dose (prednisone 10–20 mg/day, n = 14),⁷⁵ medium-dose (30 mg/day, n = 15), and high-dose (40–60 mg/day, n = 36) groups found broadly similar baseline characteristics and, after 6 months of follow-up, similar clinical, biochemical, and imaging remission across groups; complete remission rates were 71%, 40%, and 72%, respectively. In addition, intravenous methylprednisolone (initial dose 500 mg/day for 3 days per week for 2 consecutive weeks), followed by oral prednisolone (20 mg/day) maintenance, may be more beneficial in patients with concomitant common bile duct involvement.⁷⁶ The glucocorticoid dose should be individualized. Higher doses may be considered in critically ill patients, whereas lower doses may be appropriate in elderly patients or those with mild symptoms. Tapering should also be individualized. Assessment is generally performed after 2–4 weeks of initial therapy; if clinical symptoms, laboratory findings, and

imaging signs have improved significantly, tapering may begin, usually by reducing the dose by 5 mg every 1–2 weeks.

Recommendation 14. In patients with diffuse pancreatic enlargement, slow radiologic remission after treatment, or serum IgG4 levels remaining more than twice the upper limit of normal, low-dose glucocorticoid maintenance therapy is recommended to reduce the risk of relapse. (Quality of evidence: High; Strength of recommendation: Strong; Level of consensus: 96.7%)

The 2017 International Consensus for the Treatment of Autoimmune Pancreatitis by the International Association of Pancreatology recommended low-dose glucocorticoid maintenance therapy for patients with type 1 AIP who have diffuse pancreatic enlargement,⁷⁴ delayed radiologic remission, persistently elevated serum IgG4 after treatment, two or more extrapancreatic lesions, or proximal bile duct involvement. The optimal dose and duration of maintenance therapy have not yet been established, but many Chinese and international reports have used prednisone 5–10 mg/day for 6 months to 3 years.

Multiple studies have shown that low-dose glucocorticoid maintenance therapy can reduce the risk of relapse in AIP. In a Japanese multicenter randomized controlled study including 49 patients with AIP,⁷⁷ the 3-year relapse rate was significantly lower in the maintenance group (prednisolone 5–7.5 mg/day for 3 years, $n = 30$) than in the withdrawal group (prednisolone 5–7.5 mg/day for 12 weeks, $n = 19$; 23.3% vs. 57.9%, $P = 0.011$), and relapse-free survival was significantly longer in the maintenance group ($P = 0.007$). Another Japanese study involving 22 centers and 510 patients with type 1 AIP found that the 7-year relapse rate in the prednisolone 5 mg/day maintenance group was 26.1%,⁷⁸ significantly lower than that in the withdrawal group (45.2%) and the prednisolone 2.5 mg/day maintenance group (43.4%). Compared with maintenance therapy for less than 1 year, maintenance therapy for more than 1 year was associated with a lower pooled relapse rate,⁶¹ and lack of maintenance therapy is a risk factor for relapse.⁷⁹

During glucocorticoid maintenance therapy, the patient's clinical status and disease course should be dynamically assessed, and the dose and duration should be adjusted individually, with close monitoring for possible adverse events.^{80,81}

Recommendation 15. Re-treatment with glucocorticoids is suggested for patients with relapsed AIP. (Quality of evidence: Low; Strength of recommendation: Weak; Level of consensus: 100%)

Most patients with relapse can achieve remission again with the initial glucocorticoid dose. When necessary, the glucocorticoid dose may be increased or the treatment course extended to improve disease control. A Chinese systematic review showed that among patients who relapsed after discontinuation of glucocorticoid therapy,⁸ re-treatment with glucocorticoids achieved a remission rate of 94.8% (95% CI 0.882–0.988). If patients develop significant glucocorticoid-related adverse effects, the addition of immunosuppressive agents or switching to biologic therapy is recommended.

II. Immunosuppressive agents and biologic therapy

Recommendation 16. In patients with AIP who are refrac-

tory to glucocorticoids or experience repeated relapses during tapering or after withdrawal, combination treatment with immunosuppressive agents or switching to biologic therapy is suggested. (Quality of evidence: Moderate; Strength of recommendation: Weak; Level of consensus: 100%)

If glucocorticoid monotherapy fails to control disease activity in AIP, if the disease recurs during glucocorticoid tapering, or if glucocorticoid-related adverse effects are prominent, combination therapy with immunosuppressive agents is suggested to help reduce glucocorticoid exposure.^{82,83} Commonly used immunosuppressive agents include mycophenolate mofetil, azathioprine, cyclophosphamide, and leflunomide.

A Chinese randomized controlled trial included 69 treatment-naïve patients with IgG4-RD (39 with pancreatic involvement) who received glucocorticoid monotherapy or glucocorticoids plus low-dose mycophenolate mofetil. Remission rates were not significantly different between groups, but the 12-month cumulative clinical relapse rate was lower in the combination group (11.76% vs. 34.29%).⁸² Another Chinese randomized controlled trial involving 66 patients with IgG4-RD showed that the relapse rate was significantly lower in the leflunomide plus glucocorticoid group than in the glucocorticoid monotherapy group (18.2% vs. 42.4%, $P = 0.032$).⁸⁴ However, another cohort study including 277 patients with IgG4-RD who experienced re-elevation of serum IgG4 during treatment found that although 45.1% of patients receiving glucocorticoids combined with various immunosuppressive agents (cyclophosphamide, mycophenolate mofetil, azathioprine, methotrexate, or leflunomide) showed a trend toward reduced relapse,⁸⁵ the difference was not statistically significant. During treatment with conventional immunosuppressive agents, potential adverse reactions such as leukopenia, thrombocytopenia, and liver dysfunction should be closely monitored.

Regarding biologic therapy, multiple studies have shown that rituximab is effective in patients with AIP who are resistant to or dependent on glucocorticoids.^{86–88} In patients with IgG4-RD-related pancreaticobiliary disease treated with rituximab, the complete remission rate at 6 months was 88.9% (95% CI 0.805–0.939), and the relapse rate was 21.9% (95% CI 0.105–0.403), although the relapse rate was higher in patients with multiorgan involvement (35.9%, 95% CI 0.173–0.601).⁸⁹ Before biologic therapy is initiated for AIP, patients should be fully evaluated, and adverse events should be closely monitored throughout treatment.

III. Pancreatic enzyme replacement therapy

Recommendation 17. Pancreatic enzyme replacement therapy is suggested for patients with AIP who have pancreatic exocrine insufficiency (PEI). (Quality of evidence: Low; Strength of recommendation: Weak; Level of consensus: 100%)

Patients with AIP may have PEI to varying degrees. A systematic review of 18 studies showed that the prevalence of PEI at the time of diagnosis of AIP was 45% (95% CI 0.329–0.574),⁹⁰ although this estimate showed substantial heterogeneity. A French study including 44 patients reported a PEI rate of 34% during a median follow-up of 41 (5–130) months, defined as fecal elastase-1 <200 µg/g.⁹¹ A prospective Dutch cohort study including 68 patients with AIP reported a PEI rate of 82% during a

median follow-up of 75 (50–106) months, again defined as fecal elastase-1 <200 µg/g; however, a substantial proportion of patients in that study had undergone pancreatic resection, which may have contributed to the high PEI rate.⁹² Because fecal elastase-1 is not widely used in routine practice, some studies have assessed PEI in AIP on the basis of symptoms. In a Chinese study, 4% (4/100) of patients with AIP initially presented with steatorrhea.⁷⁹ In a Korean study of 138 patients with type 1 AIP followed for a median of 60 (24–197) months, four patients (2.9%) developed marked steatorrhea, and all had relapsed during the disease course.⁹³ PEI is an important independent risk factor for mortality in chronic pancreatitis⁹⁴; therefore, monitoring and treatment of PEI should be emphasized during follow-up of patients with AIP. Treatment of PEI mainly consists of exogenous pancreatic enzyme replacement therapy, such as high-lipase pancreatic enzyme preparations with enteric coating that permits release of active enzymes in the intestine; these should be taken during meals.⁹⁵

IV. Endoscopic interventional therapy

Recommendation 18. Routine biliary drainage is not recommended for patients with AIP complicated by obstructive jaundice. (Quality of evidence: Moderate; Strength of recommendation: Strong; Level of consensus: 100%)

Some investigators have proposed that biliary drainage should be performed first in AIP patients with obstructive jaundice to reduce the risk of cholangitis induced after glucocorticoid therapy.^{9,96} However, two recent Chinese studies showed that glucocorticoid monotherapy is safe and effective in AIP patients with obstructive jaundice and that combined biliary drainage is not necessary. One study included 48 patients with AIP and obstructive jaundice⁹⁷; 25 underwent ERCP biliary stenting before glucocorticoid treatment, and 23 received glucocorticoid therapy alone. Baseline total and direct bilirubin levels were higher in the stent group, but after 4 weeks there were no significant differences in liver function indices between the two groups. In the glucocorticoid-alone group, liver function indices decreased by 53–81%, and no patient developed infection or required biliary stenting. Another study included 87 patients with type 1 AIP and marked obstructive jaundice (total bilirubin ≥ 51 µmol/L),⁹⁸ of whom 48 underwent ERCP biliary drainage before glucocorticoid therapy and 39 received glucocorticoid therapy alone. There was no significant difference in the risk of cholangitis after treatment, and after a median follow-up of 41 months, the remission and relapse rates also did not differ significantly between groups. Thus, in patients with AIP and obstructive jaundice but without acute cholangitis, glucocorticoid therapy may be initiated first, and routine biliary drainage is unnecessary.

V. Multidisciplinary diagnosis and management

Recommendation 19. Because AIP may involve multiple organs, strengthening multidisciplinary collaboration is suggested during diagnosis and management. (Quality of evidence: Low; Strength of recommendation: Weak; Level of consensus: 100%)

The clinical manifestations of AIP are complex and diverse. In particular, type 1 AIP often involves other organs, and multidisciplinary collaboration among relevant specialties is frequently required to complete diagnosis, evaluation, treatment, and follow-

up. With increasing awareness of AIP, surgery is now considered only when malignancy cannot be excluded clinically. In recent years, both the misdiagnosis rate of AIP and the rate of unnecessary surgery in China have decreased significantly.⁹⁹

Follow-up and prognosis

Recommendation 20. Long-term follow-up should be emphasized in patients with AIP, with attention to the risks of relapse, pancreatic endocrine and exocrine insufficiency, pancreatic duct stones, and malignancy. (Quality of evidence: Low; Strength of recommendation: Weak; Level of consensus: 100%)

AIP has a long disease course. A considerable proportion of patients relapse, and some develop pancreatic duct stones, pancreatic endocrine or exocrine insufficiency, and related clinical manifestations. The relationship between AIP and tumors of the pancreas or other organs remains unclear. In clinical practice, regular follow-up of patients with AIP is recommended, with attention to symptoms, imaging changes, drug-related adverse effects, and long-term outcomes.

I. Relapse

The relapse rate of AIP is relatively high, particularly in type 1 AIP. Approximately 20–40% of patients relapse after discontinuation of glucocorticoid induction therapy. A nationwide Japanese survey showed that 23.4% (344/1,471) of patients with AIP experienced relapse; 49.7% of relapses involved the pancreas, 30.5% were extrapancreatic relapses, and 19.8% involved both pancreatic and extrapancreatic sites. The cumulative relapse rates at 3, 5, and 10 years were 14%, 25%, and 40%, respectively.¹¹ Reported risk factors for relapse in type 1 AIP include extrapancreatic organ involvement (especially bile duct involvement), high pretreatment serum IgG4 levels, persistently elevated IgG4 after treatment, and diffuse pancreatic enlargement.^{61,100} Type 2 AIP has a lower relapse rate. A systematic review including 107 patients with type 2 AIP reported a post-treatment relapse rate of 15.9%.⁶¹

II. Pancreatic endocrine and exocrine insufficiency

Pancreatic endocrine and exocrine insufficiency are typical long-term outcomes of chronic pancreatitis. Endocrine insufficiency commonly manifests as diabetes mellitus. A systematic review including 62 studies and 6,522 patients with AIP showed that the prevalence of diabetes at the time of AIP diagnosis was 37% (95% CI 0.320–0.420), while the prevalence of diabetes during follow-up among patients treated with glucocorticoids was 44% (95% CI 0.261–0.624). In AIP patients who have abnormal glucose metabolism at diagnosis, blood glucose levels often worsen during glucocorticoid therapy, but pancreatic endocrine function may partially improve as the disease remits and glucocorticoids are tapered. A nationwide Japanese survey showed that 66.5% of patients with AIP had diabetes at disease onset; after glucocorticoid therapy, diabetes improved in 47.2%, whereas 17.0% developed new-onset diabetes or experienced worsening diabetes after glucocorticoid treatment. The older the patient, the higher the proportion of new-onset or worsened diabetes.¹⁰¹ In addition, although long-term glucocorticoid maintenance therapy improves serum C-peptide levels, it also increases the homeostasis model assessment index for insulin resistance.¹⁰² Therefore, whether glucocorticoid maintenance

therapy is beneficial for preserving pancreatic endocrine function remains uncertain.

PEI often manifests as dyspepsia and weight loss, and severe cases may develop steatorrhea. For discussion of PEI, see Recommendation 17.

III. Pancreatic duct stone formation

Pancreatic duct stones are a characteristic pathological feature of chronic pancreatitis. It was once thought that AIP differs from typical chronic pancreatitis and rarely leads to pancreatic duct stones or pancreatic calcification. However, with increasing research, 5–41% of patients with AIP have been reported to develop new pancreatic duct stones or an increased stone burden during follow-up.^{93,103–107} A long disease course and relapse are widely recognized risk factors for pancreatic duct stone formation.

IV. Risk of malignancy

Cases of pancreatic cancer and other malignancies occurring concurrently with or subsequent to AIP have been reported. However, because available studies are limited by sample size and study design, the association between AIP and malignancy remains inconclusive. A systematic review including 17 studies and 2,746 patients with AIP reported an overall prevalence of malignancy of 9.6% (95% CI 0.057–0.135), with prevalences of 3.7% before or concurrent with the diagnosis of AIP and 4.6% after the diagnosis of AIP. The most common malignancies were gastric cancer (1.3%, 95% CI 0.005–0.021) and colorectal cancer (1.2%, 95% CI 0.006–0.010). Thirty-five cases of pancreatic cancer were reported, but it remains uncertain whether AIP is a risk factor for these tumors or represents a paraneoplastic syndrome.¹⁰⁸ A Chinese multicenter retrospective study including 602 patients with IgG4-RD, with a median follow-up of 47.0 months, found that the overall prevalence of malignancy was significantly higher than that in the general population (standardized prevalence ratio, 8.66; 95% CI 5.84–12.31), and that AIP was an independent risk factor for malignancy in patients with IgG4-RD (odds ratio = 2.400, 95% CI 1.038–5.549, $P = 0.041$).¹⁰⁹ Therefore, attention should be paid during follow-up to the risk of pancreatic cancer and other malignancies in patients with AIP.

Compared with international guidance, including the 2011 International Consensus Diagnostic Criteria for AIP, the Japanese consensus guidelines, the European guideline on IgG4-related digestive disease, and the international treatment consensus for AIP,^{9,16,73,74} the present Chinese guidelines have several distinctive features. First, they were developed using a formal guideline-development framework, including systematic literature review, GRADE assessment, and three rounds of Delphi consensus. Second, they incorporate Chinese evidence and expert experience and adapt recommendations to clinical practice in China. Third, while referring to the framework of the ICDC, they propose simplified and more clinically practical diagnostic criteria that are better suited to routine practice. Fourth, they specifically address the management of AIP with obstructive jaundice and, based on available evidence, do not recommend routine biliary drainage in the absence of specific indications.

These guidelines reflect the evidence and expert consensus that were available during their development. Because research on AIP and IgG4-RD continues to evolve, future updates should incorporate newly published evidence through systematic reviews and formal consensus procedures. In diagnostic technologies, emerging evidence suggests that tissue acquisition with newer EUS-FNB needles may improve histologic diagnosis, while radiomics

and artificial intelligence-assisted imaging may provide additional support for differentiating AIP from pancreaticobiliary malignancies.^{48,110,111} Novel biomarkers beyond serum IgG4, including circulating plasmablasts, cytokine profiles, and disease-related autoantibodies, may also contribute to diagnosis, disease-activity assessment, and relapse prediction.^{112,113} In therapeutic aspects, B-cell-targeted strategies remain an important area of development. In addition to rituximab, newer approaches such as CD19-targeted therapy with inebilizumab and B-cell inhibition with obexelimab are under clinical investigation in IgG4-RD.^{114–116} Cellular therapies, including CD19-directed CAR-T therapy, have shown proof-of-concept activity in refractory B-cell-mediated autoimmune diseases; however, their role in IgG4-RD and AIP remains to be elucidated.¹¹⁷

Acknowledgments

We thank the Guideline and Standard Research Center of the Institute of Health Data Science, Lanzhou University, for providing methodological guidance during the development of the present guidelines.

Funding

None.

Conflict of interest

ZLiao has been an Editorial Board Member of *Journal of Translational Gastroenterology* since 2023. The other authors have no conflict of interests related to this publication.

Author contributions

Manuscript drafting (LX, YC, LH, HL, DW); supervising (ZLi, XL, XG, ZLiao). All authors have approved the final version and publication of the manuscript.

Experts participating in the discussion and finalization of the revised guidelines (listed in stroke order of surnames)

Yizhi Yu (National Key Laboratory of Immunity and Inflammation), Lin Wang (Xijing Hospital, Air Force Medical University), Chenguang Bai (The First Affiliated Hospital of Naval Medical University), Xueli Bai (The Second Affiliated Hospital, Zhejiang University School of Medicine), Yun Bian (The First Affiliated Hospital of Naval Medical University), Nonghua Lü (The First Affiliated Hospital of Nanchang University), Kexiang Zhu (The First Hospital of Lanzhou University), Xu Ren (Heilongjiang Provincial Hospital), Xu Liu (General Hospital of Northern Theater Command), Shanrong Liu (The First Affiliated Hospital of Naval Medical University), Hao Sun (The First Affiliated Hospital of Xi'an Jiaotong University), Yiqi Du (The First Affiliated Hospital of Naval Medical University), Qiyong Li (Zhejiang University International Hospital), Mingyang Li (Chinese PLA General Hospital), Peng Li (Beijing Friendship Hospital, Capital Medical University), Aiming Yang (Peking Union Medical College Hospital), Bin Xiao (The First Affiliated Hospital of Nanjing Medical University), Xi Wu (Peking Union Medical College Hospital), Duowu Zou (Ruijin Hospital, Shanghai Jiao Tong University School of Medicine), Wen Zhang (Peking Union Medical College Hos-

pital), Hong Zhang (Basic Medical College, Shaanxi University of Chinese Medicine), Xuan Zhang (Beijing Hospital), Jianping Lu (The First Affiliated Hospital of Naval Medical University), Chengwei Shao (The First Affiliated Hospital of Naval Medical University), Huaizhou Wang (The First Affiliated Hospital of Naval Medical University), Gang Jin (The First Affiliated Hospital of Naval Medical University), Zhendong Jin (The First Affiliated Hospital of Naval Medical University), Jiaxin Zhou (Peking Union Medical College Hospital), Dongbao Zhao (The First Affiliated Hospital of Naval Medical University), Bing Hu (Eastern Hepatobiliary Surgery Hospital, Naval Medical University), Yin Zhu (The First Affiliated Hospital of Nanchang University), Tong Dang (The Second Affiliated Hospital of Baotou Medical College), Huji Xu (Changzheng Hospital, Naval Medical University), Jie Gao (The First Affiliated Hospital of Naval Medical University), Xuegang Guo (Xijing Hospital, Air Force Medical University), Tao Guo (Peking Union Medical College Hospital), Caixi Tang (Zhuzhou Hospital, Xiangya School of Medicine, Central South University), Haojie Huang (The First Affiliated Hospital of Naval Medical University), Bin Cheng (Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology), Yongjing Cheng (Beijing Hospital), Yue Zeng (Shanghai General Hospital).

Methodology experts

Yaolong Chen (Institute of Health Data Science, Lanzhou University), Qianling Shi (The First Clinical Medical College of Lanzhou University).

References

- Chen YL, Yang KH, Wang XQ, Kang DY, Zhan SY, Wang JY, *et al*. Chinese guiding principles for the development/revision of clinical practice guidelines (2022 edition) (in Chinese). *Natl Med J China* 2022;102(10):697–703. doi:10.3760/cma.j.cn112137-20211228-02911.
- Brouwers MC, Kho ME, Browman GP, Burgers JS, Cluzeau F, Feder G, *et al*. AGREE II: advancing guideline development, reporting and evaluation in health care. *CMAJ* 2010;182(18):E839–E842. doi:10.1503/cmaj.090449, PMID:20603348.
- Chen Y, Yang K, Marušić A, Qaseem A, Meerpohl JJ, Flottorp S, *et al*. A Reporting Tool for Practice Guidelines in Health Care: The RIGHT Statement. *Ann Intern Med* 2017;166(2):128–132. doi:10.7326/M16-1565, PMID:27893062.
- Guyatt G, Oxman AD, Akl EA, Kunz R, Vist G, Brozek J, *et al*. GRADE guidelines: 1. Introduction–GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol* 2011;64(4):383–394. doi:10.1016/j.jclinepi.2010.04.026, PMID:21195583.
- Chari ST, Longnecker DS, Klöppel G. The diagnosis of autoimmune pancreatitis: a Western perspective. *Pancreas* 2009;38(8):846–848. doi:10.1097/MPA.0b013e3181bba281, PMID:19855232.
- López-Serrano A, Crespo J, Pascual I, Salord S, Bolado F, Del-Pozo-García AJ, *et al*. Diagnosis, treatment and long-term outcomes of autoimmune pancreatitis in Spain based on the International Consensus Diagnostic Criteria: A multi-centre study. *Pancreatol* 2016;16(3):382–390. doi:10.1016/j.pan.2016.02.006, PMID:26944001.
- Song Y, Liu QD, Zhou NX, Zhang WZ, Wang DJ. Diagnosis and management of autoimmune pancreatitis: experience from China. *World J Gastroenterol* 2008;14(4):601–606. doi:10.3748/wjg.14.601, PMID:18203294.
- Meng Q, Xin L, Liu W, Lin H, Tian B, Wang L, *et al*. Diagnosis and Treatment of Autoimmune Pancreatitis in China: A Systematic Review. *PLoS One* 2015;10(6):e0130466. doi:10.1371/journal.pone.0130466, PMID:26110658.
- Okazaki K, Kawa S, Kamisawa T, Ikeura T, Itoi T, Ito T, *et al*. Amendment of the Japanese consensus guidelines for autoimmune pancreatitis, 2020. *J Gastroenterol* 2022;57(4):225–245. doi:10.1007/s00535-022-01857-9, PMID:35192048.
- Kamisawa T, Chari ST, Giday SA, Kim MH, Chung JB, Lee KT, *et al*. Clinical profile of autoimmune pancreatitis and its histological subtypes: an international multicenter survey. *Pancreas* 2011;40(6):809–814. doi:10.1097/MPA.0b013e3182258a15, PMID:21747310.
- Masamune A, Kikuta K, Hamada S, Tsuji I, Takeyama Y, Shimosegawa T, *et al*. Nationwide epidemiological survey of autoimmune pancreatitis in Japan in 2016. *J Gastroenterol* 2020;55(4):462–470. doi:10.1007/s00535-019-01658-7, PMID:31872350.
- Xin L, He YX, Zhu XF, Zhang QH, Hu LH, Zou DW, *et al*. Diagnosis and treatment of autoimmune pancreatitis: experience with 100 patients. *Hepatobiliary Pancreat Dis Int* 2014;13(6):642–648. doi:10.1016/s1499-3872(14)60263-0, PMID:25475868.
- Li M, Bai X, Xu K, Wu X, Guo T, Jiang Q, *et al*. Peripancreatic vascular involvement in patients with type 1 autoimmune pancreatitis. *Hepatobiliary Surg Nutr* 2022;11(3):355–362. doi:10.21037/hbsn-21-82, PMID:35693390.
- Webster GJ. Autoimmune Pancreatitis - A Riddle Wrapped in an Enigma. *Dig Dis* 2016;34(5):532–539. doi:10.1159/000445234, PMID:27333053.
- Khandelwal A, Inoue D, Takahashi N. Autoimmune pancreatitis: an update. *Abdom Radiol (NY)* 2020;45(5):1359–1370. doi:10.1007/s00261-019-02275-x, PMID:31650376.
- Shimosegawa T, Chari ST, Frulloni L, Kamisawa T, Kawa S, Mino-Kenudson M, *et al*. International consensus diagnostic criteria for autoimmune pancreatitis: guidelines of the International Association of Pancreatologists. *Pancreas* 2011;40(3):352–358. doi:10.1097/MPA.0b013e3182142fd2, PMID:21412117.
- Yu ZQ, Bai XY, Ruan GC, Han W, Xu TM, Zhang MY, *et al*. Autoimmune pancreatitis associated with inflammatory bowel diseases: A retrospectively bidirectional case-control study in China. *J Dig Dis* 2023;24(8-9):452–460. doi:10.1111/1751-2980.13209, PMID:37503771.
- Ueki T, Kawamoto K, Otsuka Y, Minoda R, Maruo T, Matsumura K, *et al*. Prevalence and clinicopathological features of autoimmune pancreatitis in Japanese patients with inflammatory bowel disease. *Pancreas* 2015;44(3):434–440. doi:10.1097/MPA.0000000000000261, PMID:25469544.
- Cai O, Tan S. From Pathogenesis, Clinical Manifestation, and Diagnosis to Treatment: An Overview on Autoimmune Pancreatitis. *Gastroenterol Res Pract* 2017;2017:3246459. doi:10.1155/2017/3246459, PMID:28197205.
- Huang XM, Shi ZS, Ma CL. Multifocal autoimmune pancreatitis: A retrospective study in a single tertiary center of 26 patients with a 20-year literature review. *World J Gastroenterol* 2021;27(27):4429–4440. doi:10.3748/wjg.v27.i27.4429, PMID:34366614.
- He M, Wang X, Xu J, Li J, Chang X, Zins M, *et al*. Diffuse Involvement of Pancreas is not Always Autoimmune Pancreatitis. *Acad Radiol* 2022;29(10):1523–1531. doi:10.1016/j.acra.2022.01.013, PMID:35279380.
- Proctor RD, Rofe CJ, Bryant TJ, Hacking CN, Stedman B. Autoimmune pancreatitis: an illustrated guide to diagnosis. *Clin Radiol* 2013;68(4):422–432. doi:10.1016/j.crad.2012.08.016, PMID:23177083.
- Okazaki K, Uchida K. Autoimmune Pancreatitis: The Past, Present, and Future. *Pancreas* 2015;44(7):1006–1016. doi:10.1097/MPA.0000000000000382, PMID:26355544.
- Donet JA, Barkin JA, Keihanian T, Nemeth Z, Barkin JS. Pancreatic Pseudocysts and Parenchymal Necrosis in Patients With Autoimmune Pancreatitis: A Systematic Review. *Pancreas* 2018;47(8):952–957. doi:10.1097/MPA.0000000000001121, PMID:30028447.
- Ha J, Choi SH, Byun JH, Kim KW, Kim SY, Kim JH, *et al*. Meta-analysis of CT and MRI for differentiation of autoimmune pancreatitis from pancreatic adenocarcinoma. *Eur Radiol* 2021;31(5):3427–3438. doi:10.1007/s00330-020-07416-1, PMID:33146798.
- Zhao Y, Li F, An N, Peng Z. Atypical enhanced computed tomography signs of pancreatic cancer and its differential diagnosis from autoimmune pancreatitis. *Gland Surg* 2021;10(1):347–354. doi:10.21037/gs-20-821, PMID:33633991.
- Li J, Liu F, Fang X, Cao K, Meng Y, Zhang H, *et al*. CT Radiomics Features in Differentiation of Focal-Type Autoimmune Pancreatitis from Pancreatic Ductal Adenocarcinoma: A Propensity Score Analysis.

- Acad Radiol 2022;29(3):358–366. doi:10.1016/j.acra.2021.04.014, PMID:34108115.
- [28] He C, Rong D, Hu W, Cai Q, Yang Q, Mao Y, *et al*. A feasible CT feature to differentiate focal-type autoimmune pancreatitis from pancreatic ductal adenocarcinoma. *Cancer Med* 2019;8(14):6250–6257. doi:10.1002/cam4.2526, PMID:31469256.
- [29] Irie H, Honda H, Baba S, Kuroiwa T, Yoshimitsu K, Tajima T, *et al*. Autoimmune pancreatitis: CT and MR characteristics. *AJR Am J Roentgenol* 1998;170(5):1323–1327. doi:10.2214/ajr.170.5.9574610, PMID:9574610.
- [30] Manfredi R, Frulloni L, Mantovani W, Bonatti M, Graziani R, Pozzi Mucelli R. Autoimmune pancreatitis: pancreatic and extrapancreatic MR imaging-MR cholangiopancreatography findings at diagnosis, after steroid therapy, and at recurrence. *Radiology* 2011;260(2):428–436. doi:10.1148/radiol.11101729, PMID:21613442.
- [31] Ha J, Choi SH, Kim KW, Kim JH, Kim HJ. MRI features for differentiation of autoimmune pancreatitis from pancreatic ductal adenocarcinoma: A systematic review and meta-analysis. *Dig Liver Dis* 2022;54(7):849–856. doi:10.1016/j.dld.2021.11.013, PMID:34903501.
- [32] Shi Y, Cang L, Zhang X, Cai X, Wang X, Ji R, *et al*. The use of magnetic resonance elastography in differentiating autoimmune pancreatitis from pancreatic ductal adenocarcinoma: A preliminary study. *Eur J Radiol* 2018;108:13–20. doi:10.1016/j.ejrad.2018.09.001, PMID:30396645.
- [33] Khandelwal A, Shanbhogue AK, Takahashi N, Sandrasegaran K, Prasad SR. Recent advances in the diagnosis and management of autoimmune pancreatitis. *AJR Am J Roentgenol* 2014;202(5):1007–1021. doi:10.2214/AJR.13.11247, PMID:24758653.
- [34] Kawa S, Kamisawa T, Notohara K, Fujinaga Y, Inoue D, Koyama T, *et al*. Japanese Clinical Diagnostic Criteria for Autoimmune Pancreatitis, 2018: Revision of Japanese Clinical Diagnostic Criteria for Autoimmune Pancreatitis, 2011. *Pancreas* 2020;49(1):e13–e14. doi:10.1097/MPA.0000000000001443, PMID:31856100.
- [35] Matsubayashi H, Furukawa H, Maeda A, Matsunaga K, Kanemoto H, Uesaka K, *et al*. Usefulness of positron emission tomography in the evaluation of distribution and activity of systemic lesions associated with autoimmune pancreatitis. *Pancreatol* 2009;9(5):694–699. doi:10.1159/000199439, PMID:19684434.
- [36] Saegusa H, Momose M, Kawa S, Hamano H, Ochi Y, Takayama M, *et al*. Hilar and pancreatic gallium-67 accumulation is characteristic feature of autoimmune pancreatitis. *Pancreas* 2003;27(1):20–25. doi:10.1097/00006676-200307000-00003, PMID:12826901.
- [37] Zhang J, Jia G, Zuo C, Jia N, Wang H. (18)F-FDG PET/CT helps differentiate autoimmune pancreatitis from pancreatic cancer. *BMC Cancer* 2017;17(1):695. doi:10.1186/s12885-017-3665-y, PMID:29061130.
- [38] Ohtani M, Ofuji K, Akazawa Y, Saito Y, Nosaka T, Ozaki Y, *et al*. Clinical Usefulness of [18F]-Fluoro-2-Deoxy-D-Glucose-Positron Emission Tomography/Computed Tomography for Distinguishing Between Autoimmune Pancreatitis and Pancreatic Cancer. *Pancreas* 2021;50(7):1014–1019. doi:10.1097/MPA.0000000000001873, PMID:34629452.
- [39] Zhang Y, Cheng C, Liu Z, Wang L, Pan G, Sun G, *et al*. Radiomics analysis for the differentiation of autoimmune pancreatitis and pancreatic ductal adenocarcinoma in (18) F-FDG PET/CT. *Med Phys* 2019;46(10):4520–4530. doi:10.1002/mp.13733, PMID:31348535.
- [40] Liu Z, Li M, Zuo C, Yang Z, Yang X, Ren S, *et al*. Radiomics model of dual-time 2-[(18)F]FDG PET/CT imaging to distinguish between pancreatic ductal adenocarcinoma and autoimmune pancreatitis. *Eur Radiol* 2021;31(9):6983–6991. doi:10.1007/s00330-021-07778-0, PMID:33677645.
- [41] Palazzo M, Palazzo L, Aubert A, Fabre M, Couvelard A, Vullierme MP, *et al*. Irregular narrowing of the main pancreatic duct in association with a wall thickening is a key sign at endoscopic ultrasonography for the diagnosis of autoimmune pancreatitis. *Pancreas* 2015;44(2):211–215. doi:10.1097/MPA.0000000000000242, PMID:25394223.
- [42] Guo T, Xu T, Zhang S, Lai Y, Wu X, Wu D, *et al*. The role of EUS in diagnosing focal autoimmune pancreatitis and differentiating it from pancreatic cancer. *Endosc Ultrasound* 2021;10(4):280–287. doi:10.4103/EUS-D-20-00212, PMID:34213428.
- [43] Buscarini E, De Lisi S, Arcidiacono PG, Petrone MC, Fuini A, Conigliaro R, *et al*. Endoscopic ultrasonography findings in autoimmune pancreatitis. *World J Gastroenterol* 2011;17(16):2080–2085. doi:10.3748/wjg.v17.i16.2080, PMID:21547126.
- [44] Hocke M, Ignee A, Dietrich CF. Contrast-enhanced endoscopic ultrasound in the diagnosis of autoimmune pancreatitis. *Endoscopy* 2011;43(2):163–165. doi:10.1055/s-0030-1256022, PMID:21165827.
- [45] Kobayashi G, Fujita N, Noda Y, Ito K, Horaguchi J, Koshida S, *et al*. Vascular image in autoimmune pancreatitis by contrast-enhanced color-Doppler endoscopic ultrasonography: Comparison with pancreatic cancer. *Endosc Ultrasound* 2014;3(Suppl 1):S13. PMID:26425511.
- [46] Imazu H, Kanazawa K, Mori N, Ikeda K, Kakutani H, Sumiyama K, *et al*. Novel quantitative perfusion analysis with contrast-enhanced harmonic EUS for differentiation of autoimmune pancreatitis from pancreatic carcinoma. *Scand J Gastroenterol* 2012;47(7):853–860. doi:10.3109/00365521.2012.679686, PMID:22507131.
- [47] Zhu J, Wang L, Chu Y, Hou X, Xing L, Kong F, *et al*. A new descriptor for computer-aided diagnosis of EUS imaging to distinguish autoimmune pancreatitis from chronic pancreatitis. *Gastrointest Endosc* 2015;82(5):831–836.e1. doi:10.1016/j.gie.2015.02.043, PMID:25952089.
- [48] Marya NB, Powers PD, Chari ST, Gleeson FC, Leggett CL, Abu Dayyeh BK, *et al*. Utilisation of artificial intelligence for the development of an EUS-convolutional neural network model trained to enhance the diagnosis of autoimmune pancreatitis. *Gut* 2021;70(7):1335–1344. doi:10.1136/gutjnl-2020-322821, PMID:33028668.
- [49] Cao L, Wang Y, Wang J, Guo Q, Chen Q, Wu X, *et al*. The role of EUS-guided fine needle aspiration in autoimmune pancreatitis: a single center prospective study. *Scand J Gastroenterol* 2018;53(12):1604–1610. doi:10.1080/00365521.2018.1534137, PMID:30422724.
- [50] Yoon SB, Moon SH, Song TJ, Kim JH, Kim MH. Endoscopic ultrasound-guided fine needle aspiration versus biopsy for diagnosis of autoimmune pancreatitis: Systematic review and comparative meta-analysis. *Dig Endosc* 2021;33(7):1024–1033. doi:10.1111/den.13866, PMID:33030283.
- [51] Notohara K, Kamisawa T, Fukushima N, Furukawa T, Tajiri T, Yamaguchi H, *et al*. Guidance for diagnosing autoimmune pancreatitis with biopsy tissues. *Pathol Int* 2020;70(10):699–711. doi:10.1111/pin.12994, PMID:32767550.
- [52] Sugumar A, Levy MJ, Kamisawa T, Webster GJ, Kim MH, Enders F, *et al*. Endoscopic retrograde pancreatography criteria to diagnose autoimmune pancreatitis: an international multicentre study. *Gut* 2011;60(5):666–670. doi:10.1136/gut.2010.207951, PMID:21131631.
- [53] Kamisawa T, Nakazawa T, Tazuma S, Zen Y, Tanaka A, Ohara H, *et al*. Clinical practice guidelines for IgG4-related sclerosing cholangitis. *J Hepatobiliary Pancreat Sci* 2019;26(1):9–42. doi:10.1002/jhpb.596, PMID:30575336.
- [54] Parker AM, Schaefer D, Royer A, Sonani H, Grunes D, Glover SC. A Spy took a bite to get the diagnosis right: the benefit of SpyGlass technology in the diagnosis of autoimmune pancreatitis evading traditional diagnostic methods. *BMJ Open Gastroenterol* 2021;8(1):e000694. doi:10.1136/bmjgast-2021-000694, PMID:34479950.
- [55] Jung JG, Lee JK, Lee KH, Lee KT, Woo YS, Paik WH, *et al*. Comparison of endoscopic retrograde cholangiopancreatography with papillary biopsy and endoscopic ultrasound-guided pancreatic biopsy in the diagnosis of autoimmune pancreatitis. *Pancreatol* 2015;15(3):259–264. doi:10.1016/j.pan.2015.03.011, PMID:25891790.
- [56] Chiba K, Kamisawa T, Kuruma S, Iwasaki S, Tabata T, Koizumi S, *et al*. Major and minor duodenal papillae in autoimmune pancreatitis. *Pancreas* 2014;43(8):1299–1302. doi:10.1097/MPA.0000000000000191, PMID:25058885.
- [57] Hao M, Liu M, Fan G, Yang X, Li J. Diagnostic Value of Serum IgG4 for IgG4-Related Disease: A PRISMA-compliant Systematic Review and Meta-analysis. *Medicine (Baltimore)* 2016;95(21):e3785. doi:10.1097/MD.00000000000003785, PMID:27227950.
- [58] Culver EL, Sadler R, Simpson D, Cargill T, Makuch M, Bateman AC, *et al*. Elevated Serum IgG4 Levels in Diagnosis, Treatment Response, Organ Involvement, and Relapse in a Prospective IgG4-Related Disease UK Cohort. *Am J Gastroenterol* 2016;111(5):733–743. doi:10.1038/ajg.2016.40, PMID:27091321.
- [59] Xin L, Liao Z, Hu LH, Li ZS. The sensitivity of combined IgG4 and IgG in autoimmune pancreatitis. *Am J Gastroenterol* 2010;105(8):1902. doi:10.1038/ajg.2010.205, PMID:20686470.
- [60] Lee SC, Yang CH, Chang CT, Yu KH. Diagnostic Utility of Serum IgG4

- in Autoimmune Pancreatitis: An Updated Comprehensive Systematic Review and Meta-analysis. *J Clin Gastroenterol* 2022;56(9):810–817. doi:10.1097/MCG.0000000000001612, PMID:34516462.
- [61] Tacelli M, Celsa C, Magro B, Barresi L, Guastella S, Capurso G, *et al*. Risk Factors for Rate of Relapse and Effects of Steroid Maintenance Therapy in Patients With Autoimmune Pancreatitis: Systematic Review and Meta-analysis. *Clin Gastroenterol Hepatol* 2019;17(6):1061–1072. e8. doi:10.1016/j.cgh.2018.09.051, PMID:30312787.
 - [62] Peng Y, Li JQ, Zhang PP, Zhang X, Peng LY, Chen H, *et al*. Clinical outcomes and predictive relapse factors of IgG4-related disease following treatment: a long-term cohort study. *J Intern Med* 2019;286(5):542–552. doi:10.1111/joim.12942, PMID:31121062.
 - [63] Kamisawa T, Zen Y, Nakazawa T, Okazaki K. Advances in IgG4-related pancreatobiliary diseases. *Lancet Gastroenterol Hepatol* 2018;3(8):575–585. doi:10.1016/S2468-1253(18)30121-3, PMID:30047448.
 - [64] Fukui Y, Uchida K, Sumimoto K, Kusuda T, Miyoshi H, Koyabu M, *et al*. The similarity of Type 1 autoimmune pancreatitis to pancreatic ductal adenocarcinoma with significant IgG4-positive plasma cell infiltration. *J Gastroenterol* 2013;48(6):751–761. doi:10.1007/s00535-012-0677-x, PMID:23053421.
 - [65] Hart PA, Zen Y, Chari ST. Recent Advances in Autoimmune Pancreatitis. *Gastroenterology* 2015;149(1):39–51. doi:10.1053/j.gastro.2015.03.010, PMID:25770706.
 - [66] Umehara H, Okazaki K, Kawa S, Takahashi H, Goto H, Matsui S, *et al*. The 2020 revised comprehensive diagnostic (RCD) criteria for IgG4-RD. *Mod Rheumatol* 2021;31(3):529–533. doi:10.1080/14397595.2020.1859710, PMID:33274670.
 - [67] Wallace ZS, Naden RP, Chari S, Choi H, Della-Torre E, Dicaire JF, *et al*. The 2019 American College of Rheumatology/European League Against Rheumatism Classification Criteria for IgG4-Related Disease. *Arthritis Rheumatol* 2020;72(1):7–19. doi:10.1002/art.41120, PMID:31793250.
 - [68] Moon SH, Kim MH, Park DH, Hwang CY, Park SJ, Lee SS, *et al*. Is a 2-week steroid trial after initial negative investigation for malignancy useful in differentiating autoimmune pancreatitis from pancreatic cancer? A prospective outcome study. *Gut* 2008;57(12):1704–1712. doi:10.1136/gut.2008.150979, PMID:18583399.
 - [69] Xiang P, Zhang X, Wang C, Lang Y, Xu L, Huang L, *et al*. Pancreatic tumor in type 1 autoimmune pancreatitis: a diagnostic challenge. *BMC Cancer* 2019;19(1):814. doi:10.1186/s12885-019-6027-0, PMID:31419961.
 - [70] Moon SH, Kim MH, Lee JK, Baek S, Woo YS, Cho DH, *et al*. Development of a scoring system for differentiating IgG4-related sclerosing cholangitis from primary sclerosing cholangitis. *J Gastroenterol* 2017;52(4):483–493. doi:10.1007/s00535-016-1246-5, PMID:27470434.
 - [71] Wallace ZS, Khosroshahi A, Carruthers MD, Perugino CA, Choi H, Campochiaro C, *et al*. An International Multispecialty Validation Study of the IgG4-Related Disease Responder Index. *Arthritis Care Res (Hoboken)* 2018;70(11):1671–1678. doi:10.1002/acr.23543, PMID:29457382.
 - [72] Zongfei J, Lingli C, Ying S, Lingying M, Lijuan Z, Dongmei L, *et al*. Clinical and pathological predictors of relapse in IgG4-related disease. *Arthritis Res Ther* 2022;24(1):106. doi:10.1186/s13075-022-02792-z, PMID:35546243.
 - [73] Löhr JM, Beuers U, Vujasinovic M, Alvaro D, Frøkjær JB, Buttgerit F, *et al*. European Guideline on IgG4-related digestive disease - UEG and SGF evidence-based recommendations. *United European Gastroenterol J* 2020;8(6):637–666. doi:10.1177/2050640620934911, PMID:32552502.
 - [74] Okazaki K, Chari ST, Frulloni L, Lerch MM, Kamisawa T, Kawa S, *et al*. International consensus for the treatment of autoimmune pancreatitis. *Pancreatol* 2017;17(1):1–6. doi:10.1016/j.pan.2016.12.003, PMID:28027896.
 - [75] Buijs J, van Heerde MJ, Rauws EA, de Buy Wenniger LJ, Hansen BE, Biermann K, *et al*. Comparable efficacy of low- versus high-dose induction corticosteroid treatment in autoimmune pancreatitis. *Pancreas* 2014;43(2):261–267. doi:10.1097/MPA.0000000000000044, PMID:24518506.
 - [76] Tomiyama T, Uchida K, Matsushita M, Ikeura T, Fukui T, Takaoka M, *et al*. Comparison of steroid pulse therapy and conventional oral steroid therapy as initial treatment for autoimmune pancreatitis. *J Gastroenterol* 2011;46(5):696–704. doi:10.1007/s00535-010-0361-y, PMID:21188426.
 - [77] Masamune A, Nishimori I, Kikuta K, Tsuji I, Mizuno N, Iiyama T, *et al*. Randomised controlled trial of long-term maintenance corticosteroid therapy in patients with autoimmune pancreatitis. *Gut* 2017;66(3):487–494. doi:10.1136/gutjnl-2016-312049, PMID:27543430.
 - [78] Kubota K, Kamisawa T, Okazaki K, Kawa S, Hirano K, Hirooka Y, *et al*. Low-dose maintenance steroid treatment could reduce the relapse rate in patients with type 1 autoimmune pancreatitis: a long-term Japanese multicenter analysis of 510 patients. *J Gastroenterol* 2017;52(8):955–964. doi:10.1007/s00535-016-1302-1, PMID:28062947.
 - [79] Xin L, Meng QQ, Hu LH, Lin H, Pan J, Hao L, *et al*. Prediction and Management for Relapse of Type 1 Autoimmune Pancreatitis After Initial Steroid Treatment: A Long-Term Follow-up From China. *Pancreas* 2018;47(9):1110–1114. doi:10.1097/MPA.0000000000001147, PMID:30142120.
 - [80] Nakamura A, Ozawa M, Watanabe T, Ito T, Muraki T, Hamano H, *et al*. Predictive Factors for Autoimmune Pancreatitis Relapse After 3 Years of Maintenance Therapy. *Pancreas* 2018;47(10):1337–1343. doi:10.1097/MPA.0000000000001173, PMID:30286015.
 - [81] Hirano K, Tada M, Isayama H, Sasahira N, Umefune G, Akiyama D, *et al*. Outcome of Long-term Maintenance Steroid Therapy Cessation in Patients With Autoimmune Pancreatitis: A Prospective Study. *J Clin Gastroenterol* 2016;50(4):331–337. doi:10.1097/MCG.0000000000000440, PMID:26565969.
 - [82] Yunyun F, Yu P, Panpan Z, Xia Z, Linyi P, Jiaxin Z, *et al*. Efficacy and safety of low dose Mycophenolate mofetil treatment for immunoglobulin G4-related disease: a randomized clinical trial. *Rheumatology (Oxford)* 2019;58(1):52–60. doi:10.1093/rheumatology/key227, PMID:30124952.
 - [83] Omar D, Chen Y, Cong Y, Dong L. Glucocorticoids and steroid sparing medications monotherapies or in combination for IgG4-RD: a systematic review and network meta-analysis. *Rheumatology (Oxford)* 2020;59(4):718–726. doi:10.1093/rheumatology/kez380, PMID:31511884.
 - [84] Wang Y, Zhao Z, Gao D, Wang H, Liao S, Dong C, *et al*. Additive effect of leflunomide and glucocorticoids compared with glucocorticoids monotherapy in preventing relapse of IgG4-related disease: A randomized clinical trial. *Semin Arthritis Rheum* 2020;50(6):1513–1520. doi:10.1016/j.semarthrit.2020.01.010, PMID:32113839.
 - [85] Liu Y, Zeng Q, Zhu L, Gao J, Wang Z, Wang Z, *et al*. Relapse predictors and serologically unstable condition of IgG4-related disease: a large Chinese cohort. *Rheumatology (Oxford)* 2020;59(8):2115–2123. doi:10.1093/rheumatology/kez669, PMID:32097488.
 - [86] Nikolic S, Panic N, Hintikka ES, Dani L, Rutkowski W, Hedström A, *et al*. Efficacy and safety of rituximab in autoimmune pancreatitis type 1: our experiences and systematic review of the literature. *Scand J Gastroenterol* 2021;56(11):1355–1362. doi:10.1080/00365521.2021.1963837, PMID:34410885.
 - [87] Hart PA, Topazian MD, Witzig TE, Clain JE, Gleeson FC, Klebig RR, *et al*. Treatment of relapsing autoimmune pancreatitis with immunomodulators and rituximab: the Mayo Clinic experience. *Gut* 2013;62(11):1607–1615. doi:10.1136/gutjnl-2012-302886, PMID:22936672.
 - [88] Soliman H, Vullierme MP, Maire F, Hentic O, Ruszniewski P, Lévy P, *et al*. Risk factors and treatment of relapses in autoimmune pancreatitis: Rituximab is safe and effective. *United European Gastroenterol J* 2019;7(8):1073–1083. doi:10.1177/2050640619862459, PMID:31662864.
 - [89] Lanzillotta M, Della-Torre E, Wallace ZS, Stone JH, Karadag O, Fernández-Codina A, *et al*. Efficacy and safety of rituximab for IgG4-related pancreato-biliary disease: A systematic review and meta-analysis. *Pancreatol* 2021;21(7):1395–1401. doi:10.1016/j.pan.2021.06.009, PMID:34244040.
 - [90] Lanzillotta M, Tacelli M, Falconi M, Arcidiacono PG, Capurso G, Della-Torre E. Incidence of endocrine and exocrine insufficiency in patients with autoimmune pancreatitis at diagnosis and after treatment: a systematic review and meta-analysis. *Eur J Intern Med* 2022;100:83–93. doi:10.1016/j.ejim.2022.03.014, PMID:35367110.
 - [91] Maire F, Le Baleur Y, Rebours V, Vullierme MP, Couvelard A, Voitot H, *et al*. Outcome of patients with type 1 or 2 autoimmune pan-

- creatitis. *Am J Gastroenterol* 2011;106(1):151–156. doi:10.1038/ajg.2010.314, PMID:20736934.
- [92] Buijs J, Cahen DL, van Heerde MJ, Rauws EA, de Buy Wenniger LJ, Hansen BE, *et al*. The Long-Term Impact of Autoimmune Pancreatitis on Pancreatic Function, Quality of Life, and Life Expectancy. *Pancreas* 2015;44(7):1065–1071. doi:10.1097/MPA.0000000000000451, PMID:26355549.
- [93] Lee HW, Moon SH, Kim MH, Cho DH, Jun JH, Nam K, *et al*. Relapse rate and predictors of relapse in a large single center cohort of type 1 autoimmune pancreatitis: long-term follow-up results after steroid therapy with short-duration maintenance treatment. *J Gastroenterol* 2018;53(8):967–977. doi:10.1007/s00535-018-1434-6, PMID:29362937.
- [94] de la Iglesia-García D, Vallejo-Senra N, Iglesias-García J, López-López A, Nieto L, Domínguez-Muñoz JE. Increased Risk of Mortality Associated With Pancreatic Exocrine Insufficiency in Patients With Chronic Pancreatitis. *J Clin Gastroenterol* 2018;52(8):e63–e72. doi:10.1097/MCG.0000000000000917, PMID:28858943.
- [95] Chronic Pancreatitis Group of Pancreatic Disease Committee of Chinese Medical Doctor Association. Guideline for the diagnosis and treatment of chronic pancreatitis (2018, Guangzhou) (in Chinese). *J Clin Hepatol* 2019;35(1):45–51. doi:10.3969/j.issn.1001-5256.2019.01.008.
- [96] Okazaki K, Kawa S, Kamisawa T, Shimosegawa T, Tanaka M, Research Committee for Intractable Pancreatic Disease and Japan Pancreas Society. Japanese consensus guidelines for management of autoimmune pancreatitis: I. Concept and diagnosis of autoimmune pancreatitis. *J Gastroenterol* 2010;45(3):249–265. doi:10.1007/s00535-009-0184-x, PMID:20084528.
- [97] Feng Y, Zhang S, Guo T, Zheng W, Wu D, Wu X, *et al*. Validity and safety of corticosteroids alone without biliary stenting for obstructive jaundice in autoimmune pancreatitis. *Pancreatol* 2020;20(8):1793–1795. doi:10.1016/j.pan.2020.10.034, PMID:33097432.
- [98] Meng QQ, Gao Y, Hu LH, Qian W, Lin H, Zhang YR, *et al*. Limited Role of Endoscopic Biliary Drainage Before Steroid Treatment for Autoimmune Pancreatitis With Significant Obstructive Jaundice. *Pancreas* 2021;50(8):e65–e66. doi:10.1097/MPA.0000000000001883, PMID:34714290.
- [99] Wu L, Li W, Huang X, Wang Z. Clinical features and comprehensive diagnosis of autoimmune pancreatitis in China. *Digestion* 2013;88(2):128–134. doi:10.1159/000353597, PMID:24008239.
- [100] Lu H, Peng L, Zhu L, Feng R, Zhou J, Liu Z, *et al*. Persistent enlargement of the pancreatic gland after glucocorticoid therapy increases the risk of relapse in IgG4-related autoimmune pancreatitis. *Clin Rheumatol* 2022;41(6):1709–1718. doi:10.1007/s10067-022-06091-5, PMID:35175447.
- [101] Nishimori I, Tamakoshi A, Kawa S, Tanaka S, Takeuchi K, Kamisawa T, *et al*. Influence of steroid therapy on the course of diabetes mellitus in patients with autoimmune pancreatitis: findings from a nationwide survey in Japan. *Pancreas* 2006;32(3):244–248. doi:10.1097/01.mpa.0000202950.02988.07, PMID:16628078.
- [102] Noguchi K, Nakai Y, Mizuno S, Isayama H, Hirano K, Kanai S, *et al*. Insulin secretion improvement during steroid therapy for autoimmune pancreatitis according to the onset of diabetes mellitus. *J Gastroenterol* 2020;55(2):198–204. doi:10.1007/s00535-019-01615-4, PMID:31471733.
- [103] Hirano K, Tada M, Isayama H, Watanabe T, Saito T, Uchino R, *et al*. High alcohol consumption increases the risk of pancreatic stone formation and pancreatic atrophy in autoimmune pancreatitis. *Pancreas* 2013;42(3):502–505. doi:10.1097/MPA.0b013e31826b3984, PMID:23146923.
- [104] Ito T, Kawa S, Matsumoto A, Kubota K, Kamisawa T, Okazaki K, *et al*. Risk Factors for Pancreatic Stone Formation in Type 1 Autoimmune Pancreatitis: A Long-term Japanese Multicenter Analysis of 624 Patients. *Pancreas* 2019;48(1):49–54. doi:10.1097/MPA.0000000000001210, PMID:30540679.
- [105] Maruyama M, Arakura N, Ozaki Y, Watanabe T, Ito T, Yoneda S, *et al*. Risk factors for pancreatic stone formation in autoimmune pancreatitis over a long-term course. *J Gastroenterol* 2012;47(5):553–560. doi:10.1007/s00535-011-0510-y, PMID:22183858.
- [106] Kuraishi Y, Uehara T, Watanabe T, Ashihara N, Ozawa M, Kanai K, *et al*. Corticosteroids prevent the progression of autoimmune pancreatitis to chronic pancreatitis. *Pancreatol* 2020;20(6):1062–1068. doi:10.1016/j.pan.2020.07.408, PMID:32800575.
- [107] Hart PA, Kamisawa T, Brugge WR, Chung JB, Culver EL, Czako L, *et al*. Long-term outcomes of autoimmune pancreatitis: a multicentre, international analysis. *Gut* 2013;62(12):1771–1776. doi:10.1136/gutjnl-2012-303617, PMID:23232048.
- [108] Haghbin H, Chuang J, Fatima R, Zakirkhodjaev N, Lee-Smith W, Aziz M. Correlation of Autoimmune Pancreatitis and Malignancy: Systematic Review and Meta-Analysis. *Dig Dis Sci* 2022;67(7):3252–3264. doi:10.1007/s10620-021-07179-9, PMID:34297267.
- [109] Liu Y, Fu J, Ning X, Li H, Ma X, Wang K, *et al*. Malignancy Risk of Immunoglobulin G4-Related Disease: Evidence from a Large Cohort Multicenter Retrospective Study. *Rheumatol Ther* 2021;8(3):1207–1221. doi:10.1007/s40744-021-00326-8, PMID:34184190.
- [110] Ishikawa T, Kawashima H, Ohno E, Suhara H, Hayashi D, Hiramatsu T, *et al*. Usefulness of endoscopic ultrasound-guided fine-needle biopsy for the diagnosis of autoimmune pancreatitis using a 22-gauge Franseen needle: a prospective multicenter study. *Endoscopy* 2020;52(11):978–985. doi:10.1055/a-1183-3583, PMID:32583394.
- [111] Zhong J, Hu Y, Xing Y, Ge X, Ding D, Zhang H, *et al*. A systematic review of radiomics in pancreatitis: applying the evidence level rating tool for promoting clinical transferability. *Insights Imaging* 2022;13(1):139. doi:10.1186/s13244-022-01279-4, PMID:35986798.
- [112] Hara A, Watanabe T, Minaga K, Yoshikawa T, Kamata K, Kudo M. Biomarkers in autoimmune pancreatitis and immunoglobulin G4-related disease. *World J Gastroenterol* 2021;27(19):2257–2269. doi:10.3748/wjg.v27.i19.2257, PMID:34040320.
- [113] Jarrell JA, Baker MC, Perugino CA, Liu H, Bloom MS, Maehara T, *et al*. Neutralizing anti-IL-1 receptor antagonist autoantibodies induce inflammatory and fibrotic mediators in IgG4-related disease. *J Allergy Clin Immunol* 2022;149(1):358–368. doi:10.1016/j.jaci.2021.05.002, PMID:33974929.
- [114] Lanzillotta M, Stone JH, Della-Torre E. B-Cell depletion therapy in IgG4-related disease: State of the art and future perspectives. *Mod Rheumatol* 2023;33(2):258–265. doi:10.1093/mr/roac098, PMID:35983918.
- [115] Perugino C, Culver EL, Khosroshahi A, Zhang W, Della-Torre E, Okazaki K, *et al*. Efficacy and Safety of Inebilizumab in IgG4-Related Disease: Protocol for a Randomized Controlled Trial. *Rheumatol Ther* 2023;10(6):1795–1808. doi:10.1007/s40744-023-00593-7, PMID:37792260.
- [116] Perugino CA, Wallace ZS, Zack DJ, Quinn SM, Poma A, Fernandes AD, *et al*. Evaluation of the safety, efficacy, and mechanism of action of obexelimab for the treatment of patients with IgG4-related disease: an open-label, single-arm, single centre, phase 2 pilot trial. *Lancet Rheumatol* 2023;5(8):e442–e450. doi:10.1016/S2665-9913(23)00157-1, PMID:38251576.
- [117] Mackensen A, Müller F, Mougiakakos D, Böltz S, Wilhelm A, Aigner M, *et al*. Anti-CD19 CAR T cell therapy for refractory systemic lupus erythematosus. *Nat Med* 2022;28(10):2124–2132. doi:10.1038/s41591-022-02017-5, PMID:36109639.