



Case Report

Pediatric Atypical Teratoid/Rhabdoid Tumor of the Cerebellum: A Case Report and Literature Review



Xing Huang, Yike Cai and Hong Lin*

Department of Neurosurgery, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei, China

Received: January 15, 2025 | Revised: April 07, 2025 | Accepted: April 29, 2025 | Published online: June 30, 2025

Abstract

Atypical teratoid/rhabdoid tumor (AT/RT) is a rare and highly aggressive embryonal tumor that predominantly affects infants and young children. This malignancy arises from primitive neuroectodermal cells and exhibits heterogeneous differentiation into various embryonic tissues. Due to its rarity and complexity, diagnosing and managing AT/RT present significant challenges. Recent studies have summarized the key features of cerebellar and supratentorial AT/RT cases; however, critical gaps remain in understanding their diffuse leptomeningeal variants and long-term functional outcomes. Here, we report a case of a two-year-old child diagnosed with cerebellar AT/RT, who presented with vomiting and gait instability. The patient underwent a gross total resection followed by adjuvant radiotherapy and chemotherapy. Despite achieving radiological remission, the patient survived for only eight months and experienced severe neurological deficits, including persistent ataxia and recurrent infections. This case highlights the disconnect between surgical success and long-term quality of life. It underscores the importance of integrating molecular diagnostics and palliative care to address the multifaceted burden of AT/RT.

Introduction

Atypical teratoid/rhabdoid tumor (AT/RT) is one of the most aggressive central nervous system (CNS) malignancies in pediatric patients, accounting for approximately 1–2% of all pediatric brain tumors.¹ It predominantly affects children under the age of three, with a slight male predominance.² AT/RT is characterized by rapid progression, resistance to conventional therapies, and poor prognosis, with a median survival of less than one year despite aggressive treatment.³ The tumor typically arises in the posterior fossa, particularly in the cerebellum, but can also occur in supratentorial regions.⁴ Histologically, AT/RT is composed of undifferentiated small round cells with rhabdoid features and often shows heterogeneous differentiation into mesenchymal, epithelial, and neural tissues.⁵ Immunohistochemically, the tumor frequently expresses vimentin and other intermediate filaments, reflecting its neural crest origin.⁵

The molecular pathogenesis of AT/RT involves the inactivation of the SMARCB1 (INI1) or SMARCA4 (BRG1) genes, which are critical components of the SWI/SNF chromatin remodeling com-

plex.⁶ These genetic alterations lead to dysregulation of cell cycle control and differentiation, contributing to the tumor's aggressive behavior.⁷ Despite advances in molecular diagnostics and multimodal therapies, including surgery, radiation, and chemotherapy, the prognosis for AT/RT remains dismal.⁷ This case report aimed to provide a detailed account of the clinical presentation, diagnostic challenges, surgical management, and postoperative outcomes in a two-year-old girl with cerebellar AT/RT, contributing to the limited literature on this rare and devastating disease.

Case presentation

A two-year-old girl presented to Wuhan Union Hospital in November 2023 with a one-week history of frequent vomiting and gait instability. Brain magnetic resonance imaging (MRI) revealed a large mass in the fourth ventricle, accompanied by supratentorial hydrocephalus (Fig. 1). On physical examination, the patient was confused, with bilaterally equal and round pupils measuring 3 mm in diameter and reactive to light. Muscle strength in all four limbs was graded as 3. Following admission, comprehensive evaluations were completed, and no contraindications to surgery were identified. A posterior midline approach was employed to resect the cerebellar lesion. Intraoperatively, the tumor appeared slightly firm and fleshy, with moderate vascularity. Portions of the tumor were poorly demarcated from the inferior vermis of the cerebellum, while anteriorly it compressed the brainstem with clearly defined margins. Superiorly, it obstructed the aqueduct of Sylvius. After a three-hour procedure, gross total resection of the

Keywords: Atypical teratoid/rhabdoid tumor; Clinical manifestation; Surgical treatment; Case report.

*Correspondence to: Hong Lin, Department of Neurosurgery, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei 430022, China. ORCID: <https://orcid.org/0000-0003-1046-0325>. Tel: +86-027-85350775, E-mail: lh6309@126.com

How to cite this article: Huang X, Cai Y, Lin H. Pediatric Atypical Teratoid/Rhabdoid Tumor of the Cerebellum: A Case Report and Literature Review. *Neurosurg Subspec* 2025;000(000):000–000. doi: 10.14218/NSSS.2025.00002.




Fig. 1. Preoperative imaging findings of the atypical teratoid/rhabdoid tumor. (a) T1-enhanced presentation of tumors in the sagittal plane. (b) T2 manifestation of tumors in the sagittal plane. (c) T1 manifestation of tumors in the horizontal plane. (d–g, i) Imaging manifestations of tumors on continuous T1-enhanced scans. (h) Localized magnification indicates the unclear boundary between the tumor and the cerebellum (The red arrow indicates the tumor).

tumor was achieved.

Postoperative care and rehabilitation

Following surgical resection, the patient was closely monitored in the intensive care unit for 24 h to manage potential complications, such as intracranial pressure fluctuations and cerebrospinal fluid leakage. Postoperative imaging confirmed complete tumor removal (Fig. 2), and the patient was transferred to the pediatric

ward for continued recovery. Over the subsequent two weeks, the patient underwent physical therapy to address gait instability and muscle weakness. Despite these efforts, the patient continued to experience significant neurological deficits, including persistent ataxia and impaired fine motor function. Psychological support was provided to the family to assist in coping with the emotional and physical challenges posed by the illness. The patient was discharged after three weeks with a plan for outpatient rehabilitation




Fig. 2. Postoperative pathological and imaging manifestations of atypical teratoid/rhabdoid tumors. (a–f) The MRI in the sagittal and horizontal planes shows that the tumor has been completely removed. (g, h) The tumor did not metastasize to the distant spinal cord. (i, j) Postoperative pathological manifestations (200×). MRI, magnetic resonance imaging.

and scheduled follow-up visits.

Prognosis and survival

The patient was referred to the oncology department for adjuvant radiotherapy and chemotherapy, receiving a regimen that included vincristine and carboplatin. Despite aggressive multimodal therapy, the patient's quality of life remained poor, with persistent neurological deficits and limited functional recovery. The patient survived for eight months post-surgery but required frequent hospital readmissions due to complications, including infections and progressive neurological decline. This outcome highlights the aggressive nature of AT/RT and the limited effectiveness of current

treatment modalities in improving long-term survival and quality of life.

Discussion

AT/RT represents one of the most challenging pediatric CNS malignancies due to its aggressive biology, resistance to conventional therapies, and dismal prognosis (median survival <12 months).⁷ While it predominantly affects children under three years of age, its rarity (1–2% of pediatric brain tumors) further complicates standardized management.² To contextualize our findings within the existing literature, we systematically reviewed recent AT/RT

Table 1. Summary of similar case reports of atypical teratoid/rhabdoid tumor (AT/RT) in pediatric patients

Study	Age/Sex	Location	Symptoms	Treatment	Outcome	Key findings
Guo <i>et al.</i> , 2022 ⁸	27 months/M	Cerebellum	Vomiting, ataxia	Subtotal resection, chemotherapy (vincristine, carboplatin)	Survived nine months, poor quality of life	Tumor recurrence leads to a poor prognosis
Shah <i>et al.</i> , 2023 ⁹	10 months/F	Sellar	Polyuria, polydipsia, drowsiness	Biopsy surgery	Abandoning treatment	Early identification and timely multidisciplinary management are crucial
El Malih <i>et al.</i> , 2023 ¹⁰	1 month/M	Cerebellum	Eye malformation	chemotherapy (vincristine and cyclophosphamide)	Survived one week	The importance of radiological imaging in evaluating this rare disease
Fujigasaki <i>et al.</i> , 2012 ¹¹	6 months/F	Fourth ventricle	Vomiting	Total resection, radiotherapy	Survived 26 years	Highlighted the importance of complete tumor resection
Syed <i>et al.</i> , 2023 ¹²	15 months/F	Lumbar vertebra	Fussiness	Subtotal resection, chemotherapy, radiotherapy	Survived two years, no recurrence	Aggressive adjuvant therapy can allow patients to achieve remission
Sharma <i>et al.</i> , 2020 ¹³	4 years/M	Lateral ventricle	Vomiting, headache	Gross total resection	Survived three months, recurrence at two months	AT/RT can have an aggressive course with a tendency for early leptomeningeal spread

F, female; M, male.

case reports focusing on cerebellar presentations.

We conducted a systematic literature search of the PubMed and Web of Science databases for case reports published between 2000 and 2023. The search terms included “atypical teratoid/rhabdoid tumor”, “AT/RT”, “pediatric”, and “cerebellum”. A total of 15 relevant case reports were identified, and 6 representative cases involving children under three years old were selected for inclusion based on their relevance to cerebellar AT/RT and the availability of detailed clinical data.

As summarized in Table 1, AT/RT is among the most malignant forms of childhood CNS neoplasms.^{8–13} The selected reports provide a comprehensive overview of the clinical features, treatment strategies, and outcomes of pediatric AT/RT. They highlight the challenges in managing this aggressive tumor and underscore the urgent need for further research into targeted therapies and personalized treatment approaches.

Patients with AT/RT usually present within the first three years of life, with a slight male predominance.² In our case, the patient was a two-year-old girl, aligning with the typical age of onset but contrasting with the reported male predilection. Symptoms commonly include signs of increased intracranial pressure, such as vomiting, headaches, and papilledema, along with focal neurological deficits like seizures and motor or sensory impairments.¹⁴ Our patient exhibited frequent vomiting and gait instability, consistent with cerebellar dysfunction due to the tumor’s location in the fourth ventricle. Radiological imaging typically shows a heterogeneous mass with mixed density on computed tomography and variable signal intensity on T1-weighted and T2-weighted MRI images.⁵ Post-contrast enhancement often reveals irregular patterns of tumor vascularity. In this case, preoperative MRI revealed a well-circumscribed cerebellar mass with mixed T1/T2 signals and post-contrast enhancement (Fig. 1a–c), consistent with classical imaging features.

Definitive diagnosis relies on surgical resection and histopathological examination. Distinguishing AT/RT from other pediatric brain tumors, particularly medulloblastoma and primitive neuroectodermal tumors, requires careful evaluation of immunohistochemical markers.¹⁵ Genetic studies have identified specific alterations in AT/RT, with SMARCB1 deletion being a key event in its pathogenesis,⁶ providing potential therapeutic targets.

Given the aggressive nature of AT/RT, multimodal therapy is recommended.¹⁶ Our patient underwent a gross total resection followed by adjuvant radiotherapy and chemotherapy. Despite achieving radiological remission (Fig. 2a–f), survival was limited to eight months, with severe neurological deficits and recurrent infections. Adjuvant radiotherapy targets residual disease and has demonstrated some efficacy in delaying progression.¹⁷ Chemotherapy regimens, including high-dose methotrexate, vincristine, and carboplatin, are commonly employed, but response rates remain suboptimal.⁷ Targeted therapies, including enhancer of zeste homolog 2 inhibitors (e.g., UNC1999), mesenchymal-epithelial transition factor inhibitors, and immune checkpoint inhibitors, are currently under investigation in clinical trials and offer hope for improved survival outcomes.⁶

Despite aggressive treatment, our patient’s poor quality of life and limited survival underscore the grim prognosis associated with AT/RT. The tumor’s rapid progression and resistance to conventional therapies continue to pose significant therapeutic challenges.⁶ A deeper understanding of the molecular mechanisms underlying this disease is crucial for the development of more effective treatments.

Conclusions

Conclusions

This case highlights the challenges associated with managing AT/RT, particularly in pediatric patients. Complete surgical resection remains the cornerstone of AT/RT treatment, as demonstrated in this case, where total tumor removal was achieved. However, despite successful surgery and multimodal therapy, the prognosis for AT/RT remains poor, underscoring the urgent need for more effective treatment strategies. Postoperative care and rehabilitation are

crucial for optimizing patient recovery and quality of life. Collaborative efforts among clinicians, pathologists, and researchers are essential to advancing our understanding of this devastating disease and to the development of innovative therapeutic approaches.

Acknowledgments

None.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflict of interest

The authors declare that they have no known competing financial interests or personal relationships that could have influenced the work reported in this paper.

Author contributions

Conceptualization, methodology, investigation of this study (XH, HL), data curation, formal analysis, writing the original draft (XH), validation, resources, and supervision (YKC). All authors participated in the review and editing of the manuscript and approved the final version for publication.

Ethical statement

This study was conducted in accordance with the ethical standards of Huazhong University of Science and Technology Tongji Medical College Medical Ethics Committee (TJ-IRB2025068), as well as the 2024 Helsinki Declaration and its later amendments. Informed consent was obtained from the patient's legal guardian for the publication of this case report and any accompanying images.

References

- [1] Li D, Heiferman DM, Syed HR, Santos JG, Bowman RM, DiPatri AJ, *et al*. Pediatric primary spinal atypical teratoid rhabdoid tumor: a case series and review of the literature. *J Neurosurg Pediatr* 2019;24(3):267–283. doi:10.3171/2019.4.PEDS19113, PMID:31299639.
- [2] Park M, Han JW, Hahn SM, Lee JA, Kim JY, Shin SH, *et al*. Atypical Teratoid/Rhabdoid Tumor of the Central Nervous System in Children under the Age of 3 Years. *Cancer Res Treat* 2021;53(2):378–388. doi:10.4143/crt.2020.756, PMID:33138347.
- [3] Wu WW, Bi WL, Kang YJ, Ramkissoon SH, Prasad S, Shih HA, *et al*. Adult Atypical Teratoid/Rhabdoid Tumors. *World Neurosurg* 2016;85:197–204. doi:10.1016/j.wneu.2015.08.076, PMID:26344637.
- [4] Corvino S, Del Basso De Caro M, Franca RA, Corazzelli G, Della Monica R, Chiariotti L, *et al*. Atypical Teratoid/Rhabdoid Tumor of the Nervous System in Adults: Location-Related Features and Outcome. *World Neurosurg* 2023;179:e404–e415. doi:10.1016/j.wneu.2023.08.107, PMID:37659753.
- [5] Zhao RJ, Wu KY, Zhang JG, Ma YH, Kong LF. Primary Intracranial Atypical Teratoid/Rhabdoid Tumors: A Clinicopathologic and Neuroradiologic Study. *J Child Neurol* 2015;30(8):1017–1023. doi:10.1177/0883073814551795, PMID:25323739.
- [6] Nesvick CL, Lafay-Cousin L, Raghunathan A, Bouffet E, Huang AA, Daniels DJ. Atypical teratoid rhabdoid tumor: molecular insights and translation to novel therapeutics. *J Neurooncol* 2020;150(1):47–56. doi:10.1007/s11060-020-03639-w, PMID:33021733.
- [7] Frühwald MC, Biegel JA, Bourdeaut F, Roberts CW, Chi SN. Atypical teratoid/rhabdoid tumors-current concepts, advances in biology, and potential future therapies. *Neuro Oncol* 2016;18(6):764–778. doi:10.1093/neuonc/nov264, PMID:26755072.
- [8] Guo G, Zhuang J, Zhang K, Zhou Z, Wang Y, Zhang Z. Atypical Teratoid/Rhabdoid Tumor of the Central Nervous System in Children: Case Reports and Literature Review. *Front Surg* 2022;9:864518. doi:10.3389/fsurg.2022.864518, PMID:36034392.
- [9] Shah R, Ambalavanan N, Chatterjee D, Dutta P. Sellar Suprasellar Surprise: A Rare Case of Atypical Teratoid/Rhabdoid Tumor in an Infant. *Cureus* 2023;15(10):e47632. doi:10.7759/cureus.47632, PMID:38022150.
- [10] El Malih S, Akammar A, El Mounssefe G, Tahiri L, El Bouardi N, Haloua M, *et al*. Synchronous congenital malignant rhabdoid tumor of the orbit and cerebellar atypical teratoid/rhabdoid tumor: Case report. *Radiol Case Rep* 2023;18(11):3979–3983. doi:10.1016/j.radcr.2023.07.074, PMID:37680666.
- [11] Takahashi-Fujigasaki J, Matumoto M, Kan I, Oka H, Yasue M. Atypical teratoid/rhabdoid tumor with 26-year overall survival: case report. *J Neurosurg Pediatr* 2012;9(4):400–405. doi:10.3171/2012.1.PEDS11350, PMID:22462705.
- [12] Syed H, Teferi N, Hanson A, Challa M, Eschbacher K, Hitchon P. Clinical diagnostic and radiographic features of primary spinal atypical teratoid rhabdoid tumors tumor in a pediatric patient: A case report and review of the literature. *J Cent Nerv Syst Dis* 2023;15:11795735231209199. doi:10.1177/11795735231209199, PMID:37876767.
- [13] Sharma A, Sable MN, Singla R, Dash C, Sahu RN. Lateral Ventricle Atypical Teratoid/Rhabdoid Tumor (AT/RT): Case Report and Review of Literature. *Neurol India* 2020;68(6):1465–1468. doi:10.4103/0028-3886.304127, PMID:33342894.
- [14] Greenaway GP, Page PS, Patel V, Ahmed AS. Atypical Teratoid/Rhabdoid Tumor of the Cerebellum in an Adult: Case Report and Literature Review. *World Neurosurg* 2021;145:57–63. doi:10.1016/j.wneu.2020.08.079, PMID:32810633.
- [15] Calandrelli R, Massimi L, Pilato F, Verdolotti T, Ruggiero A, Attinà G, *et al*. Atypical Teratoid Rhabdoid Tumor: Proposal of a Diagnostic Pathway Based on Clinical Features and Neuroimaging Findings. *Diagnostics (Basel)* 2023;13(3):475. doi:10.3390/diagnostics13030475, PMID:36766580.
- [16] Schrey D, Carceller Lechón F, Malietzis G, Moreno L, Dufour C, Chi S, *et al*. Multimodal therapy in children and adolescents with newly diagnosed atypical teratoid rhabdoid tumor: individual pooled data analysis and review of the literature. *J Neurooncol* 2016;126(1):81–90. doi:10.1007/s11060-015-1904-0, PMID:26608522.
- [17] Yang WC, Yen HJ, Liang ML, Chen HH, Lee YY, Wong TT, *et al*. Role of early and aggressive post-operative radiation therapy in improving outcome for pediatric central nervous system atypical teratoid/rhabdoid tumor. *Childs Nerv Syst* 2019;35(6):1013–1020. doi:10.1007/s00381-019-04126-y, PMID:30982172.